



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 03:39 AM EDT

PDB ID : 1MW9  
Title : Crystal Structure of H365R mutant of 67 kDA N-terminal fragment of E. coli DNA Topoisomerase I  
Authors : Perry, K.; Mondragon, A.  
Deposited on : 2002-09-27  
Resolution : 1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	592	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA Topoisomerase I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	X	556	4643	2912	827	885	19	0	27	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	365	ARG	HIS	engineered mutation	UNP P06612
X	549	ASN	SER	SEE REMARK 999	UNP P06612

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		
2	X	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	X	577	Total	O	0	0
			577	577		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.99Å 78.70Å 139.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.39 – 1.67 49.18 – 1.67	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.39-1.67) 98.2 (49.18-1.67)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 1.67Å)	Xtrriage
Refinement program	REFMAC 5.0, CNS	Depositor
R, $R_{free}$	0.208 , 0.216 0.215 , 0.221	Depositor DCC
$R_{free}$ test set	7623 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.9	Xtrriage
Anisotropy	0.366	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 47.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5265	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	X	0.27	0/4734	0.64	4/6392 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	36	ASP	CB-CG-OD2	5.33	123.10	118.30
1	X	243	ASP	CB-CG-OD2	5.22	123.00	118.30
1	X	581	ASP	CB-CG-OD2	5.15	122.94	118.30
1	X	386	ASP	CB-CG-OD2	5.14	122.93	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	4643	0	4580	117	0
2	X	45	0	0	0	0
3	X	577	0	0	18	0
All	All	5265	0	4580	117	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (117) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:279[B]:LYS:HD2	1:X:280:PRO:HD2	1.35	1.08
1:X:521:LYS:HE3	1:X:521:LYS:N	1.83	0.93
1:X:35:ARG:HE	1:X:122:HIS:HD2	1.26	0.84
1:X:539:ASN:HD22	1:X:542:PHE:H	1.25	0.83
1:X:428:ILE:HD11	1:X:449:ILE:HG23	1.64	0.80
1:X:258:VAL:O	1:X:262:GLU:HG3	1.85	0.76
1:X:381:LYS:HB2	1:X:381:LYS:HZ3	1.53	0.74
1:X:250:ASN:HD21	1:X:253:GLN:HG3	1.51	0.74
1:X:432:ASP:OD2	1:X:436:LYS:HE3	1.88	0.74
1:X:220:TRP:HB2	1:X:248[B]:PRO:HG2	1.69	0.73
1:X:138:VAL:HG12	1:X:365:ARG:NH2	2.04	0.72
1:X:225:SER:HB2	1:X:463[B]:GLU:HG3	1.74	0.69
1:X:250:ASN:ND2	1:X:253:GLN:H	1.91	0.68
1:X:138:VAL:HG12	1:X:365:ARG:HH22	1.59	0.68
1:X:7[B]:ILE:HD13	1:X:123:LEU:HD13	1.75	0.67
1:X:436:LYS:NZ	3:X:1544:HOH:O	2.29	0.66
1:X:511:ARG:HD3	1:X:520[A]:GLU:OE1	1.96	0.66
1:X:295[B]:THR:HG22	3:X:1165:HOH:O	1.96	0.65
1:X:422:LEU:C	1:X:423:LYS:HD2	2.17	0.65
1:X:7[B]:ILE:HD11	1:X:30:SER:HB3	1.79	0.64
1:X:34:ILE:HD11	1:X:93:VAL:HG22	1.77	0.64
1:X:35:ARG:HE	1:X:122:HIS:CD2	2.12	0.64
1:X:97:LYS:O	1:X:101:GLU:HG2	2.01	0.61
1:X:202:ARG:O	1:X:206:GLU:HG3	2.01	0.61
1:X:310:ARG:NH2	1:X:384[B]:GLU:OE2	2.33	0.61
1:X:168:ARG:HA	1:X:171:MET:CE	2.32	0.60
1:X:220:TRP:CZ2	1:X:251:LYS:HG3	2.37	0.60
1:X:422:LEU:O	1:X:423:LYS:HD2	2.04	0.58
1:X:583:GLU:HB2	1:X:584:GLU:OE1	2.04	0.57
1:X:521:LYS:HE2	3:X:1575:HOH:O	2.04	0.57
1:X:427:ARG:HE	1:X:447:ASP:CG	2.08	0.56
1:X:577:LYS:HA	1:X:580:LYS:HD3	1.88	0.56
1:X:250:ASN:C	1:X:250:ASN:HD22	2.08	0.55
1:X:340[B]:SER:OG	1:X:348:LEU:HD22	2.06	0.55
1:X:582[B]:PRO:HG2	1:X:583:GLU:OE2	2.07	0.54
1:X:236:LEU:HD23	1:X:424:ALA:HB2	1.88	0.54
1:X:250:ASN:ND2	1:X:253:GLN:HG3	2.22	0.53
1:X:436:LYS:HE2	3:X:1209:HOH:O	2.08	0.53
1:X:167:ALA:O	1:X:171:MET:HG3	2.10	0.52
1:X:168:ARG:HA	1:X:171:MET:HE2	1.91	0.52

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:263:LYS:NZ	3:X:1299:HOH:O	2.41	0.52
1:X:271:ARG:HD2	1:X:453:VAL:O	2.10	0.52
1:X:441:LEU:H	1:X:441:LEU:HD12	1.74	0.52
1:X:421:ARG:CD	1:X:423:LYS:HE3	2.41	0.51
1:X:112:LEU:HD22	1:X:365:ARG:NH1	2.26	0.51
1:X:239:THR:CG2	1:X:423:LYS:HD3	2.40	0.51
1:X:279[A]:LYS:N	1:X:279[A]:LYS:HD2	2.26	0.51
1:X:63:GLU:HB2	3:X:1328:HOH:O	2.10	0.51
1:X:278[B]:SER:OG	1:X:406:ALA:HB3	2.10	0.50
1:X:392[A]:GLN:HE21	1:X:396[A]:ARG:HG3	1.76	0.50
1:X:28:LYS:HE3	3:X:1184:HOH:O	2.11	0.50
1:X:239:THR:HG21	1:X:423:LYS:HD3	1.93	0.50
1:X:250:ASN:HD21	1:X:253:GLN:H	1.56	0.49
1:X:354:GLN:HE22	1:X:356:ALA:HB3	1.77	0.49
1:X:34:ILE:HD11	1:X:93:VAL:CG2	2.40	0.49
1:X:513:GLU:HG2	1:X:518:TYR:CE1	2.47	0.49
1:X:166[B]:GLN:NE2	3:X:1040:HOH:O	2.38	0.49
1:X:225:SER:CB	1:X:463[B]:GLU:HG3	2.42	0.49
1:X:233:ALA:O	1:X:451[A]:PRO:HG2	2.12	0.49
1:X:346:LYS:HG2	3:X:1292:HOH:O	2.11	0.49
1:X:421:ARG:HD2	1:X:423:LYS:HE3	1.93	0.49
1:X:91:LYS:NZ	3:X:1256:HOH:O	2.46	0.48
1:X:532:GLU:OE1	1:X:588:ARG:HD3	2.14	0.48
1:X:112:LEU:HD22	1:X:365:ARG:HH12	1.79	0.48
1:X:381:LYS:HE3	3:X:1442:HOH:O	2.13	0.48
1:X:493:ARG:NH2	1:X:547:GLU:HB3	2.29	0.48
1:X:580:LYS:O	1:X:586:GLY:HA3	2.14	0.48
1:X:497:TYR:O	1:X:501:ILE:HD13	2.13	0.47
1:X:168:ARG:HA	1:X:171:MET:HE3	1.97	0.47
1:X:582[A]:PRO:CG	1:X:589[A]:PRO:HD3	2.44	0.47
1:X:140[B]:ASN:HD21	1:X:365:ARG:HG3	1.80	0.46
1:X:206:GLU:O	1:X:210:GLU:HG3	2.15	0.46
1:X:537:LEU:C	1:X:537:LEU:HD23	2.36	0.46
1:X:169:ARG:HG3	1:X:169:ARG:NH1	2.30	0.46
1:X:171:MET:HE1	1:X:550:LEU:HD11	1.97	0.46
1:X:244:LYS:NZ	1:X:244:LYS:HB3	2.31	0.46
1:X:334:MET:HG3	1:X:373:VAL:HG12	1.98	0.46
1:X:525:ILE:O	1:X:529:ARG:HG2	2.16	0.45
1:X:279[A]:LYS:NZ	3:X:1369:HOH:O	2.48	0.45
1:X:333:ASN:HB2	3:X:1272:HOH:O	2.15	0.45
1:X:112:LEU:C	1:X:112:LEU:HD12	2.37	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:316:TYR:OH	1:X:381:LYS:NZ	2.50	0.45
1:X:421:ARG:HG2	1:X:423:LYS:HE3	1.98	0.45
1:X:441:LEU:HD12	1:X:441:LEU:N	2.31	0.45
1:X:295[B]:THR:HG21	3:X:1360:HOH:O	2.15	0.44
1:X:423:LYS:NZ	3:X:1574:HOH:O	2.49	0.44
1:X:181:PRO:HA	1:X:184:TRP:CE3	2.52	0.44
1:X:332:VAL:HG11	1:X:352:PRO:HB3	1.99	0.44
1:X:491:ILE:HD12	1:X:543:THR:HB	1.99	0.44
1:X:279[A]:LYS:HE2	3:X:1339:HOH:O	2.17	0.44
1:X:319:TYR:CE2	1:X:321:ARG:HB2	2.53	0.44
1:X:491:ILE:HD12	1:X:543:THR:CG2	2.48	0.43
1:X:169:ARG:HG3	1:X:169:ARG:HH11	1.83	0.43
1:X:428:ILE:CD1	1:X:449:ILE:HG12	2.48	0.43
1:X:35:ARG:NE	1:X:122:HIS:HD2	2.05	0.43
1:X:582[A]:PRO:HG2	1:X:589[A]:PRO:HD3	1.99	0.43
1:X:463[B]:GLU:OE1	1:X:465:THR:OG1	2.36	0.43
1:X:428:ILE:HD13	1:X:449:ILE:HG12	2.01	0.42
1:X:95:GLU:O	1:X:98:GLN:HG2	2.19	0.42
1:X:346:LYS:HG3	1:X:347:TYR:CD1	2.54	0.42
1:X:539:ASN:HD21	1:X:541:ASP:HB2	1.85	0.42
1:X:407:LYS:HE3	3:X:1113:HOH:O	2.20	0.42
1:X:67:LEU:HD22	1:X:71:MET:CE	2.50	0.41
1:X:180:SER:HB2	1:X:181:PRO:HD3	2.02	0.41
1:X:441:LEU:H	1:X:441:LEU:CD1	2.33	0.41
1:X:345:LYS:HG2	3:X:1311:HOH:O	2.20	0.41
1:X:539:ASN:ND2	1:X:542:PHE:H	2.03	0.41
1:X:513:GLU:HG2	1:X:518:TYR:CD1	2.56	0.41
1:X:121:TRP:O	1:X:125:GLU:HG2	2.21	0.41
1:X:271:ARG:HG3	1:X:413:LEU:CD2	2.51	0.41
1:X:379[A]:SER:O	1:X:381:LYS:HD3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	575/592 (97%)	568 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	499/501 (100%)	487 (98%)	12 (2%)	49	28

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	63	GLU
1	X	67	LEU
1	X	78	ASN
1	X	136	ARG
1	X	189	ARG
1	X	250	ASN
1	X	320	MET
1	X	354	GLN
1	X	381	LYS
1	X	423	LYS
1	X	470	PHE
1	X	521	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	69	ASN
1	X	78	ASN
1	X	98	GLN
1	X	122	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	X	145	ASN
1	X	237	GLN
1	X	241	GLN
1	X	250	ASN
1	X	255	GLN
1	X	329	GLN
1	X	454	ASN
1	X	539	ASN
1	X	545	GLN
1	X	548	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	X	2003	-	4,4,4	0.13	0	6,6,6	0.07	0
2	SO4	X	2007	-	4,4,4	0.18	0	6,6,6	0.17	0
2	SO4	X	2008	-	4,4,4	0.15	0	6,6,6	0.09	0
2	SO4	X	2005	-	4,4,4	0.14	0	6,6,6	0.06	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	X	2004	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	X	2002	-	4,4,4	0.14	0	6,6,6	0.06	0
2	SO4	X	2001	-	4,4,4	0.15	0	6,6,6	0.08	0
2	SO4	X	2009	-	4,4,4	0.15	0	6,6,6	0.11	0
2	SO4	X	2006	-	4,4,4	0.14	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	X	556/592 (93%)	0.53	37 (6%) <b>17</b> <b>18</b>	16, 25, 37, 60	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	441	LEU	9.5
1	X	470	PHE	8.5
1	X	2	GLY	6.8
1	X	39	THR	5.4
1	X	440	ALA	5.2
1	X	101	GLU	5.2
1	X	592	MET	4.2
1	X	446	GLU	4.1
1	X	447	ASP	3.9
1	X	449	ILE	3.7
1	X	102	LYS	3.7
1	X	24	ASP	3.5
1	X	153	LYS	3.2
1	X	542	PHE	3.2
1	X	439	PRO	3.0
1	X	130	ASP	3.0
1	X	98	GLN	3.0
1	X	581	ASP	2.9
1	X	152	ASN	2.7
1	X	273	ASP	2.7
1	X	448	ARG	2.6
1	X	381	LYS	2.6
1	X	148	ARG	2.6
1	X	513	GLU	2.5
1	X	345	LYS	2.4
1	X	189	ARG	2.4
1	X	77	HIS	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	X	506	ASP	2.3
1	X	149	GLN	2.3
1	X	514	ASN	2.3
1	X	104	ASP	2.3
1	X	365	ARG	2.2
1	X	169	ARG	2.1
1	X	428	ILE	2.1
1	X	382	ASP	2.1
1	X	491	ILE	2.1
1	X	584	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	X	2008	5/5	0.89	0.17	64,64,65,65	0
2	SO4	X	2009	5/5	0.90	0.12	50,50,51,51	0
2	SO4	X	2006	5/5	0.92	0.34	66,66,66,66	0
2	SO4	X	2002	5/5	0.93	0.17	71,71,71,71	0
2	SO4	X	2003	5/5	0.94	0.18	42,43,43,43	0
2	SO4	X	2004	5/5	0.95	0.14	58,59,59,59	0
2	SO4	X	2005	5/5	0.97	0.13	38,38,38,39	0
2	SO4	X	2001	5/5	0.98	0.11	38,38,39,40	0
2	SO4	X	2007	5/5	0.98	0.10	29,30,30,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.