

Full wwPDB X-ray Structure Validation Report (i)

Oct 2, 2023 - 03:31 AM EDT

PDB ID : 6MXS

Title : Crystal structure of the dimeric bH1-Fab variant [HC-Y33W,HC-D98F,HC-G

99M]

Authors: Shi, R.; Picard, M.-E.; Manenda, M.S.

Deposited on : 2018-10-31

Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.95 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 7555 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called anti-VEGF-A Fab fragment bH1 heavy chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Н	217	Total 1654	C 1050	11	O 317	S 9	0	3	0
1	A	218	Total 1657	C 1051		O 318	S 9	0	2	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Н	220	GLY	-	expression tag	UNP V9HW68
Н	221	HIS	-	expression tag	UNP V9HW68
Н	222	HIS	-	expression tag	UNP V9HW68
Н	223	HIS	-	expression tag	UNP V9HW68
Н	224	HIS	-	expression tag	UNP V9HW68
Н	225	HIS	-	expression tag	UNP V9HW68
Н	226	HIS	-	expression tag	UNP V9HW68
Н	227	HIS	-	expression tag	UNP V9HW68
Н	228	HIS	-	expression tag	UNP V9HW68
Н	229	GLY	-	expression tag	UNP V9HW68
A	220	GLY	-	expression tag	UNP V9HW68
A	221	HIS	-	expression tag	UNP V9HW68
A	222	HIS	-	expression tag	UNP V9HW68
A	223	HIS	-	expression tag	UNP V9HW68
A	224	HIS	-	expression tag	UNP V9HW68
A	225	HIS	-	expression tag	UNP V9HW68
A	226	HIS	-	expression tag	UNP V9HW68
A	227	HIS	-	expression tag	UNP V9HW68
A	228	HIS		expression tag	UNP V9HW68
A	229	GLY	-	expression tag	UNP V9HW68

• Molecule 2 is a protein called anti-VEGF-A Fab fragment bH1 light chain.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	Т	218	Total	С	N	О	S	0	2	0
	L	210	1698	1066	282	343	7	U		
9	D	217	Total	С	N	О	S	0	2	0
	D	211	1691	1062	282	340	7	U		


• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	L	1	Total Cl 1 1	0	0
3	В	1	Total Cl 1 1	0	0

• Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	1	Total Na 1 1	0	0

 \bullet Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total C O 4 2 2	0	0
5	В	1	Total C O 4 2 2	0	0



• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	Н	189	Total O 189 189	0	0
6	L	232	Total O 232 232	0	0
6	A	175	Total O 175 175	0	0
6	В	248	Total O 248 248	0	0

Mol Probity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	65.68Å 95.96Å 110.11Å	Donositor	
a, b, c, α , β , γ	90.00° 103.68° 90.00°	Depositor	
Resolution (Å)	50.00 - 1.95	Depositor	
% Data completeness	98.7 (50.00-1.95)	Depositor	
(in resolution range)	, , ,	Depositor	
R_{merge}	0.22	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	4.53 (at 1.95Å)	Xtriage	
Refinement program	REFMAC 5.8.0073	Depositor	
R, R_{free}	0.191 , 0.229	Depositor	
Wilson B-factor (\mathring{A}^2)	25.2	Xtriage	
Anisotropy	0.508	Xtriage	
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.32$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	7555	wwPDB-VP	
Average B, all atoms (\mathring{A}^2)	29.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trunc	Chain	Dag	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	В	302	-	3,3,3	0.50	0	2,2,2	0.40	0
5	EDO	В	301	-	3,3,3	0.83	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	В	302	-	-	1/1/1/1	-
5	EDO	В	301	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	В	302	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

