wwPDB X-ray Structure Validation Summary Report (i)

Oct 17, 2021 – 07:11 AM EDT

PDB ID	:	1N5N
Title	:	Crystal Structure of Peptide Deformylase from Pseudomonas aeruginosa
Authors	:	Kreusch, A.; Spraggon, G.; Lee, C.C.; Klock, H.; McMullan, D.; Ng, K.; Shin, T.; Vincent, J.; Warner, I.; Ericson, C.; Lesley, S.A.
Deposited on	:	2002-11-06
Resolution	:	1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	180	65% 22% 5% • 7%
1	B	180	67% 17% 7% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	301	-	X	-	-
3	GOL	A	303	-	X	-	-
3	GOL	B	302	-	X	X	-

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptide deformylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	167	1347	849	235	256	7	0	0	0
1	B	164	1328	838	231	252	7	0	0	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	UNP Q9I7A8
A	-10	GLY	-	expression tag	UNP Q9I7A8
A	-9	SER	-	expression tag	UNP Q9I7A8
A	-8	ASP	-	expression tag	UNP Q9I7A8
A	-7	LYS	-	expression tag	UNP Q9I7A8
A	-6	ILE	-	expression tag	UNP Q9I7A8
A	-5	HIS	-	expression tag	UNP Q9I7A8
A	-4	HIS	-	expression tag	UNP Q9I7A8
A	-3	HIS	-	expression tag	UNP Q9I7A8
A	-2	HIS	-	expression tag	UNP Q9I7A8
A	-1	HIS	-	expression tag	UNP Q9I7A8
A	0	HIS	-	expression tag	UNP Q9I7A8
A	84	GLU	ASP	engineered mutation	UNP Q9I7A8
B	-11	MET	-	expression tag	UNP Q9I7A8
B	-10	GLY	-	expression tag	UNP Q9I7A8
B	-9	SER	-	expression tag	UNP Q9I7A8
B	-8	ASP	-	expression tag	UNP Q9I7A8
B	-7	LYS	-	expression tag	UNP Q9I7A8
B	-6	ILE	-	expression tag	UNP Q9I7A8
B	-5	HIS	-	expression tag	UNP Q9I7A8
B	-4	HIS	-	expression tag	UNP Q9I7A8
B	-3	HIS	-	expression tag	UNP Q9I7A8
B	-2	HIS	-	expression tag	UNP Q9I7A8
B	-1	HIS	-	expression tag	UNP Q9I7A8
B	0	HIS	-	expression tag	UNP Q9I7A8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	84	GLU	ASP	engineered mutation	UNP Q9I7A8

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	140	Total O 140 140	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	B	122	122	122	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Peptide deformylase

Chain A: 65% 22% 5% • 7%



• Molecule 1: Peptide deformylase

Chain B: 67% 17% 7% 9%



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	68.61Å 73.56Å 76.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.87 – 1.80	Depositor
% Data completeness (in resolution range)	99.1 (41.87-1.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.196 , 0.244	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2957	wwPDB-VP
Average B, all atoms ($Å^2$)	24.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.92	29/1371 (2.1%)	1.66	19/1853 (1.0%)
1	B	1.79	12/1352 (0.9%)	1.80	28/1827 (1.5%)
All	All	1.86	41/2723 (1.5%)	1.73	47/3680 (1.3%)

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	90	GLU	CG-CD	10.61	1.67	1.51
1	B	37	GLU	CG-CD	9.55	1.66	1.51
1	A	120	PHE	CE1-CZ	9.26	1.54	1.37
1	A	78	PHE	CE2-CZ	7.65	1.51	1.37
1	A	88	TYR	CD1-CE1	7.51	1.50	1.39

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	157	ARG	NE-CZ-NH2	-21.82	109.39	120.30
1	B	157	ARG	NE-CZ-NH1	20.01	130.31	120.30
1	A	165	ARG	NE-CZ-NH1	-13.95	113.33	120.30
1	B	109	ARG	NE-CZ-NH2	-9.18	115.71	120.30
1	A	71	ARG	NE-CZ-NH2	-8.95	115.83	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1347	0	1339	13	0
1	B	1328	0	1324	23	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	12	0	8	2	0
3	B	6	0	7	8	0
4	A	140	0	0	7	0
4	B	122	0	0	13	0
All	All	2957	0	2678	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 41 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:302:GOL:O1	3:B:302:GOL:C1	1.63	1.41
1:B:165:ARG:HD3	4:B:502:HOH:O	1.57	1.03
1:B:101:ASN:HA	4:B:518:HOH:O	1.61	1.00
1:B:5:ASN:HB2	4:B:504:HOH:O	1.67	0.93
1:B:91:GLY:H	3:B:302:GOL:H31	1.35	0.89

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/180 (92%)	164 (99%)	1 (1%)	0	100	100
1	B	162/180 (90%)	158 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	327/360 (91%)	322 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	149/162 (92%)	142 (95%)	7 (5%)	26	12
1	B	148/162 (91%)	143 (97%)	5 (3%)	37	22
All	All	297/324 (92%)	285 (96%)	12 (4%)	31	16

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	41	GLU
1	B	65	GLU
1	B	101	ASN
1	B	86	ASP
1	A	67	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	GOL	A	303	-	5,5,5	4.67	3 (60%)	5,5,5	2.22	2 (40%)
3	GOL	A	301	-	5,5,5	5.34	4 (80%)	5,5,5	2.36	1 (20%)
3	GOL	B	302	-	5,5,5	4.09	5 (100%)	5,5,5	3.65	4 (80%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	303	-	-	1/4/4/4	-
3	GOL	A	301	-	-	2/4/4/4	-
3	GOL	B	302	-	-	0/4/4/4	-

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	GOL	C3-C2	-9.63	1.12	1.51
3	A	303	GOL	C3-C2	-8.02	1.18	1.51
3	A	301	GOL	O1-C1	-5.62	1.18	1.42
3	A	303	GOL	C1-C2	-5.19	1.30	1.51
3	B	302	GOL	O1-C1	5.01	1.63	1.42

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	302	GOL	C3-C2-C1	5.19	131.90	111.70
3	A	301	GOL	O2-C2-C1	5.12	131.66	109.12
3	B	302	GOL	O2-C2-C1	-4.62	88.77	109.12
3	A	303	GOL	O3-C3-C2	3.76	128.23	110.20
3	B	302	GOL	O3-C3-C2	-3.13	95.20	110.20

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	GOL	O1-C1-C2-O2
3	A	301	GOL	O2-C2-C3-O3
3	A	303	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	GOL	2	0
3	B	302	GOL	8	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.