



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 12, 2023 – 02:44 PM EDT

PDB ID : 4N7T
Title : Crystal structure of phosphorylated phosphopentomutase from streptococcus mutans
Authors : Fedorov, A.A.; Fedorov, E.V.; Bonanno, J.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC); New York Structural Genomics Research Consortium (NYSGRC)
Deposited on : 2013-10-16
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	413	 6% 93% 5%
1	B	413	 5% 92% 5%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphopentomutase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	402	Total	C	N	O	P	S	0	0	0
			3094	1956	529	598	1	10			
1	B	401	Total	C	N	O	P	S	0	0	0
			3088	1953	528	596	1	10			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP Q8DTU0
A	0	SER	-	expression tag	UNP Q8DTU0
A	1	LEU	-	expression tag	UNP Q8DTU0
A	404	GLU	-	expression tag	UNP Q8DTU0
A	405	GLY	-	expression tag	UNP Q8DTU0
A	406	HIS	-	expression tag	UNP Q8DTU0
A	407	HIS	-	expression tag	UNP Q8DTU0
A	408	HIS	-	expression tag	UNP Q8DTU0
A	409	HIS	-	expression tag	UNP Q8DTU0
A	410	HIS	-	expression tag	UNP Q8DTU0
A	411	HIS	-	expression tag	UNP Q8DTU0
B	-1	MET	-	expression tag	UNP Q8DTU0
B	0	SER	-	expression tag	UNP Q8DTU0
B	1	LEU	-	expression tag	UNP Q8DTU0
B	404	GLU	-	expression tag	UNP Q8DTU0
B	405	GLY	-	expression tag	UNP Q8DTU0
B	406	HIS	-	expression tag	UNP Q8DTU0
B	407	HIS	-	expression tag	UNP Q8DTU0
B	408	HIS	-	expression tag	UNP Q8DTU0
B	409	HIS	-	expression tag	UNP Q8DTU0
B	410	HIS	-	expression tag	UNP Q8DTU0
B	411	HIS	-	expression tag	UNP Q8DTU0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mn 2 2	0	0
2	B	2	Total Mn 2 2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	1

- Molecule 5 is AZIDE ION (three-letter code: AZI) (formula: N₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total N 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total N 3 3	0	0

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 7 4 3	0	1

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	195	Total O 199 199	0	4
7	B	166	Total O 170 170	0	4

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.79Å 63.69Å 104.77Å 90.00° 98.85° 90.00°	Depositor
Resolution (Å)	29.27 – 2.00 29.27 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (29.27-2.00) 99.0 (29.27-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.93 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.171 , 0.203 0.171 , 0.202	Depositor DCC
R_{free} test set	1872 reflections (3.05%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtrriage
Anisotropy	0.557	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 53.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6596	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, SO4, GOL, PEG, AZI, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/3154	0.51	0/4290
1	B	0.35	0/3148	0.52	0/4282
All	All	0.36	0/6302	0.52	0/8572

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3094	0	3022	11	0
1	B	3088	0	3017	10	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	18	0	24	0	0
5	A	3	0	0	0	0
5	B	3	0	0	0	0
6	A	7	0	10	1	0
7	A	199	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	170	0	0	1	0
All	All	6596	0	6073	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:193:THR:HG21	1:A:200:LEU:HD12	1.83	0.60
1:B:5:ASN:ND2	1:B:286:THR:O	2.28	0.60
1:A:3:THR:HG21	1:A:403:ILE:HG12	1.84	0.59
1:A:41:HIS:NE2	6:A:507[A]:PEG:H11	2.24	0.53
1:B:173:HIS:CE1	1:B:175:ASP:HB2	2.45	0.51
1:B:14:SER:HB2	1:B:342:ASN:HB2	1.93	0.49
1:A:14:SER:HB2	1:A:342:ASN:HB2	1.93	0.49
1:A:183:TYR:OH	1:A:220:ASN:O	2.22	0.44
1:B:99:ILE:O	1:B:380:SER:HB3	2.18	0.44
1:B:149:GLY:O	1:B:153:MET:HG3	2.18	0.44
1:B:145:ILE:HD11	1:B:171:ALA:HB1	1.99	0.44
1:B:117:GLU:OE1	1:B:133:ARG:NE	2.37	0.43
1:B:187:GLU:HG3	7:B:741:HOH:O	2.18	0.43
1:B:172:ALA:O	1:B:207:PRO:HA	2.18	0.43
1:A:99:ILE:O	1:A:380:SER:HB3	2.18	0.43
1:A:194:LEU:HD13	1:B:194:LEU:HD11	2.01	0.42
1:A:165:ASP:OD1	7:A:753:HOH:O	2.22	0.41
1:A:162:THR:OG1	1:A:163:SER:N	2.54	0.40
1:A:200:LEU:HD23	1:A:200:LEU:HA	1.94	0.40
1:A:3:THR:HG21	1:A:403:ILE:CG1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/413 (97%)	380 (95%)	18 (4%)	1 (0%)	41	37
1	B	398/413 (96%)	378 (95%)	20 (5%)	0	100	100
All	All	797/826 (96%)	758 (95%)	38 (5%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	214	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	332/342 (97%)	329 (99%)	3 (1%)	78	83
1	B	331/342 (97%)	327 (99%)	4 (1%)	71	76
All	All	663/684 (97%)	656 (99%)	7 (1%)	73	78

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	139	TYR
1	A	291	PHE
1	A	310	TYR
1	B	139	TYR
1	B	222	HIS
1	B	291	PHE
1	B	310	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	92	1,2	8,10,11	1.08	0	10,14,16	1.02	0
1	TPO	B	92	1,2	8,10,11	1.16	0	10,14,16	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	92	1,2	-	4/9/11/13	-
1	TPO	B	92	1,2	-	3/9/11/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	92	TPO	C-CA-CB-CG2
1	A	92	TPO	O-C-CA-CB
1	B	92	TPO	O-C-CA-CB
1	B	92	TPO	C-CA-CB-CG2
1	A	92	TPO	CB-OG1-P-O1P
1	A	92	TPO	CB-OG1-P-O2P
1	B	92	TPO	CB-OG1-P-O2P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	AZI	B	504	-	0,2,2	-	-	0,1,1	-	-
3	SO4	B	503	-	4,4,4	0.15	0	6,6,6	0.09	0
4	GOL	A	505	-	5,5,5	0.30	0	5,5,5	0.25	0
4	GOL	A	504	-	5,5,5	0.41	0	5,5,5	0.41	0
5	AZI	A	506	-	0,2,2	-	-	0,1,1	-	-
3	SO4	A	503	-	4,4,4	0.15	0	6,6,6	0.13	0
6	PEG	A	507[A]	-	6,6,6	0.38	0	5,5,5	0.42	0
4	GOL	A	508[B]	-	5,5,5	0.67	0	5,5,5	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	505	-	-	0/4/4/4	-
6	PEG	A	507[A]	-	-	2/4/4/4	-
4	GOL	A	508[B]	-	-	0/4/4/4	-
4	GOL	A	504	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	504	GOL	C1-C2-C3-O3
6	A	507[A]	PEG	O2-C3-C4-O4
4	A	504	GOL	O2-C2-C3-O3
6	A	507[A]	PEG	C1-C2-O2-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	507[A]	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	401/413 (97%)	0.07	26 (6%) 18 18	12, 26, 60, 86	0
1	B	400/413 (96%)	0.13	19 (4%) 30 29	13, 27, 58, 79	0
All	All	801/826 (96%)	0.10	45 (5%) 24 23	12, 26, 60, 86	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	112	TRP	5.5
1	A	2	SER	4.6
1	B	214	ASN	4.2
1	B	403	ILE	4.2
1	A	142	THR	3.7
1	A	213	ARG	3.6
1	B	210	GLY	3.4
1	B	11	VAL	3.2
1	A	216	THR	3.1
1	B	213	ARG	3.1
1	A	219	ALA	3.1
1	A	150	PRO	2.9
1	A	215	PHE	2.9
1	A	197	PRO	2.9
1	A	9	LEU	2.8
1	A	11	VAL	2.8
1	B	336	ILE	2.8
1	A	335	LEU	2.7
1	B	215	PHE	2.7
1	B	75	HIS	2.6
1	B	176	VAL	2.6
1	A	217	ARG	2.5
1	B	12	LEU	2.5
1	B	99	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	12	LEU	2.4
1	A	222	HIS	2.4
1	B	9	LEU	2.3
1	A	403	ILE	2.3
1	B	74	ASN	2.3
1	A	214	ASN	2.2
1	A	291	PHE	2.2
1	B	366	THR	2.2
1	B	113	ASN	2.2
1	A	218	THR	2.2
1	B	110	THR	2.2
1	A	336	ILE	2.1
1	A	338	ALA	2.1
1	A	99	ILE	2.1
1	B	142	THR	2.1
1	A	240	ALA	2.1
1	A	111	PHE	2.1
1	A	220	ASN	2.1
1	B	139	TYR	2.1
1	A	10	VAL	2.0
1	A	172	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	A	92	11/12	0.95	0.14	16,24,40,42	4
1	TPO	B	92	11/12	0.96	0.11	12,21,36,43	4

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	504	6/6	0.68	0.20	40,49,50,51	0
4	GOL	A	505	6/6	0.86	0.21	34,38,42,43	0
3	SO4	A	503	5/5	0.90	0.15	69,71,74,76	0
3	SO4	B	503	5/5	0.91	0.12	81,82,84,85	0
6	PEG	A	507[A]	7/7	0.91	0.20	16,18,21,22	7
5	AZI	B	504	3/3	0.94	0.12	7,7,22,25	0
4	GOL	A	508[B]	6/6	0.94	0.15	6,15,19,21	6
5	AZI	A	506	3/3	0.97	0.12	7,7,30,32	0
2	MN	B	502	1/1	0.99	0.04	22,22,22,22	0
2	MN	A	501	1/1	1.00	0.09	18,18,18,18	0
2	MN	A	502	1/1	1.00	0.07	21,21,21,21	0
2	MN	B	501	1/1	1.00	0.08	15,15,15,15	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.