



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2023 – 10:50 PM EDT

PDB ID : 3N8W  
Title : Crystal Structure of R120Q/Native Cyclooxygenase-1 Heterodimer mutant in complex with Flurbiprofen  
Authors : Sidhu, R.S.  
Deposited on : 2010-05-28  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	553	 3% 88% 11%
2	B	553	 3% 89% 10%
3	C	2	 100%
3	F	2	 50% 50%
4	D	3	 100%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	E	5	 100%
6	G	3	 100%
7	H	3	 67%  33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	C	1	X	-	-	-
9	FLP	A	701	X	-	-	-

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 9505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prostaglandin G/H synthase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	553	4437	2877	747	785	28	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	92	LEU	MET	conflict	UNP P05979

- Molecule 2 is a protein called Prostaglandin G/H synthase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	553	4472	2901	744	800	27	0	8	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	92	LEU	MET	conflict	UNP P05979
B	120	GLN	ARG	conflict	UNP P05979

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	3	39	22	2	15	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	E	5	61	34	2	25	0	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	G	3	39	22	2	15	0	0	0

- Molecule 7 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
7	H	3	39	22	2	15	0	0	0

- Molecule 8 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
8	A	1	43	34	1	4	4	0	0
8	B	1	43	34	1	4	4	0	0

- Molecule 9 is FLURBIPROFEN (three-letter code: FLP) (formula:  $C_{15}H_{13}FO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	F	O		
9	A	1	18	15	1	2	0	0

- Molecule 10 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula:  $C_{14}H_{28}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
10	A	1	20	14	6	0	0
10	B	1	13	7	6	0	0
10	B	1	12	6	6	0	0

- Molecule 11 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
11	A	105	Total 105	O 105	0	0
11	B	108	Total 108	O 108	0	0




### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Prostaglandin G/H synthase 1



- Molecule 2: Prostaglandin G/H synthase 1



- Molecule 3: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 3: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%

MAG1  
MAG2

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1  
MAG2  
BMA3

- Molecule 5: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1  
MAG2  
BMA3  
MAN4  
MAN5

- Molecule 6: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

MAG1  
MAG2  
MAN3

- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  67% 33%

MAG1  
MAG2  
BMA3

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	182.48Å 182.48Å 103.09Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.91 – 2.75 46.91 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.0 (46.91-2.75) 99.5 (46.91-2.75)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 2.77Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.178 , 0.201 0.191 , 0.211	Depositor DCC
$R_{free}$ test set	1712 reflections (3.38%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.6	Xtrriage
Anisotropy	0.048	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 32.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.064 for h,-h-k,-l	Xtrriage
Reported twinning fraction	0.532 for H, K, L 0.468 for -H-K, K, -L	Depositor
Outliers	0 of 50719 reflections	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9505	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BOG, BMA, FLP, NAG, NDG, MAN, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/4576	0.51	0/6224
2	B	0.35	0/4612	0.51	0/6281
All	All	0.35	0/9188	0.51	0/12505

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4437	0	4280	36	0
2	B	4472	0	4299	47	0
3	C	28	0	24	0	0
3	F	28	0	24	0	0
4	D	39	0	34	0	0
5	E	61	0	51	0	0
6	G	39	0	33	0	0
7	H	39	0	33	1	0
8	A	43	0	30	1	0
8	B	43	0	30	1	0
9	A	18	0	12	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	A	20	0	28	1	0
10	B	25	0	22	0	0
11	A	105	0	0	5	0
11	B	108	0	0	5	0
All	All	9505	0	8900	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:124[B]:ILE:O	2:B:126[B]:SER:N	1.71	1.21
2:B:122[B]:ASN:O	2:B:124[B]:ILE:N	1.87	1.07
2:B:185:ARG:HG3	2:B:185:ARG:HH11	1.24	1.02
2:B:122[B]:ASN:C	2:B:124[B]:ILE:H	1.63	1.01
2:B:124[B]:ILE:C	2:B:126[B]:SER:H	1.65	0.99
2:B:41:CYS:HB2	11:B:640:HOH:O	1.63	0.98
2:B:126[B]:SER:HA	2:B:127[B]:PRO:C	1.91	0.90
2:B:124[B]:ILE:C	2:B:126[B]:SER:N	2.25	0.83
1:A:563:THR:HG22	1:A:566:LYS:H	1.43	0.82
2:B:125[B]:PRO:O	2:B:127[B]:PRO:O	1.99	0.79
2:B:124[B]:ILE:O	2:B:124[B]:ILE:HG22	1.84	0.77
1:A:369:ALA:HB3	11:A:628:HOH:O	1.87	0.74
2:B:122[B]:ASN:OD1	2:B:123[B]:LEU:N	2.22	0.70
2:B:122[B]:ASN:CG	2:B:123[B]:LEU:H	1.97	0.69
2:B:122[B]:ASN:C	2:B:124[B]:ILE:N	2.34	0.68
2:B:185:ARG:HG3	2:B:185:ARG:NH1	2.03	0.68
2:B:57:CYS:SG	11:B:640:HOH:O	2.54	0.66
2:B:122[B]:ASN:CG	2:B:123[B]:LEU:N	2.50	0.65
2:B:124[B]:ILE:O	2:B:125[B]:PRO:C	2.33	0.65
2:B:563:THR:HG22	2:B:566:LYS:H	1.62	0.65
1:A:459:ARG:HD3	11:A:24:HOH:O	2.01	0.59
1:A:344:VAL:HA	1:A:348:TYR:HB3	1.88	0.56
1:A:342:LYS:HG2	1:A:562:ALA:HB3	1.87	0.55
1:A:185:ARG:HH21	1:A:438:ARG:HG2	1.73	0.54
2:B:113:MET:O	2:B:117:LEU:HD13	2.08	0.54
1:A:119:VAL:HG12	10:A:751:BOG:H62	1.90	0.53
1:A:387:TRP:HB2	8:A:801:HEM:HAC	1.91	0.53
2:B:344:VAL:HA	2:B:348:TYR:HB3	1.90	0.52
1:A:121:SER:HB2	1:A:532:LYS:HD3	1.91	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:433:ARG:HH11	2:B:436:GLY:HA3	1.75	0.51
1:A:504:TYR:HB3	1:A:505:PRO:HD3	1.92	0.51
2:B:126[A]:SER:HA	2:B:127[A]:PRO:C	2.31	0.51
1:A:287:VAL:HG11	1:A:302:ALA:HB1	1.92	0.51
1:A:388:HIS:N	1:A:389:PRO:CD	2.75	0.50
1:A:275:TYR:CE2	1:A:284:GLN:HB2	2.47	0.50
2:B:41:CYS:SG	2:B:68:ASN:HA	2.53	0.49
2:B:387:TRP:HB2	8:B:601:HEM:HAC	1.94	0.49
1:A:507:LEU:HB3	1:A:522:MET:CE	2.43	0.48
2:B:344:VAL:O	2:B:348:TYR:HB3	2.14	0.48
2:B:447:VAL:O	2:B:451:VAL:HG23	2.13	0.48
1:A:281:PRO:HA	1:A:284:GLN:HG2	1.96	0.48
2:B:124[B]:ILE:HG22	2:B:126[B]:SER:N	2.28	0.48
2:B:124[B]:ILE:HG22	2:B:126[B]:SER:H	1.78	0.48
2:B:339:GLU:HG2	2:B:562:ALA:HB2	1.96	0.47
1:A:216:MET:HE1	11:A:614:HOH:O	2.13	0.47
1:A:532:LYS:HE3	11:A:615:HOH:O	2.15	0.47
2:B:319:GLU:OE2	2:B:554:VAL:HG21	2.14	0.47
1:A:487:MET:HB3	1:A:487:MET:HE2	1.78	0.47
2:B:34:ASN:HA	2:B:49:ARG:HH21	1.80	0.47
2:B:124[B]:ILE:O	2:B:126[B]:SER:O	2.34	0.46
2:B:183:LEU:HD13	2:B:445:LEU:HD22	1.98	0.46
1:A:389:PRO:HB2	1:A:434:ILE:HA	1.97	0.46
1:A:182:LEU:HB3	1:A:440:ILE:HD12	1.98	0.45
2:B:122[B]:ASN:OD1	2:B:123[B]:LEU:HD23	2.15	0.45
1:A:183:LEU:HD13	1:A:445:LEU:HD22	1.97	0.45
1:A:97:ARG:HB3	1:A:97:ARG:NH1	2.32	0.44
1:A:507:LEU:HD22	1:A:522:MET:HE2	2.00	0.43
1:A:238:LEU:HD13	7:H:2:NDG:H6C2	2.00	0.43
2:B:84:PRO:HG2	2:B:89:ILE:HD11	2.00	0.43
2:B:181:PHE:HB3	2:B:509:LEU:HD21	2.00	0.43
2:B:388:HIS:N	2:B:389:PRO:CD	2.82	0.43
1:A:577:TYR:CE2	1:A:583:PRO:HD3	2.54	0.43
2:B:320:HIS:HB3	2:B:323:TRP:CG	2.53	0.43
2:B:518:PHE:CD1	2:B:522:MET:HG2	2.54	0.42
1:A:344:VAL:O	1:A:348:TYR:HB3	2.18	0.42
2:B:126[B]:SER:CA	2:B:127[B]:PRO:C	2.72	0.42
1:A:330:GLN:HB3	2:B:138:SER:HB2	2.00	0.42
2:B:185:ARG:NH1	2:B:185:ARG:CG	2.72	0.42
1:A:84:PRO:HG2	1:A:89:ILE:HD11	2.01	0.42
2:B:389:PRO:HB2	2:B:434:ILE:HA	2.01	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:388:HIS:CE1	1:A:447:VAL:HG11	2.56	0.41
2:B:124[B]:ILE:HG22	2:B:126[B]:SER:O	2.20	0.41
2:B:442:HIS:CD2	11:B:671:HOH:O	2.72	0.41
2:B:513:HIS:HB2	2:B:516:SER:OG	2.21	0.41
1:A:530:SER:O	1:A:534:LEU:HD22	2.20	0.41
1:A:34:ASN:HA	1:A:35:PRO:HD2	1.95	0.41
1:A:117:LEU:HB3	1:A:366:LEU:HD21	2.03	0.41
1:A:137:ILE:HD12	11:B:635:HOH:O	2.21	0.41
1:A:215:LYS:H	1:A:215:LYS:HD3	1.86	0.41
2:B:455:SER:C	11:B:623:HOH:O	2.59	0.41
1:A:154:SER:HB2	1:A:459:ARG:HB2	2.03	0.40
1:A:120:ARG:HG2	11:A:663:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	551/553 (100%)	525 (95%)	26 (5%)	0	100	100
2	B	559/553 (101%)	528 (94%)	26 (5%)	5 (1%)	17	31
All	All	1110/1106 (100%)	1053 (95%)	52 (5%)	5 (0%)	41	47

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	123[A]	LEU
2	B	123[B]	LEU
2	B	125[A]	PRO
2	B	125[B]	PRO
2	B	204	HIS

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	475/488 (97%)	449 (94%)	26 (6%)	21	37
2	B	480/488 (98%)	462 (96%)	18 (4%)	33	53
All	All	955/976 (98%)	911 (95%)	44 (5%)	27	46

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	78	LEU
1	A	117	LEU
1	A	185	ARG
1	A	215	LYS
1	A	238	LEU
1	A	244	LEU
1	A	289	GLN
1	A	298	LEU
1	A	300	LEU
1	A	316	LEU
1	A	384	LEU
1	A	385	TYR
1	A	405	GLU
1	A	450	ASP
1	A	455	SER
1	A	458	LEU
1	A	484	GLU
1	A	518	PHE
1	A	522	MET
1	A	534	LEU
1	A	556	PHE
1	A	563	THR
1	A	570	LEU
1	A	575	CYS
1	A	584	ASP
2	B	185	ARG

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
2	B	213	SER
2	B	232	HIS
2	B	238	LEU
2	B	246	LEU
2	B	289	GLN
2	B	298	LEU
2	B	300	LEU
2	B	316	LEU
2	B	347	GLU
2	B	360	LYS
2	B	385	TYR
2	B	492	GLU
2	B	518	PHE
2	B	534	LEU
2	B	556	PHE
2	B	563	THR
2	B	564	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	400	GLN
2	B	581	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	C	1	3,1	14,14,15	0.43	0	17,19,21	1.25	1 (5%)
3	NDG	C	2	3	14,14,15	0.56	0	17,19,21	1.62	4 (23%)
4	NAG	D	1	1,4	14,14,15	1.90	1 (7%)	17,19,21	2.20	4 (23%)
4	NAG	D	2	4	14,14,15	0.39	0	17,19,21	1.33	2 (11%)
4	BMA	D	3	4	11,11,12	0.57	0	15,15,17	1.55	3 (20%)
5	NAG	E	1	1,5	14,14,15	0.61	0	17,19,21	1.32	2 (11%)
5	NDG	E	2	5	14,14,15	0.61	0	17,19,21	1.95	2 (11%)
5	BMA	E	3	5	11,11,12	0.61	0	15,15,17	0.92	1 (6%)
5	MAN	E	4	5	11,11,12	0.54	0	15,15,17	1.65	3 (20%)
5	MAN	E	5	5	11,11,12	0.62	0	15,15,17	1.53	3 (20%)
3	NAG	F	1	3,2	14,14,15	0.46	0	17,19,21	0.85	0
3	NDG	F	2	3	14,14,15	0.48	0	17,19,21	1.07	1 (5%)
6	NAG	G	1	2,6	14,14,15	0.72	0	17,19,21	1.70	4 (23%)
6	NAG	G	2	6	14,14,15	0.48	0	17,19,21	1.03	1 (5%)
6	MAN	G	3	6	11,11,12	0.55	0	15,15,17	1.12	1 (6%)
7	NAG	H	1	2,7	14,14,15	0.50	0	17,19,21	1.20	1 (5%)
7	NDG	H	2	7	14,14,15	0.64	0	17,19,21	1.98	5 (29%)
7	BMA	H	3	7	11,11,12	0.60	0	15,15,17	1.04	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	1/1/5/7	0/6/23/26	0/1/1/1
3	NDG	C	2	3	-	1/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
5	NAG	E	1	1,5	-	0/6/23/26	0/1/1/1
5	NDG	E	2	5	-	2/6/23/26	0/1/1/1
5	BMA	E	3	5	-	2/2/19/22	0/1/1/1
5	MAN	E	4	5	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MAN	E	5	5	-	0/2/19/22	0/1/1/1
3	NAG	F	1	3,2	-	1/6/23/26	0/1/1/1
3	NDG	F	2	3	-	0/6/23/26	0/1/1/1
6	NAG	G	1	2,6	-	2/6/23/26	0/1/1/1
6	NAG	G	2	6	-	0/6/23/26	0/1/1/1
6	MAN	G	3	6	-	0/2/19/22	0/1/1/1
7	NAG	H	1	2,7	-	0/6/23/26	0/1/1/1
7	NDG	H	2	7	-	1/6/23/26	0/1/1/1
7	BMA	H	3	7	-	1/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1	NAG	O5-C1	-6.85	1.32	1.43

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	2	NDG	C1-O5-C5	6.44	120.92	112.19
4	D	1	NAG	C1-O5-C5	-5.84	104.28	112.19
4	D	1	NAG	O5-C1-C2	-4.92	103.53	111.29
5	E	4	MAN	C1-O5-C5	4.69	118.55	112.19
6	G	1	NAG	C4-C3-C2	4.69	117.89	111.02
4	D	3	BMA	C1-O5-C5	4.43	118.19	112.19
7	H	1	NAG	C1-O5-C5	4.41	118.16	112.19
7	H	2	NDG	C4-C3-C2	4.14	117.08	111.02
7	H	2	NDG	C1-O5-C5	3.90	117.48	112.19
3	C	1	NAG	C1-O5-C5	3.78	117.31	112.19
5	E	1	NAG	C1-O5-C5	3.73	117.25	112.19
7	H	2	NDG	O5-C5-C6	3.69	112.99	107.20
3	C	2	NDG	C3-C4-C5	3.43	116.36	110.24
5	E	5	MAN	C1-O5-C5	3.39	116.78	112.19
6	G	3	MAN	C1-O5-C5	3.27	116.63	112.19
5	E	5	MAN	C3-C4-C5	3.09	115.74	110.24
3	C	2	NDG	C4-C3-C2	3.03	115.46	111.02
3	C	2	NDG	O5-C1-C2	-3.00	106.55	111.29
6	G	1	NAG	C3-C4-C5	2.95	115.50	110.24
4	D	2	NAG	C1-O5-C5	2.88	116.09	112.19
3	F	2	NDG	C1-O5-C5	2.87	116.09	112.19
5	E	4	MAN	C3-C4-C5	2.77	115.18	110.24
5	E	4	MAN	O5-C5-C6	2.75	111.52	107.20
4	D	3	BMA	C1-C2-C3	2.67	112.95	109.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	2	NAG	C4-C3-C2	-2.62	107.18	111.02
5	E	3	BMA	C1-O5-C5	2.60	115.71	112.19
6	G	1	NAG	C2-N2-C7	2.58	126.58	122.90
4	D	1	NAG	C3-C4-C5	2.45	114.61	110.24
5	E	5	MAN	O5-C5-C6	2.36	110.91	107.20
7	H	3	BMA	C3-C4-C5	2.33	114.39	110.24
7	H	2	NDG	C2-N2-C7	2.31	126.19	122.90
5	E	2	NDG	C2-N2-C7	2.27	126.14	122.90
7	H	2	NDG	C3-C4-C5	2.26	114.27	110.24
6	G	2	NAG	C1-O5-C5	2.25	115.23	112.19
5	E	1	NAG	O4-C4-C3	2.23	115.50	110.35
4	D	3	BMA	O5-C5-C6	2.20	110.65	107.20
4	D	1	NAG	O5-C5-C6	2.17	110.60	107.20
3	C	2	NDG	C1-O5-C5	2.11	115.05	112.19
6	G	1	NAG	O5-C1-C2	-2.05	108.05	111.29

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	C	1	NAG	C1

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	E	2	NDG	C3-C2-N2-C7
6	G	1	NAG	C1-C2-N2-C7
4	D	1	NAG	O5-C5-C6-O6
5	E	3	BMA	O5-C5-C6-O6
7	H	3	BMA	O5-C5-C6-O6
4	D	1	NAG	C4-C5-C6-O6
3	C	2	NDG	C4-C5-C6-O6
5	E	3	BMA	C4-C5-C6-O6
3	F	1	NAG	C4-C5-C6-O6
7	H	2	NDG	C3-C2-N2-C7
5	E	2	NDG	C4-C5-C6-O6
6	G	1	NAG	C3-C2-N2-C7


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	H	2	NDG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.















## 5.6 Ligand geometry i

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
10	BOG	A	751	-	20,20,20	0.76	0	25,25,25	1.85	4 (16%)
10	BOG	B	1751	-	12,12,20	0.98	0	17,17,25	1.72	4 (23%)
9	FLP	A	701	-	18,19,19	1.09	1 (5%)	22,26,26	2.20	3 (13%)
8	HEM	A	801	1,11	41,50,50	2.03	6 (14%)	45,82,82	1.61	6 (13%)
8	HEM	B	601	2,11	41,50,50	2.01	7 (17%)	45,82,82	1.64	6 (13%)
10	BOG	B	1750	-	13,13,20	0.87	0	18,18,25	1.69	3 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	BOG	A	751	-	-	7/11/31/31	0/1/1/1
10	BOG	B	1751	-	-	2/2/22/31	0/1/1/1
9	FLP	A	701	-	1/1/2/2	0/8/12/12	0/2/2/2
8	HEM	A	801	1,11	-	5/12/54/54	-
8	HEM	B	601	2,11	-	4/12/54/54	-
10	BOG	B	1750	-	-	3/4/24/31	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	801	HEM	C3D-C2D	8.24	1.54	1.36
8	B	601	HEM	C3D-C2D	8.14	1.54	1.36
8	A	801	HEM	C3C-C2C	-4.57	1.34	1.40
8	B	601	HEM	C3C-C2C	-4.24	1.34	1.40
8	A	801	HEM	C3C-CAC	3.65	1.55	1.47
8	B	601	HEM	C3C-CAC	3.53	1.55	1.47

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	601	HEM	FE-ND	3.50	2.14	1.96
8	A	801	HEM	FE-ND	3.32	2.13	1.96
8	A	801	HEM	CAB-C3B	3.01	1.55	1.47
8	B	601	HEM	CAB-C3B	2.92	1.55	1.47
8	B	601	HEM	CAA-C2A	2.61	1.55	1.52
8	A	801	HEM	CAA-C2A	2.54	1.55	1.52
9	A	701	FLP	C10-C11	2.41	1.40	1.36
8	B	601	HEM	CMB-C2B	2.04	1.55	1.50

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	701	FLP	C10-C11-C6	-8.30	120.11	124.07
8	B	601	HEM	C4D-ND-C1D	6.26	111.54	105.07
8	A	801	HEM	C4D-ND-C1D	6.22	111.49	105.07
10	A	751	BOG	C1'-O1-C1	-5.74	104.32	113.84
10	B	1751	BOG	O5-C1-C2	-5.02	101.32	110.28
9	A	701	FLP	F-C11-C6	4.65	120.71	116.90
10	A	751	BOG	O5-C1-C2	-4.40	101.03	110.35
10	B	1750	BOG	C1'-O1-C1	-3.88	107.29	113.27
10	B	1750	BOG	O5-C1-C2	-3.88	102.14	110.35
8	A	801	HEM	C4B-CHC-C1C	3.35	126.98	122.56
8	B	601	HEM	C1B-NB-C4B	3.07	108.24	105.07
8	B	601	HEM	C4B-CHC-C1C	3.02	126.54	122.56
8	A	801	HEM	C1B-NB-C4B	2.92	108.09	105.07
10	B	1750	BOG	C4-C3-C2	-2.89	105.77	110.82
10	A	751	BOG	C4-C3-C2	-2.85	105.84	110.82
8	B	601	HEM	C4C-CHD-C1D	2.83	126.30	122.56
10	B	1751	BOG	C4-C3-C2	-2.82	105.90	110.82
8	A	801	HEM	C4C-CHD-C1D	2.82	126.28	122.56
10	A	751	BOG	C1-C2-C3	-2.59	104.61	110.00
8	B	601	HEM	CMA-C3A-C4A	-2.52	124.59	128.46
10	B	1751	BOG	C1-C2-C3	-2.30	105.55	110.31
8	A	801	HEM	C3B-C2B-C1B	2.23	108.14	106.49
8	B	601	HEM	C3B-C2B-C1B	2.12	108.06	106.49
10	B	1751	BOG	O5-C5-C4	-2.10	105.88	109.69
9	A	701	FLP	C11-C6-C7	2.05	119.49	116.47
8	A	801	HEM	CBA-CAA-C2A	-2.04	109.14	112.62

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
9	A	701	FLP	C12

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	801	HEM	C2A-CAA-CBA-CGA
10	B	1750	BOG	O5-C1-O1-C1'
10	B	1750	BOG	C2-C1-O1-C1'
10	B	1751	BOG	O5-C5-C6-O6
10	A	751	BOG	O5-C1-O1-C1'
10	B	1751	BOG	C4-C5-C6-O6
10	A	751	BOG	C3'-C4'-C5'-C6'
10	A	751	BOG	C2-C1-O1-C1'
10	A	751	BOG	C2'-C3'-C4'-C5'
10	A	751	BOG	O1-C1'-C2'-C3'
10	A	751	BOG	C4'-C5'-C6'-C7'
8	B	601	HEM	CAA-CBA-CGA-O1A
8	B	601	HEM	CAA-CBA-CGA-O2A
10	A	751	BOG	C5'-C6'-C7'-C8'
8	A	801	HEM	C2B-C3B-CAB-CBB
8	B	601	HEM	C2B-C3B-CAB-CBB
8	A	801	HEM	CAA-CBA-CGA-O2A
8	A	801	HEM	C4B-C3B-CAB-CBB
8	B	601	HEM	C4B-C3B-CAB-CBB
8	A	801	HEM	CAA-CBA-CGA-O1A
10	B	1750	BOG	O5-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	A	751	BOG	1	0
8	A	801	HEM	1	0
8	B	601	HEM	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	553/553 (100%)	0.07	15 (2%) 54 63	43, 56, 71, 79	0
2	B	553/553 (100%)	0.01	16 (2%) 51 61	44, 58, 74, 81	0
All	All	1106/1106 (100%)	0.04	31 (2%) 53 62	43, 56, 72, 81	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	280	PRO	4.2
2	B	123[A]	LEU	4.1
2	B	124[A]	ILE	3.3
2	B	488	ALA	3.2
1	A	107	PHE	3.2
2	B	483	GLY	3.0
2	B	489	ALA	3.0
1	A	281	PRO	3.0
1	A	75	TRP	2.8
1	A	493	GLU	2.8
1	A	261	VAL	2.8
2	B	75	TRP	2.8
2	B	265	SER	2.7
2	B	157	ARG	2.7
1	A	256	MET	2.6
2	B	491	LEU	2.6
2	B	107	PHE	2.5
2	B	32	PRO	2.5
1	A	91	PHE	2.4
1	A	243	GLN	2.4
2	B	98	TRP	2.3
2	B	297	GLY	2.3
1	A	247	PHE	2.3
2	B	487	MET	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	493	GLU	2.2
1	A	489	ALA	2.2
2	B	496	GLY	2.2
1	A	270	PRO	2.1
1	A	246	LEU	2.1
1	A	267	GLU	2.1
1	A	98	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	MAN	E	4	11/12	0.65	0.33	89,91,92,94	0
3	NDG	F	2	14/15	0.84	0.28	74,76,77,77	0
6	MAN	G	3	11/12	0.84	0.31	79,80,80,80	0
7	NDG	H	2	14/15	0.84	0.33	61,64,65,68	0
5	MAN	E	5	11/12	0.85	0.36	95,95,96,96	0
3	NDG	C	2	14/15	0.86	0.32	75,77,79,79	0
7	BMA	H	3	11/12	0.86	0.39	69,70,70,70	0
5	NDG	E	2	14/15	0.88	0.34	64,67,69,73	0
6	NAG	G	1	14/15	0.90	0.27	56,59,63,67	0
4	BMA	D	3	11/12	0.90	0.33	77,78,79,79	0
6	NAG	G	2	14/15	0.91	0.28	71,73,75,77	0
5	BMA	E	3	11/12	0.91	0.38	77,80,84,87	0
3	NAG	F	1	14/15	0.92	0.29	59,65,67,71	0
3	NAG	C	1	14/15	0.92	0.20	60,66,68,72	0
4	NAG	D	1	14/15	0.93	0.19	56,60,62,66	0
4	NAG	D	2	14/15	0.93	0.17	69,71,73,75	0
5	NAG	E	1	14/15	0.94	0.14	46,49,53,58	0
7	NAG	H	1	14/15	0.95	0.13	45,48,51,56	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
10	BOG	A	751	20/20	0.88	0.29	78,82,82,83	0
10	BOG	B	1750	13/20	0.90	0.27	94,94,94,95	0
10	BOG	B	1751	12/20	0.91	0.19	87,88,88,89	0
8	HEM	B	601	43/43	0.96	0.19	54,58,64,66	0
8	HEM	A	801	43/43	0.96	0.18	52,56,62,66	0
9	FLP	A	701	18/18	0.97	0.18	53,55,57,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around BOG B 1750:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around BOG B 1751:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.