



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2024 – 02:27 PM EST

PDB ID : 3NBC  
Title : Clitocybe nebularis ricin B-like lectin (CNL) in complex with lactose, crystallized at pH 4.4  
Authors : Renko, M.; Pohleven, J.; Sabotic, J.; Kos, J.; Turk, D.  
Deposited on : 2010-06-03  
Resolution : 1.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1188 (1.08-0.96)
Clashscore	141614	1253 (1.08-0.96)
Ramachandran outliers	138981	1178 (1.08-0.96)
Sidechain outliers	138945	1180 (1.08-0.96)
RSRZ outliers	127900	1158 (1.08-0.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	 50% (red), 89% (green), 11% (yellow)
1	B	148	 53% (red), 87% (green), 12% (yellow)
2	C	2	 100% (yellow)

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2936 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ricin B-like lectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	148	1156	724	197	234	1	21	7	0
1	B	148	1161	731	195	234	1	12	9	0

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	2	23	12	11	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	349	349	349	0	0
3	B	247	247	247	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ricin B-like lectin



- Molecule 1: Ricin B-like lectin



- Molecule 2: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.79Å 55.90Å 112.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.95 – 1.01 27.95 – 1.01	Depositor EDS
% Data completeness (in resolution range)	98.3 (27.95-1.01) 98.3 (27.95-1.01)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.73 (at 1.01Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.157 , 0.172 0.149 , 0.165	Depositor DCC
$R_{free}$ test set	8906 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.2	Xtrriage
Anisotropy	0.181	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 48.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.010 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	2936	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	2.21	7/1195 (0.6%)	0.93	3/1641 (0.2%)
1	B	2.35	19/1211 (1.6%)	0.98	3/1664 (0.2%)
All	All	2.28	26/2406 (1.1%)	0.95	6/3305 (0.2%)

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	70	TYR	CD2-CE2	5.98	1.48	1.39
1	B	60	TYR	CG-CD2	5.96	1.46	1.39
1	A	105	ARG	CZ-NH2	5.95	1.40	1.33
1	B	135	SER	CB-OG	5.94	1.50	1.42
1	B	44	TYR	CD2-CE2	5.91	1.48	1.39
1	B	125	PRO	N-CD	5.89	1.56	1.47
1	B	96	GLU	CD-OE1	5.87	1.32	1.25
1	A	49	TRP	CD2-CE2	5.81	1.48	1.41
1	B	7	TYR	CE2-CZ	5.73	1.46	1.38
1	B	44	TYR	CZ-OH	5.68	1.47	1.37
1	B	44	TYR	CB-CG	5.67	1.60	1.51
1	B	146	SER	CB-OG	-5.66	1.34	1.42
1	A	28	GLU	CD-OE1	5.61	1.31	1.25
1	A	69	SER	CA-CB	5.59	1.61	1.52
1	B	130[A]	GLU	CD-OE1	5.45	1.31	1.25
1	B	130[B]	GLU	CD-OE1	5.45	1.31	1.25
1	B	2[A]	ILE	N-CA	5.43	1.57	1.46
1	B	2[B]	ILE	N-CA	5.43	1.57	1.46
1	A	26	PRO	N-CD	5.35	1.55	1.47
1	B	70	TYR	CD1-CE1	5.28	1.47	1.39
1	A	70	TYR	CE2-CZ	5.27	1.45	1.38
1	B	133	GLU	CB-CG	5.23	1.62	1.52
1	B	74[A]	ARG	CZ-NH2	5.23	1.39	1.33
1	B	74[B]	ARG	CZ-NH2	5.23	1.39	1.33

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	140	TRP	CD2-CE3	5.12	1.48	1.40
1	A	104	PHE	CE2-CZ	5.09	1.47	1.37

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	120	ASP	CB-CG-OD1	-5.92	112.97	118.30
1	A	100	ASN	N-CA-CB	5.75	120.94	110.60
1	A	120	ASP	CB-CG-OD2	5.37	123.13	118.30
1	B	120	ASP	CB-CG-OD2	5.24	123.01	118.30
1	A	120	ASP	CB-CG-OD1	-5.16	113.66	118.30
1	B	75	ASP	CB-CG-OD2	5.07	122.86	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1156	0	1148	10	0
1	B	1161	0	1173	3	0
2	C	23	0	21	0	0
3	A	349	0	0	7	0
3	B	247	0	0	2	0
All	All	2936	0	2342	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36[A]:HIS:HD2	1:A:122:ASN:HD21	1.04	0.91
1:A:36[A]:HIS:CD2	1:A:122:ASN:HD21	1.93	0.84

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:GLN:HG3	3:A:542:HOH:O	1.78	0.83
1:B:68:GLN:HG3	3:B:554:HOH:O	1.78	0.83
1:A:89:GLN:CG	3:A:542:HOH:O	2.38	0.65
1:A:11:ASN:HD22	1:A:14:TYR:H	1.47	0.61
1:A:107[B]:LYS:HG2	3:A:486:HOH:O	2.04	0.58
1:A:89:GLN:CD	3:A:542:HOH:O	2.50	0.49
1:A:107[B]:LYS:HD2	3:A:518:HOH:O	2.14	0.47
1:B:116[B]:THR:HG21	3:B:441:HOH:O	2.16	0.45
1:A:107[B]:LYS:CD	3:A:486:HOH:O	2.67	0.42
1:B:2[A]:ILE:HD11	1:B:104:PHE:CE2	2.55	0.42
1:A:107[B]:LYS:CG	3:A:486:HOH:O	2.66	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	153/148 (103%)	148 (97%)	5 (3%)	0	100	100
1	B	154/148 (104%)	152 (99%)	2 (1%)	0	100	100
All	All	307/296 (104%)	300 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/127 (106%)	131 (98%)	3 (2%)	52	17
1	B	136/127 (107%)	135 (99%)	1 (1%)	84	58
All	All	270/254 (106%)	266 (98%)	4 (2%)	62	30

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	100	ASN
1	A	146	SER
1	B	100	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	B	29	ASN
1	B	137	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BGC	C	1	2	12,12,12	1.50	2 (16%)	17,17,17	1.86	3 (17%)
2	GAL	C	2	2	11,11,12	2.32	4 (36%)	15,15,17	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	0/2/22/22	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	GAL	O5-C1	4.94	1.51	1.43
2	C	2	GAL	C2-C3	3.77	1.58	1.52
2	C	1	BGC	C4-C3	3.07	1.60	1.52
2	C	1	BGC	C4-C5	2.89	1.59	1.53
2	C	2	GAL	C4-C3	2.73	1.59	1.52
2	C	2	GAL	C4-C5	2.20	1.57	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	BGC	O1-C1-O5	4.28	123.23	110.38
2	C	1	BGC	O5-C1-C2	4.00	117.43	110.28
2	C	1	BGC	O1-C1-C2	2.68	116.59	109.03


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	146/148 (98%)	2.53	74 (50%) 0 1	8, 11, 19, 31	2 (1%)
1	B	147/148 (99%)	2.59	79 (53%) 0 1	8, 11, 19, 30	1 (0%)
All	All	293/296 (98%)	2.56	153 (52%) 0 1	8, 11, 20, 31	3 (1%)

All (153) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	148	VAL	36.8
1	B	148	VAL	33.8
1	B	99	GLY	13.5
1	A	44	TYR	8.7
1	A	99	GLY	8.4
1	B	101	SER	7.9
1	A	146	SER	6.8
1	B	146	SER	6.1
1	B	98	ALA	6.1
1	A	101	SER	5.2
1	B	100	ASN	4.9
1	A	19	ILE	4.6
1	B	115	LEU	4.5
1	B	19	ILE	4.4
1	B	106	ILE	4.4
1	B	126	ILE	4.4
1	B	1[A]	SER	4.3
1	A	126	ILE	4.3
1	B	97	SER	4.2
1	A	106	ILE	4.2
1	A	98	ALA	4.1
1	B	40[A]	THR	4.1
1	A	108	ILE	4.0
1	A	116[A]	THR	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	51[A]	LEU	3.9
1	B	44	TYR	3.8
1	B	71	VAL	3.8
1	B	133	GLU	3.8
1	A	9	ILE	3.7
1	B	93	VAL	3.7
1	A	49	TRP	3.6
1	B	31	LEU	3.6
1	B	84	LEU	3.6
1	A	140	TRP	3.6
1	B	140	TRP	3.6
1	A	127	VAL	3.6
1	B	95	ILE	3.6
1	A	115	LEU	3.5
1	B	9	ILE	3.5
1	A	59	ILE	3.5
1	B	134	THR	3.4
1	A	113	LEU	3.4
1	B	139	LEU	3.4
1	B	59	ILE	3.4
1	B	32	ILE	3.3
1	A	10[A]	THR	3.3
1	B	142	PHE	3.3
1	A	32	ILE	3.3
1	B	73	VAL	3.3
1	A	95	ILE	3.2
1	B	2[A]	ILE	3.2
1	B	33	ILE	3.2
1	A	52	VAL	3.2
1	B	49	TRP	3.2
1	A	79	VAL	3.2
1	A	60	TYR	3.2
1	A	117	LEU	3.2
1	A	30	THR	3.1
1	A	142	PHE	3.1
1	A	51	LEU	3.0
1	A	114	ALA	3.0
1	A	1	SER	3.0
1	A	7	TYR	3.0
1	A	84	LEU	3.0
1	B	108	ILE	3.0
1	A	36[A]	HIS	3.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	71	VAL	3.0
1	A	93	VAL	3.0
1	A	21	LEU	3.0
1	A	70	TYR	3.0
1	B	54	LEU	2.9
1	B	18	LEU	2.9
1	A	133	GLU	2.8
1	B	131	VAL	2.8
1	A	3	THR	2.8
1	A	33	ILE	2.8
1	B	15	THR	2.8
1	B	127	VAL	2.7
1	B	22	THR	2.7
1	B	141	ALA	2.7
1	A	139	LEU	2.7
1	B	74[A]	ARG	2.7
1	A	58	THR	2.7
1	B	89	GLN	2.7
1	B	21	LEU	2.7
1	B	82	ALA	2.7
1	B	128	LEU	2.7
1	B	91	THR	2.7
1	A	12	VAL	2.6
1	B	109	PRO	2.6
1	A	40[A]	THR	2.6
1	A	61	THR	2.6
1	A	73	VAL	2.6
1	B	61	THR	2.6
1	A	31	LEU	2.6
1	A	85	VAL	2.6
1	A	54	LEU	2.6
1	B	64	ALA	2.6
1	B	116[A]	THR	2.5
1	A	89	GLN	2.5
1	B	12	VAL	2.5
1	B	52	VAL	2.5
1	B	113	LEU	2.5
1	B	70	TYR	2.5
1	B	144[A]	SER	2.5
1	A	37	LEU	2.5
1	B	37	LEU	2.5
1	A	64	ALA	2.4

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	117	LEU	2.4
1	B	60	TYR	2.4
1	A	141	ALA	2.4
1	B	7	TYR	2.4
1	A	92	PRO	2.4
1	B	111	LEU	2.4
1	A	104	PHE	2.4
1	A	18	LEU	2.3
1	A	128	LEU	2.3
1	B	119	SER	2.3
1	A	5	GLY	2.3
1	A	74	ARG	2.3
1	A	2	ILE	2.3
1	A	45	GLY	2.3
1	A	86	GLY	2.3
1	B	79	VAL	2.3
1	A	50	THR	2.3
1	A	91	THR	2.3
1	B	10	THR	2.2
1	B	62	MET	2.2
1	A	39	LYS	2.2
1	B	66	ASN	2.2
1	A	131	VAL	2.2
1	B	125	PRO	2.2
1	B	6	THR	2.2
1	A	135	SER	2.2
1	B	13	ALA	2.2
1	A	6	THR	2.2
1	B	3[A]	THR	2.2
1	A	67	PRO	2.1
1	B	92	PRO	2.1
1	A	83	ALA	2.1
1	B	14	TYR	2.1
1	B	30	THR	2.1
1	B	114	ALA	2.1
1	A	14	TYR	2.1
1	B	50	THR	2.1
1	B	56	HIS	2.1
1	A	82	ALA	2.1
1	A	119	SER	2.0
1	B	8	ASN	2.0
1	A	22	THR	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	58	THR	2.0
1	B	102	GLY	2.0
1	B	85	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BGC	C	1	12/12	0.77	0.21	11,16,20,27	12
2	GAL	C	2	11/12	0.90	0.17	8,10,12,13	11

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.