



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2023 – 07:45 PM EDT

PDB ID : 3NBD
Title : Clitocybe nebularis ricin B-like lectin (CNL) in complex with lactose, crystallized at pH 7.1
Authors : Renko, M.; Pohleven, J.; Sabotic, J.; Kos, J.; Turk, D.
Deposited on : 2010-06-03
Resolution : 1.15 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1492 (1.18-1.10)
Clashscore	141614	1537 (1.18-1.10)
Ramachandran outliers	138981	1483 (1.18-1.10)
Sidechain outliers	138945	1480 (1.18-1.10)
RSRZ outliers	127900	1464 (1.18-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	
1	B	148	
2	C	2	
2	D	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	D	1	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2937 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ricin B-like lectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	148	1173	738	196	238	1	12	13	0
1	B	148	1158	726	195	236	1	12	7	0

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-alpha-D-glucopyranoside.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	2	23	12	11	0	0	0
2	D	2	23	12	11	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	283	Total	O	0	0
			283	283		
4	B	252	Total	O	0	0
			252	252		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ricin B-like lectin




- Molecule 1: Ricin B-like lectin



- Molecule 2: beta-D-galactopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: beta-D-galactopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	43.12Å 85.08Å 97.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.03 – 1.15 23.03 – 1.15	Depositor EDS
% Data completeness (in resolution range)	99.5 (23.03-1.15) 99.5 (23.03-1.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 1.15Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.138 , 0.160 0.145 , 0.163	Depositor DCC
R_{free} test set	6456 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	10.6	Xtrriage
Anisotropy	0.282	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2937	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, GLC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	2.15	10/1235 (0.8%)	1.15	9/1693 (0.5%)
1	B	2.09	7/1208 (0.6%)	1.02	4/1655 (0.2%)
All	All	2.12	17/2443 (0.7%)	1.09	13/3348 (0.4%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	102	GLY	C-O	8.88	1.37	1.23
1	B	130	GLU	CD-OE2	7.51	1.33	1.25
1	A	112	GLY	N-CA	7.03	1.56	1.46
1	B	131	VAL	CB-CG1	6.64	1.66	1.52
1	A	119	SER	CA-CB	6.61	1.62	1.52
1	A	140	TRP	CD2-CE2	6.02	1.48	1.41
1	A	74	ARG	CD-NE	5.79	1.56	1.46
1	A	130	GLU	CD-OE2	5.38	1.31	1.25
1	B	60	TYR	CD2-CE2	5.29	1.47	1.39
1	A	105	ARG	NE-CZ	5.27	1.40	1.33
1	A	112	GLY	CA-C	5.25	1.60	1.51
1	B	87	SER	CA-CB	5.14	1.60	1.52
1	A	28	GLU	CD-OE2	5.06	1.31	1.25
1	A	130	GLU	CD-OE1	5.06	1.31	1.25
1	B	52	VAL	CB-CG2	5.05	1.63	1.52
1	B	12	VAL	CB-CG1	5.03	1.63	1.52
1	A	74	ARG	CZ-NH2	5.01	1.39	1.33

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74	ARG	NE-CZ-NH1	16.57	128.59	120.30
1	B	28	GLU	OE1-CD-OE2	11.15	136.68	123.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	51	LEU	CB-CG-CD2	10.23	128.39	111.00
1	A	17	ARG	NE-CZ-NH1	9.32	124.96	120.30
1	B	51	LEU	CB-CG-CD2	6.84	122.63	111.00
1	A	17	ARG	NE-CZ-NH2	-6.65	116.98	120.30
1	A	16[A]	ASN	CB-CG-OD1	6.48	134.57	121.60
1	A	16[B]	ASN	CB-CG-OD1	6.48	134.57	121.60
1	A	120[A]	ASP	CB-CG-OD2	5.85	123.57	118.30
1	A	120[B]	ASP	CB-CG-OD2	5.85	123.57	118.30
1	A	74	ARG	NH1-CZ-NH2	-5.84	112.97	119.40
1	B	102	GLY	O-C-N	5.30	131.18	122.70
1	B	74	ARG	CD-NE-CZ	5.01	130.61	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1173	0	1191	12	0
1	B	1158	0	1159	9	0
2	C	23	0	21	0	0
2	D	23	0	21	0	0
3	A	15	0	0	0	0
3	B	10	0	0	0	0
4	A	283	0	0	1	0
4	B	252	0	0	1	0
All	All	2937	0	2392	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:87:SER:OG	1:B:88[A]:GLN:OE1	2.03	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88[A]:GLN:OE1	1:B:89:GLN:N	2.18	0.76
1:A:137[B]:ASN:OD1	4:A:208:HOH:O	2.07	0.72
1:B:100:ASN:O	1:B:101:SER:CB	2.36	0.72
1:A:127[B]:VAL:HG12	1:A:128:LEU:N	2.06	0.70
1:A:82:ALA:O	1:A:127[B]:VAL:HG13	1.92	0.69
1:B:100:ASN:O	1:B:101:SER:HB2	2.00	0.60
1:B:98:ALA:HB2	4:B:286:HOH:O	2.01	0.60
1:A:107[B]:LYS:HZ3	1:A:107[B]:LYS:HB3	1.69	0.58
1:A:107[B]:LYS:HB3	1:A:107[B]:LYS:NZ	2.24	0.52
1:B:35:HIS:HD2	1:B:36:HIS:O	1.93	0.51
1:A:81:GLY:O	1:A:127[B]:VAL:HG11	2.11	0.50
1:A:81:GLY:C	1:A:127[B]:VAL:CG1	2.80	0.49
1:A:107[B]:LYS:NZ	1:A:107[B]:LYS:CB	2.76	0.49
1:B:100:ASN:O	1:B:101:SER:OG	2.30	0.48
1:A:100:ASN:OD1	1:A:100:ASN:N	2.46	0.47
1:A:127[B]:VAL:CG1	1:A:128:LEU:N	2.75	0.47
1:A:81:GLY:O	1:A:127[B]:VAL:CG1	2.66	0.44
1:B:107[B]:LYS:HB3	1:B:107[B]:LYS:HE3	1.32	0.42
1:A:14:TYR:CD1	1:A:120[A]:ASP:HB3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	158/148 (107%)	153 (97%)	4 (2%)	1 (1%)	25 5
1	B	155/148 (105%)	153 (99%)	1 (1%)	1 (1%)	25 5
All	All	313/296 (106%)	306 (98%)	5 (2%)	2 (1%)	22 5

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	147	ALA
1	B	101	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	140/127 (110%)	138 (99%)	2 (1%)	67 29
1	B	136/127 (107%)	132 (97%)	4 (3%)	42 7
All	All	276/254 (109%)	270 (98%)	6 (2%)	55 12

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	ASN
1	A	148	VAL
1	B	88[A]	GLN
1	B	88[B]	GLN
1	B	100	ASN
1	B	133	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	8	ASN
1	B	29	ASN
1	B	35	HIS
1	B	100	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	C	1	2	12,12,12	1.10	0	17,17,17	1.56	5 (29%)
2	GAL	C	2	2	11,11,12	2.09	5 (45%)	15,15,17	0.60	0
2	GLC	D	1	2	12,12,12	1.30	2 (16%)	17,17,17	0.47	0
2	GAL	D	2	2	11,11,12	1.57	2 (18%)	15,15,17	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	2/2/22/22	0/1/1/1
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	GAL	C1-C2	3.28	1.59	1.52
2	C	2	GAL	O5-C5	3.24	1.50	1.43
2	C	2	GAL	O5-C1	2.98	1.48	1.43
2	C	2	GAL	C2-C3	2.88	1.56	1.52
2	D	1	GLC	C4-C5	2.80	1.58	1.53
2	C	2	GAL	C1-C2	2.79	1.58	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	GAL	O5-C1	-2.39	1.39	1.43
2	C	2	GAL	O4-C4	2.30	1.48	1.43
2	D	1	GLC	O5-C5	2.19	1.49	1.44

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GLC	C3-C4-C5	3.13	115.83	110.24
2	C	1	GLC	O1-C1-O5	2.51	117.93	110.38
2	C	1	GLC	O3-C3-C4	2.47	116.06	110.35
2	C	1	GLC	C4-C3-C2	2.42	115.04	110.82
2	C	1	GLC	O2-C2-C1	2.22	114.32	109.16

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	1	GLC	C4-C5-C6-O6
2	D	1	GLC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	671	-	4,4,4	0.93	0	6,6,6	0.82	0
3	SO4	B	678	-	4,4,4	0.56	0	6,6,6	0.39	0
3	SO4	A	679	-	4,4,4	1.95	1 (25%)	6,6,6	1.79	2 (33%)
3	SO4	A	672	-	4,4,4	0.63	0	6,6,6	0.82	0
3	SO4	A	677	-	4,4,4	1.07	0	6,6,6	0.96	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	679	SO4	O1-S	3.54	1.65	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	679	SO4	O4-S-O2	-3.22	92.52	109.31
3	A	679	SO4	O2-S-O1	2.18	125.52	109.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	147/148 (99%)	1.45	22 (14%) 2 3	6, 9, 24, 50	1 (0%)
1	B	147/148 (99%)	1.64	28 (19%) 1 2	8, 12, 26, 54	1 (0%)
All	All	294/296 (99%)	1.54	50 (17%) 1 2	6, 10, 26, 54	2 (0%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	148	VAL	22.9
1	A	100	ASN	15.2
1	B	146	SER	14.1
1	B	100	ASN	13.3
1	B	99	GLY	11.3
1	B	101	SER	11.0
1	A	101	SER	10.0
1	A	99	GLY	9.5
1	A	148	VAL	9.3
1	A	98	ALA	5.2
1	A	146	SER	5.0
1	A	1[A]	SER	4.5
1	B	98	ALA	3.6
1	B	1	SER	3.5
1	B	19	ILE	3.2
1	B	115	LEU	3.0
1	B	106	ILE	3.0
1	A	127[A]	VAL	2.9
1	B	9	ILE	2.9
1	B	102	GLY	2.9
1	A	102	GLY	2.9
1	B	93	VAL	2.9
1	B	126	ILE	2.9
1	B	41	PRO	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	145	VAL	2.5
1	A	145	VAL	2.5
1	A	32	ILE	2.4
1	B	88[A]	GLN	2.4
1	B	51	LEU	2.4
1	B	144[A]	SER	2.3
1	B	133	GLU	2.3
1	A	85	VAL	2.3
1	B	12	VAL	2.3
1	A	133	GLU	2.3
1	B	14	TYR	2.3
1	A	33	ILE	2.3
1	A	126	ILE	2.2
1	B	56	HIS	2.2
1	B	40	THR	2.2
1	A	79	VAL	2.1
1	B	119[A]	SER	2.1
1	A	9	ILE	2.0
1	B	95	ILE	2.0
1	B	120	ASP	2.0
1	A	16[A]	ASN	2.0
1	A	31	LEU	2.0
1	A	115	LEU	2.0
1	A	116[A]	THR	2.0
1	A	19	ILE	2.0
1	B	140	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	1	12/12	0.54	0.42	22,40,47,49	0
2	GAL	D	2	11/12	0.84	0.18	12,16,20,20	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	C	1	12/12	0.95	0.11	8,12,16,17	0
2	GAL	C	2	11/12	0.97	0.11	6,7,8,12	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	678	5/5	0.88	0.46	64,66,67,68	0
3	SO4	A	679	5/5	0.89	0.14	14,21,23,24	0
3	SO4	A	672	5/5	0.89	0.25	22,25,32,34	0
3	SO4	B	671	5/5	0.90	0.30	24,26,34,35	0
3	SO4	A	677	5/5	0.97	0.18	19,20,25,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.