



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2023 – 04:22 AM EDT

PDB ID : 3NBE
Title : Clitocybe nebularis ricin B-like lectin (CNL) in complex with N,N'-diacetylla ctosediamine
Authors : Renko, M.; Pohleven, J.; Sabotic, J.; Kos, J.; Turk, D.
Deposited on : 2010-06-03
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	 6% 74% 20%
1	B	148	 6% 80% 18%
2	C	2	 100%
2	D	2	 50% 50%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ricin B-like lectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	147	1138	712	193	232	1	0	6	0
1	B	147	1128	706	195	226	1	3	3	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-2-azidoethyl 2-acetamido-2-deoxy-beta-D-glucopyranoside.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	2	34	18	5	11	1	0	0
2	D	2	34	18	5	11	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	215	Total	O	0	0
			215	215		
4	B	146	Total	O	0	0
			146	146		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ricin B-like lectin



- Molecule 1: Ricin B-like lectin



- Molecule 2: 2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-2-azidoethyl 2-acetamido-2-deoxy-beta-D-glucopyranoside



- Molecule 2: 2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-2-azidoethyl 2-acetamido-2-deoxy-beta-D-glucopyranoside



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.88Å 80.95Å 95.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	16.47 – 1.86 16.47 – 1.86	Depositor EDS
% Data completeness (in resolution range)	97.5 (16.47-1.86) 97.5 (16.47-1.86)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.178 , 0.219 0.186 , 0.229	Depositor DCC
R_{free} test set	1396 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	27.3	Xtrriage
Anisotropy	0.086	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2705	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: Z3Q, NGA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.69	16/1181 (1.4%)	1.44	12/1621 (0.7%)
1	B	1.53	6/1161 (0.5%)	1.41	12/1593 (0.8%)
All	All	1.61	22/2342 (0.9%)	1.43	24/3214 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	85	VAL	CB-CG2	9.81	1.73	1.52
1	A	28	GLU	CB-CG	6.87	1.65	1.52
1	A	39[A]	LYS	CD-CE	6.77	1.68	1.51
1	A	39[B]	LYS	CD-CE	6.77	1.68	1.51
1	A	131	VAL	CB-CG2	6.63	1.66	1.52
1	A	44	TYR	CG-CD1	-6.06	1.31	1.39
1	A	24	SER	N-CA	5.96	1.58	1.46
1	A	44	TYR	CB-CG	5.93	1.60	1.51
1	A	65	VAL	CB-CG2	-5.82	1.40	1.52
1	A	44	TYR	CD1-CE1	5.77	1.48	1.39
1	B	32	ILE	CB-CG2	5.72	1.70	1.52
1	A	147	ALA	C-O	5.69	1.34	1.23
1	B	70	TYR	CD2-CE2	5.68	1.47	1.39
1	A	79	VAL	CB-CG2	5.56	1.64	1.52
1	B	108	ILE	CB-CG2	5.55	1.70	1.52
1	A	19	ILE	CA-CB	5.39	1.67	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	72	ARG	CZ-NH2	5.38	1.40	1.33
1	A	85	VAL	CA-CB	5.33	1.66	1.54
1	B	49	TRP	CZ2-CH2	5.22	1.47	1.37
1	A	28	GLU	CD-OE1	-5.18	1.20	1.25
1	B	65	VAL	CB-CG2	5.17	1.63	1.52
1	A	49	TRP	CG-CD1	5.00	1.43	1.36

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	74[A]	ARG	NE-CZ-NH1	-10.14	115.23	120.30
1	B	74[B]	ARG	NE-CZ-NH1	-10.14	115.23	120.30
1	B	72	ARG	NE-CZ-NH1	-7.30	116.65	120.30
1	A	21	LEU	CB-CG-CD2	-7.23	98.71	111.00
1	A	74	ARG	NE-CZ-NH2	-7.10	116.75	120.30
1	B	74[A]	ARG	NE-CZ-NH2	6.98	123.79	120.30
1	B	74[B]	ARG	NE-CZ-NH2	6.98	123.79	120.30
1	A	20	ASP	CB-CG-OD2	-6.95	112.05	118.30
1	B	74[A]	ARG	CD-NE-CZ	6.78	133.09	123.60
1	B	74[B]	ARG	CD-NE-CZ	6.78	133.09	123.60
1	A	100	ASN	N-CA-CB	-6.60	98.71	110.60
1	B	72	ARG	NE-CZ-NH2	6.33	123.47	120.30
1	B	73	VAL	CG1-CB-CG2	6.24	120.88	110.90
1	B	54	LEU	CB-CG-CD2	-6.18	100.50	111.00
1	A	100	ASN	CB-CG-OD1	-5.65	110.31	121.60
1	A	72	ARG	NE-CZ-NH2	5.65	123.12	120.30
1	A	100	ASN	N-CA-C	5.53	125.94	111.00
1	A	101	SER	N-CA-C	-5.44	96.31	111.00
1	A	8	ASN	N-CA-C	-5.39	96.45	111.00
1	A	98	ALA	N-CA-C	-5.34	96.59	111.00
1	B	34	GLY	N-CA-C	-5.27	99.92	113.10
1	A	37	LEU	CB-CG-CD1	-5.19	102.18	111.00
1	A	146	SER	C-N-CA	-5.17	108.79	121.70
1	B	17	ARG	NE-CZ-NH2	5.12	122.86	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	39[B]	LYS	Mainchain
1	A	45	GLY	Mainchain
1	A	54	LEU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1138	0	1127	13	0
1	B	1128	0	1119	12	0
2	C	34	0	13	0	0
2	D	34	0	13	2	0
3	A	5	0	0	1	0
3	B	5	0	0	0	0
4	A	215	0	0	1	2
4	B	146	0	0	3	2
All	All	2705	0	2272	27	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:1:Z3Q:N1C	2:D:1:Z3Q:N1B	1.61	1.48
1:B:74[A]:ARG:HD2	4:B:158:HOH:O	1.82	0.78
1:A:38:ASN:O	1:A:39[B]:LYS:HB3	1.94	0.66
1:B:105:ARG:HH11	1:B:105:ARG:HG3	1.62	0.64
1:A:14:TYR:CD1	1:A:120[B]:ASP:HB3	2.38	0.59
1:A:130[B]:GLU:HG3	4:A:228:HOH:O	2.04	0.58
1:A:16[B]:ASN:ND2	3:A:355:SO4:O1	2.37	0.57
1:B:10[A]:THR:HG22	4:B:194:HOH:O	2.05	0.55
1:A:100:ASN:N	1:A:100:ASN:OD1	2.37	0.52
1:A:14:TYR:CD1	1:A:120[A]:ASP:HB2	2.44	0.51
1:A:98:ALA:O	1:A:100:ASN:N	2.43	0.51
4:B:202:HOH:O	2:D:1:Z3Q:C8	2.58	0.51
1:B:35:HIS:HD2	1:B:123:SER:OG	1.95	0.50
1:B:107:LYS:HE2	1:B:111:LEU:O	2.13	0.49
1:A:146:SER:O	1:A:147:ALA:C	2.52	0.47
1:B:100:ASN:HB3	1:B:103:GLN:HG3	1.96	0.47
1:B:10[A]:THR:HG23	1:B:141:ALA:HB3	1.96	0.47
1:B:105:ARG:HG3	1:B:105:ARG:NH1	2.29	0.46
1:A:58:THR:O	1:A:94:SER:HA	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:ALA:C	1:A:100:ASN:H	2.20	0.44
1:A:14:TYR:CE1	1:A:120[B]:ASP:HB3	2.52	0.44
1:B:14:TYR:CD1	1:B:120:ASP:HB3	2.54	0.43
1:A:139:LEU:HD12	1:A:139:LEU:HA	1.82	0.43
1:B:102:GLY:O	1:B:141:ALA:HA	2.21	0.41
1:B:54:LEU:HA	1:B:55:PRO:HD3	1.93	0.40
1:A:2:ILE:HG12	1:A:104:PHE:CZ	2.56	0.40
1:B:40:THR:HA	1:B:41:PRO:HA	1.55	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:330:HOH:O	4:B:235:HOH:O[1_655]	2.07	0.13
4:A:359:HOH:O	4:B:154:HOH:O[3_555]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	151/148 (102%)	145 (96%)	1 (1%)	5 (3%)	4 0
1	B	148/148 (100%)	141 (95%)	6 (4%)	1 (1%)	22 9
All	All	299/296 (101%)	286 (96%)	7 (2%)	6 (2%)	9 1

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	100	ASN
1	A	101	SER
1	A	99	GLY
1	B	101	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	39[A]	LYS
1	A	39[B]	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	132/127 (104%)	128 (97%)	4 (3%)	41 24
1	B	129/127 (102%)	127 (98%)	2 (2%)	62 49
All	All	261/254 (103%)	255 (98%)	6 (2%)	49 34

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR
1	A	4	PRO
1	A	67	PRO
1	A	139	LEU
1	B	41	PRO
1	B	133	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	103	GLN
1	B	35	HIS
1	B	100	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	Z3Q	C	1	2	19,20,20	1.85	5 (26%)	24,26,26	1.20	3 (12%)
2	NGA	C	2	2	14,14,15	1.29	1 (7%)	17,19,21	1.30	0
2	Z3Q	D	1	2	19,20,20	3.11	3 (15%)	24,26,26	1.65	4 (16%)
2	NGA	D	2	2	14,14,15	1.50	2 (14%)	17,19,21	1.43	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	Z3Q	C	1	2	-	5/12/32/32	0/1/1/1
2	NGA	C	2	2	-	0/6/23/26	0/1/1/1
2	Z3Q	D	1	2	-	2/12/32/32	0/1/1/1
2	NGA	D	2	2	-	1/6/23/26	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	Z3Q	N1B-N1A	12.84	1.57	1.23
2	D	2	NGA	O7-C7	4.47	1.33	1.23
2	C	1	Z3Q	O1-C11	3.99	1.54	1.43
2	C	1	Z3Q	C2-N2	3.57	1.51	1.45
2	C	1	Z3Q	O4-C4	3.30	1.50	1.43
2	C	1	Z3Q	O7-C7	2.99	1.30	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	Z3Q	O1-C1	2.84	1.45	1.40
2	D	2	NGA	C2-N2	2.73	1.51	1.46
2	D	1	Z3Q	C8-C7	-2.32	1.45	1.50
2	C	2	NGA	O3-C3	2.08	1.47	1.43
2	D	1	Z3Q	C7-N2	2.00	1.41	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	Z3Q	O7-C7-C8	-4.89	112.98	122.06
2	D	2	NGA	C1-C2-N2	3.37	116.25	110.49
2	D	1	Z3Q	C8-C7-N2	2.95	121.09	116.10
2	C	1	Z3Q	C12-N1A-N1B	2.87	129.55	115.95
2	C	1	Z3Q	O7-C7-N2	2.45	126.46	121.95
2	D	2	NGA	C4-C3-C2	-2.31	107.63	111.02
2	D	2	NGA	O5-C5-C6	2.29	110.80	107.20
2	D	1	Z3Q	C12-N1A-N1B	-2.26	105.22	115.95
2	D	1	Z3Q	O7-C7-N2	2.17	125.94	121.95
2	C	1	Z3Q	C6-C5-C4	2.07	117.86	113.00

There are no chirality outliers.

All (8) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	C	1	Z3Q	C2-C1-O1-C11
2	C	1	Z3Q	O5-C1-O1-C11
2	D	1	Z3Q	C8-C7-N2-C2
2	D	1	Z3Q	O7-C7-N2-C2
2	C	1	Z3Q	C11-C12-N1A-N1B
2	C	1	Z3Q	C3-C2-N2-C7
2	D	2	NGA	O5-C5-C6-O6
2	C	1	Z3Q	C1-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	Z3Q	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	355	-	4,4,4	1.84	2 (50%)	6,6,6	1.14	1 (16%)
3	SO4	B	356	-	4,4,4	1.23	0	6,6,6	1.51	1 (16%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	355	SO4	O2-S	2.90	1.61	1.46
3	A	355	SO4	O1-S	2.04	1.57	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	356	SO4	O4-S-O2	2.59	122.84	109.31
3	A	355	SO4	O4-S-O3	2.37	119.19	109.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	355	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	147/148 (99%)	0.17	9 (6%) 21 20	17, 23, 45, 61	0
1	B	147/148 (99%)	0.66	9 (6%) 21 20	21, 30, 53, 81	1 (0%)
All	All	294/296 (99%)	0.41	18 (6%) 21 20	17, 26, 53, 81	1 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	147	ALA	12.2
1	B	100	ASN	10.4
1	A	100	ASN	10.1
1	B	147	ALA	9.2
1	B	99	GLY	9.0
1	B	146	SER	7.7
1	B	101	SER	7.7
1	A	99	GLY	6.2
1	A	146	SER	5.1
1	A	101	SER	4.6
1	A	1	SER	3.1
1	B	98	ALA	2.9
1	B	145	VAL	2.9
1	B	133	GLU	2.3
1	A	102	GLY	2.2
1	B	1	SER	2.1
1	A	134	THR	2.0
1	A	98	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	Z3Q	D	1	20/20	0.89	0.18	34,42,46,48	0
2	NGA	C	2	14/15	0.93	0.10	18,21,25,26	0
2	NGA	D	2	14/15	0.93	0.10	23,31,33,34	0
2	Z3Q	C	1	20/20	0.94	0.10	20,30,37,39	1

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	356	5/5	0.96	0.19	44,44,47,51	0
3	SO4	A	355	5/5	0.98	0.07	29,31,34,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.