

Full wwPDB X-ray Structure Validation Report (i)

Feb 21, 2024 – 05:58 AM EST

:	4NB5
:	Crystal Structure of a transcriptional regulator
:	Su, CC.; Radhakrishnan, A.; Yu, E.W.
	2013-10-22
:	1.64 Å(reported)
	: : :

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics Refmac CCP4 Ideal geometry (proteins) Ideal geometry (DNA, RNA)	:::::::::::::::::::::::::::::::::::::::	20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove) Engh & Huber (2001) Parkinson et al. (1996)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	3122(1.66-1.62)
Clashscore	141614	3268(1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	171	2% 75 %	9%	15%
1	В	171	6%	9%	•• 13%
1	С	171	2% 7 6%	8% •	14%
1	D	171	77%	9%	15%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5396 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	145	Total	С	Ν	0	\mathbf{S}	0	9	0
	А	140	1195	742	225	219	9	0	9	
1	В	149	Total	С	Ν	0	S	0	6	0
	ГБ		1210	751	229	220	10	0	0	0
1	С	147	Total	С	Ν	0	S	0	3	0
	U	141	1183	732	225	218	8	0	5	0
1	П	146	Total	С	Ν	0	S	0	5	0
	I D	146	1177	732	222	214	9	U	0	U


• Molecule 1 is a protein called DNA binding protein.

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	3	VAL	-	expression tag	UNP I6Y8F7
А	168	HIS	-	expression tag	UNP I6Y8F7
А	169	HIS	-	expression tag	UNP I6Y8F7
А	170	HIS	-	expression tag	UNP I6Y8F7
А	171	HIS	-	expression tag	UNP I6Y8F7
А	172	HIS	-	expression tag	UNP I6Y8F7
А	173	HIS	-	expression tag	UNP I6Y8F7
В	3	VAL	-	expression tag	UNP I6Y8F7
В	168	HIS	-	expression tag	UNP I6Y8F7
В	169	HIS	-	expression tag	UNP I6Y8F7
В	170	HIS	-	expression tag	UNP I6Y8F7
В	171	HIS	-	expression tag	UNP I6Y8F7
В	172	HIS	-	expression tag	UNP I6Y8F7
В	173	HIS	-	expression tag	UNP I6Y8F7
С	3	VAL	-	expression tag	UNP I6Y8F7
С	168	HIS	-	expression tag	UNP I6Y8F7
С	169	HIS	-	expression tag	UNP I6Y8F7
С	170	HIS	-	expression tag	UNP I6Y8F7
С	171	HIS	-	expression tag	UNP I6Y8F7
С	172	HIS	-	expression tag	UNP I6Y8F7
С	173	HIS	-	expression tag	UNP I6Y8F7

Continu	Continued from previous page						
Chain	Residue	Modelled	Actual	Comment	Reference		
D	3	VAL	-	expression tag	UNP I6Y8F7		
D	168	HIS	-	expression tag	UNP I6Y8F7		
D	169	HIS	-	expression tag	UNP I6Y8F7		
D	170	HIS	-	expression tag	UNP I6Y8F7		
D	171	HIS	-	expression tag	UNP I6Y8F7		
D	172	HIS	-	expression tag	UNP I6Y8F7		
D	173	HIS	-	expression tag	UNP I6Y8F7		

• Molecule 2 is 1,3-dihydroxy propan-2-yl octadecanoate (three-letter code: 2JT) (formula: $\rm C_{21}H_{42}O_4).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total C O 25 21 4	0	0
2	С	1	Total C O 25 21 4	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	149	Total O 149 149	0	0
3	В	133	Total O 133 133	0	0
3	С	146	Total O 146 146	0	0




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	153	Total O 153 153	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: DNA binding protein



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	54.54Å 57.24Å 61.44Å	Depositor
a, b, c, α , β , γ	82.24° 68.40° 72.20°	Depositor
Resolution (Å)	41.16 – 1.64	Depositor
Resolution (A)	41.17 - 1.64	EDS
% Data completeness	97.0 (41.16-1.64)	Depositor
(in resolution range)	97.0(41.17-1.64)	EDS
R _{merge}	0.04	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.05 (at 1.64 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
D D.	0.166 , 0.191	Depositor
R, R_{free}	0.181 , 0.202	DCC
R_{free} test set	3933 reflections $(5.04%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	16.8	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.37, 49.9	EDS
L-test for twinning ²	$ \langle L \rangle = 0.50, \langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5396	wwPDB-VP
Average B, all atoms $(Å^2)$	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.61% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: $2\mathrm{JT}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain Bor		nd lengths	Bond angles	
IVIOI	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.37	1/1234~(0.1%)	0.56	2/1653~(0.1%)
1	В	0.42	2/1244~(0.2%)	0.65	3/1669~(0.2%)
1	С	0.33	0/1202	0.53	1/1612~(0.1%)
1	D	0.30	0/1208	0.47	0/1620
All	All	0.36	3/4888~(0.1%)	0.56	6/6554~(0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
1	А	36	ARG	CZ-NH1	6.69	1.41	1.33
1	В	96	ARG	CZ-NH2	-5.94	1.25	1.33
1	В	96	ARG	CD-NE	-5.59	1.36	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	96	ARG	NE-CZ-NH2	-10.13	115.23	120.30
1	В	96	ARG	CG-CD-NE	-8.81	93.30	111.80
1	А	36	ARG	NE-CZ-NH2	-8.14	116.23	120.30
1	С	36	ARG	NE-CZ-NH1	8.10	124.35	120.30
1	А	36	ARG	NE-CZ-NH1	7.72	124.16	120.30
1	В	96	ARG	NE-CZ-NH1	5.62	123.11	120.30

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1195	0	1216	11	0
1	В	1210	0	1230	21	0
1	С	1183	0	1186	19	0
1	D	1177	0	1197	13	1
2	В	25	0	41	15	0
2	С	25	0	42	7	0
3	А	149	0	0	6	1
3	В	133	0	0	1	0
3	С	146	0	0	4	0
3	D	153	0	0	4	0
All	All	5396	0	4912	60	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:109:ARG:HG3	2:B:201:2JT:H19	1.44	1.00
1:D:75[A]:MET:SD	3:D:276:HOH:O	2.42	0.77
1:A:19:MET:SD	3:A:320:HOH:O	2.44	0.76
1:B:96:ARG:NH1	1:D:20:GLU:OE2	2.20	0.75
1:B:108:GLU:OE2	1:C:111:ARG:NH1	2.25	0.69
1:B:30:GLU:OE1	1:B:36:ARG:NH2	2.27	0.67
1:A:96:ARG:HD2	3:A:327:HOH:O	1.97	0.64
1:A:17:ASP:N	3:A:295:HOH:O	2.32	0.63
1:D:75[B]:MET:SD	1:D:79:PHE:CE2	2.94	0.61
1:B:161:GLN:O	1:B:162:ARG:HB3	2.04	0.58
1:B:18:ILE:HG23	1:B:47:VAL:HG21	1.87	0.57
1:B:32:ARG:HE	2:B:201:2JT:C18	2.18	0.57
1:B:32:ARG:HH21	2:B:201:2JT:H10	1.69	0.57
1:B:32:ARG:HD3	2:B:201:2JT:H6	1.87	0.56
2:C:201:2JT:H14	1:D:112:ALA:HB2	1.88	0.56
1:C:36:ARG:CZ	3:C:368:HOH:O	2.53	0.55
3:A:266:HOH:O	2:B:201:2JT:H38	2.06	0.54



Atom-1	Atom-2	Interatomic	Clash	
		distance (Å)	overlap (Å)	
1:C:32:ARG:NH2	1:D:115:GLU:OE1	2.40	0.54	
1:D:52:ARG:NE	3:D:337:HOH:O	2.38	0.54	
1:B:109:ARG:HG3	2:B:201:2JT:C05	2.28	0.52	
1:C:36:ARG:NH2	3:C:368:HOH:O	2.43	0.52	
1:C:75:MET:SD	3:D:294:HOH:O	2.59	0.52	
1:B:51:GLU:OE1	1:B:96:ARG:NH2	2.42	0.52	
1:D:108:GLU:OE2	1:D:111[A]:ARG:NH1	2.44	0.51	
1:A:115:GLU:OE1	2:B:201:2JT:H16	2.10	0.51	
1:C:28:TYR:CZ	1:D:116:LEU:HG	2.46	0.51	
1:C:32:ARG:HD3	2:C:201:2JT:H33	1.94	0.50	
1:A:96:ARG:CD	3:A:327:HOH:O	2.55	0.50	
1:A:109:ARG:NH2	1:B:109:ARG:HD2	2.27	0.49	
1:B:32:ARG:HE	2:B:201:2JT:H34	1.77	0.49	
1:C:32:ARG:HE	2:C:201:2JT:C19	2.26	0.49	
1:D:75[B]:MET:SD	1:D:79:PHE:CZ	3.06	0.48	
1:B:87:VAL:O	1:B:90:ASP:O	2.32	0.47	
1:C:31:SER:OG	1:C:32:ARG:NH1	2.47	0.47	
1:C:78:GLN:NE2	3:C:438:HOH:O	2.43	0.47	
1:C:32:ARG:HE	2:C:201:2JT:H37	1.79	0.47	
1:B:161:GLN:O	1:B:162:ARG:CB	2.63	0.47	
1:A:112:ALA:HB2	2:B:201:2JT:H34	1.96	0.46	
2:C:201:2JT:H6	1:D:109:ARG:NH1	2.30	0.46	
1:C:30:GLU:OE2	1:C:36:ARG:NH1	2.48	0.46	
1:A:110:ILE:HD11	1:A:152:VAL:HG12	1.99	0.45	
1:B:81:PHE:CZ	2:B:201:2JT:H17	2.51	0.45	
3:A:266:HOH:O	2:B:201:2JT:H9	2.17	0.44	
1:C:144:LEU:C	1:C:144:LEU:HD13	2.38	0.44	
1:A:18:ILE:HD13	1:A:18:ILE:HA	1.91	0.43	
1:B:32:ARG:HE	2:B:201:2JT:H35	1.84	0.43	
1:B:32:ARG:NE	2:B:201:2JT:H35	2.33	0.43	
1:B:154:ASP:HB2	3:B:409:HOH:O	2.19	0.42	
1:C:36:ARG:NE	3:C:368:HOH:O	2.52	0.42	
2:B:201:2JT:C12	2:B:201:2JT:O01	2.67	0.42	
2:C:201:2JT:H37	1:D:112:ALA:HB2	2.02	0.42	
1:C:96:ARG:HG2	1:C:97:LEU:N	2.34	0.42	
1:C:91:ARG:H	1:C:91:ARG:CD	2.32	0.41	
1:C:102:PHE:HD2	2:C:201:2JT:H18	1.85	0.41	
1:B:32:ARG:NH2	2:B:201:2JT:H10	2.34	0.41	
1:D:139:ARG:HD3	3:D:334:HOH:O	2.19	0.41	
1:C:97:LEU:HG	1:C:101:ALA:HB2	2.03	0.41	
1:A:140[B]:GLU:HG3	1:B:100:ASN:HA	2.03	0.41	



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:LEU:C	1:A:144:LEU:HD13	2.40	0.41
1:C:109[B]:ARG:NE	1:D:109:ARG:CZ	2.84	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:154:ASP:OD1	3:A:309:HOH:O[1_545]	2.14	0.06

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	152/171~(89%)	152 (100%)	0	0	100	100
1	В	153/171~(90%)	152~(99%)	0	1 (1%)	22	6
1	\mathbf{C}	148/171~(86%)	147~(99%)	1 (1%)	0	100	100
1	D	149/171~(87%)	149 (100%)	0	0	100	100
All	All	602/684~(88%)	600 (100%)	1 (0%)	1 (0%)	47	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	162	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	124/137~(90%)	119~(96%)	5(4%)	31 7
1	В	125/137~(91%)	123~(98%)	2(2%)	62 39
1	С	120/137~(88%)	115~(96%)	5 (4%)	30 7
1	D	121/137~(88%)	118 (98%)	3~(2%)	47 20
All	All	490/548~(89%)	475 (97%)	15 (3%)	43 13

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	32[A]	ARG
1	А	32[B]	ARG
1	А	85	LEU
1	А	133	GLN
1	А	161	GLN
1	В	15	GLU
1	В	96	ARG
1	С	85	LEU
1	С	91	ARG
1	С	109[A]	ARG
1	С	109[B]	ARG
1	С	133	GLN
1	D	33	SER
1	D	133	GLN
1	D	136	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Res Link		Bo	ond leng	\mathbf{ths}	B	ond ang	les
	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	2JT	В	201	-	24,24,24	2.30	5 (20%)	25,25,25	1.76	4 (16%)
2	2JT	С	201	-	24,24,24	0.66	0	25,25,25	1.09	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2JT	В	201	-	-	14/25/25/25	-
2	2JT	С	201	-	-	11/25/25/25	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	В	201	2 JT	O03-C04	-8.34	1.25	1.46
2	В	201	2JT	O08-C07	-5.23	1.20	1.42
2	В	201	2JT	C07-C04	3.19	1.59	1.51
2	В	201	2JT	C09-C02	2.23	1.57	1.50
2	В	201	2JT	C10-C09	2.04	1.59	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^{o})$	$Ideal(^{o})$
2	В	201	2 JT	C04-O03-C02	4.96	129.99	117.79



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	201	2JT	O03-C02-O01	-4.94	111.76	123.70
2	В	201	2JT	O03-C04-C07	-3.97	94.10	108.36
2	С	201	2JT	O06-C05-C04	-3.03	103.76	111.78
2	С	201	2JT	O08-C07-C04	-2.59	104.92	111.78
2	В	201	2JT	C07-C04-C05	2.07	116.58	112.80

Continued from previous page...

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
2	В	201	2JT	O01-C02-O03-C04
2	В	201	2JT	C09-C02-O03-C04
2	В	201	2JT	O03-C04-C05-O06
2	В	201	2JT	C07-C04-C05-O06
2	С	201	2JT	O01-C02-O03-C04
2	С	201	2JT	C09-C02-O03-C04
2	С	201	2JT	C16-C17-C18-C19
2	С	201	2JT	C02-C09-C10-C11
2	В	201	2JT	C18-C19-C20-C21
2	С	201	2JT	C13-C14-C15-C16
2	С	201	2JT	C19-C20-C21-C22
2	С	201	2JT	C09-C10-C11-C12
2	В	201	2JT	C11-C12-C13-C14
2	С	201	2JT	C18-C19-C20-C21
2	В	201	2JT	C05-C04-O03-C02
2	В	201	2JT	C02-C09-C10-C11
2	В	201	2JT	C16-C17-C18-C19
2	В	201	2JT	C12-C13-C14-C15
2	В	201	2JT	C17-C18-C19-C20
2	С	201	2JT	C07-C04-C05-O06
2	В	201	2JT	C14-C15-C16-C17
2	С	201	2JT	C14-C15-C16-C17
2	С	201	2JT	O03-C04-C05-O06
2	В	201	2JT	C21-C22-C23-C24
2	В	201	2JT	O03-C02-C09-C10

All (25) torsion outliers are listed below:

There are no ring outliers.

2 monomers are involved in 22 short contacts:


Mol	Chain	\mathbf{Res}	Type	Clashes	Symm-Clashes
2	В	201	2JT	15	0




Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	201	2JT	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	145/171~(84%)	-0.07	3 (2%) 63 64	7, 14, 35, 61	0
1	В	149/171~(87%)	0.24	10 (6%) 17 15	8, 18, 40, 66	0
1	С	147/171~(85%)	-0.08	3 (2%) 65 65	7, 16, 36, 72	0
1	D	146/171~(85%)	-0.06	3 (2%) 63 64	11, 20, 43, 58	0
All	All	587/684~(85%)	0.01	19 (3%) 47 46	7, 17, 40, 72	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	163	THR	7.1
1	В	89	GLY	6.7
1	В	162	ARG	5.4
1	В	88	ALA	4.4
1	В	91	ARG	4.1
1	В	87	VAL	3.6
1	С	162	ARG	3.6
1	В	15	GLU	3.5
1	С	91	ARG	3.3
1	С	161	GLN	3.2
1	В	161	GLN	3.0
1	А	91	ARG	2.8
1	D	161	GLN	2.8
1	А	161	GLN	2.7
1	В	19[A]	MET	2.6
1	А	129	ASP	2.3
1	D	66	SER	2.3
1	В	90	ASP	2.2
1	D	16	PRO	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	2JT	С	201	25/25	0.74	0.28	$28,\!50,\!67,\!77$	0
2	2JT	В	201	25/25	0.76	0.24	19,41,69,82	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

