



Full wwPDB X-ray Structure Validation Report i

Oct 6, 2021 – 09:20 am BST

PDB ID : 7NB7
Title : Structure of Mcl-1 complex with compound 6b
Authors : Dokurno, P.; Surgenor, A.E.; Kotschy, A.
Deposited on : 2021-01-25
Resolution : 2.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3617 (2.84-2.80)
Clashscore	141614	4060 (2.84-2.80)
Ramachandran outliers	138981	3978 (2.84-2.80)
Sidechain outliers	138945	3980 (2.84-2.80)
RSRZ outliers	127900	3552 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 3 unique types of molecules in this entry. The entry contains 4891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Induced myeloid leukemia cell differentiation protein Mcl-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	151	Total	C	N	O	S	0	0	0
			1187	745	214	224	4			
1	B	151	Total	C	N	O	S	0	0	0
			1184	744	214	222	4			
1	C	151	Total	C	N	O	S	0	0	0
			1184	744	214	222	4			
1	D	151	Total	C	N	O	S	0	0	0
			1190	747	217	222	4			

There are 52 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	158	MET	-	initiating methionine	UNP Q07820
A	159	HIS	-	expression tag	UNP Q07820
A	160	HIS	-	expression tag	UNP Q07820
A	161	HIS	-	expression tag	UNP Q07820
A	162	HIS	-	expression tag	UNP Q07820
A	163	HIS	-	expression tag	UNP Q07820
A	164	HIS	-	expression tag	UNP Q07820
A	165	LEU	-	expression tag	UNP Q07820
A	166	VAL	-	expression tag	UNP Q07820
A	167	PRO	-	expression tag	UNP Q07820
A	168	ARG	-	expression tag	UNP Q07820
A	169	GLY	-	expression tag	UNP Q07820
A	170	SER	-	expression tag	UNP Q07820
B	158	MET	-	initiating methionine	UNP Q07820
B	159	HIS	-	expression tag	UNP Q07820
B	160	HIS	-	expression tag	UNP Q07820
B	161	HIS	-	expression tag	UNP Q07820
B	162	HIS	-	expression tag	UNP Q07820
B	163	HIS	-	expression tag	UNP Q07820
B	164	HIS	-	expression tag	UNP Q07820
B	165	LEU	-	expression tag	UNP Q07820

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	166	VAL	-	expression tag	UNP Q07820
B	167	PRO	-	expression tag	UNP Q07820
B	168	ARG	-	expression tag	UNP Q07820
B	169	GLY	-	expression tag	UNP Q07820
B	170	SER	-	expression tag	UNP Q07820
C	158	MET	-	initiating methionine	UNP Q07820
C	159	HIS	-	expression tag	UNP Q07820
C	160	HIS	-	expression tag	UNP Q07820
C	161	HIS	-	expression tag	UNP Q07820
C	162	HIS	-	expression tag	UNP Q07820
C	163	HIS	-	expression tag	UNP Q07820
C	164	HIS	-	expression tag	UNP Q07820
C	165	LEU	-	expression tag	UNP Q07820
C	166	VAL	-	expression tag	UNP Q07820
C	167	PRO	-	expression tag	UNP Q07820
C	168	ARG	-	expression tag	UNP Q07820
C	169	GLY	-	expression tag	UNP Q07820
C	170	SER	-	expression tag	UNP Q07820
D	158	MET	-	initiating methionine	UNP Q07820
D	159	HIS	-	expression tag	UNP Q07820
D	160	HIS	-	expression tag	UNP Q07820
D	161	HIS	-	expression tag	UNP Q07820
D	162	HIS	-	expression tag	UNP Q07820
D	163	HIS	-	expression tag	UNP Q07820
D	164	HIS	-	expression tag	UNP Q07820
D	165	LEU	-	expression tag	UNP Q07820
D	166	VAL	-	expression tag	UNP Q07820
D	167	PRO	-	expression tag	UNP Q07820
D	168	ARG	-	expression tag	UNP Q07820
D	169	GLY	-	expression tag	UNP Q07820
D	170	SER	-	expression tag	UNP Q07820

- Molecule 2 is (2 {R})-2-[[7-but-2-ynyl-5-(3-chloranyl-2-methyl-phenyl)-6-ethyl-pyrrolo [2,3-d]pyrimidin-4-yl]amino]-3-phenyl-propanoic acid (three-letter code: U6N) (formula: C₂₈H₂₇ClN₄O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total		C	Cl	N	O	
			35	28	1	4	2		
2	B	1	Total		C	Cl	N	O	
			35	28	1	4	2		
2	C	1	Total		C	Cl	N	O	
			35	28	1	4	2		
2	D	1	Total		C	Cl	N	O	
			35	28	1	4	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total		O	
			2	2	0	0
3	B	2	Total		O	
			2	2	0	0
3	C	1	Total		O	
			1	1	0	0
3	D	1	Total		O	
			1	1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Induced myeloid leukemia cell differentiation protein Mcl-1




- Molecule 1: Induced myeloid leukemia cell differentiation protein Mcl-1



- Molecule 1: Induced myeloid leukemia cell differentiation protein Mcl-1



- Molecule 1: Induced myeloid leukemia cell differentiation protein Mcl-1





4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	89.28Å 166.11Å 44.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.82 19.90 – 2.82	Depositor EDS
% Data completeness (in resolution range)	97.2 (20.00-2.82) 97.5 (19.90-2.82)	Depositor EDS
R_{merge}	0.28	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.39 (at 2.83Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R , R_{free}	0.252 , 0.291 0.254 , 0.289	Depositor DCC
R_{free} test set	802 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtriage
Anisotropy	0.156	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	4891	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: U6N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/1206	0.90	1/1626 (0.1%)
1	B	0.64	0/1203	0.81	0/1622
1	C	0.64	0/1203	0.84	0/1622
1	D	0.65	0/1209	0.83	0/1629
All	All	0.66	0/4821	0.85	1/6499 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	198	PRO	N-CA-CB	-6.99	94.91	102.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1187	0	1169	22	0
1	B	1184	0	1167	24	0
1	C	1184	0	1167	16	0
1	D	1190	0	1178	21	0
2	A	35	0	0	0	0
2	B	35	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	35	0	0	0	0
2	D	35	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	2	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
All	All	4891	0	4681	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:312:TRP:O	1:D:316:VAL:HG23	1.65	0.95
1:B:312:TRP:O	1:B:316:VAL:HG23	1.67	0.95
1:D:261:TRP:O	1:D:265:VAL:HG23	1.68	0.93
1:C:312:TRP:O	1:C:316:VAL:HG23	1.67	0.93
3:B:501:HOH:O	1:C:224:HIS:HA	1.75	0.86
2:B:401:U6N:O1	3:B:501:HOH:O	1.96	0.82
1:D:184:ARG:HD2	1:D:199:MET:HE1	1.73	0.71
1:A:251:ILE:HD11	1:B:251:ILE:HD13	1.73	0.70
1:B:259:THR:OG1	1:B:302:LYS:NZ	2.31	0.64
1:C:175:TYR:HE2	1:C:296:ASP:OD1	1.82	0.63
1:A:300:ARG:HB2	1:B:256:ASP:HB2	1.82	0.62
1:D:208:LYS:HD3	1:D:211:GLU:OE1	2.00	0.61
1:A:251:ILE:HD13	1:B:251:ILE:HD11	1.81	0.61
1:D:300:ARG:HD2	1:D:301:THR:HG23	1.83	0.60
1:D:184:ARG:HD2	1:D:199:MET:CE	2.30	0.60
1:B:310:ARG:NH2	1:B:317:GLU:OE1	2.35	0.59
1:A:180:GLU:CD	1:A:199:MET:O	2.42	0.59
1:C:175:TYR:CE2	1:C:296:ASP:OD1	2.56	0.58
1:A:251:ILE:CD1	1:B:251:ILE:CD1	2.83	0.56
1:A:251:ILE:CD1	1:B:251:ILE:HD13	2.36	0.56
1:A:258:VAL:HG13	1:D:226:THR:OG1	2.06	0.56
1:B:180:GLU:OE2	1:B:199:MET:HA	2.05	0.56
1:C:284:GLU:H	1:C:284:GLU:CD	2.11	0.55
1:B:284:GLU:H	1:B:284:GLU:CD	2.10	0.54
1:C:213:LEU:HD21	1:C:268:ILE:HG21	1.90	0.54
1:A:284:GLU:H	1:A:284:GLU:CD	2.10	0.54
1:C:208:LYS:HB3	1:C:316:VAL:HG11	1.87	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:213:LEU:HD21	1:D:268:ILE:HG21	1.89	0.54
1:D:284:GLU:CD	1:D:284:GLU:H	2.11	0.53
1:A:257:GLY:HA2	1:D:226:THR:HG21	1.89	0.53
1:A:251:ILE:HD13	1:B:251:ILE:CD1	2.39	0.52
1:A:305:TRP:O	1:A:309:GLN:HG2	2.10	0.52
1:C:305:TRP:O	1:C:309:GLN:HG2	2.10	0.52
1:B:213:LEU:HD21	1:B:268:ILE:HG21	1.90	0.52
1:A:180:GLU:OE2	1:A:199:MET:O	2.28	0.52
1:D:199:MET:HB2	1:D:203:GLY:HA3	1.91	0.52
1:B:305:TRP:O	1:B:309:GLN:HG2	2.09	0.52
1:D:276:LYS:O	1:D:280:THR:HG23	2.10	0.51
1:D:296:ASP:O	1:D:300:ARG:HB3	2.10	0.51
1:D:305:TRP:O	1:D:309:GLN:HG2	2.10	0.51
1:A:276:LYS:O	1:A:280:THR:HG23	2.11	0.51
1:C:216:VAL:HG12	1:C:265:VAL:HG11	1.92	0.51
1:C:276:LYS:O	1:C:280:THR:HG23	2.11	0.51
1:A:310:ARG:NH2	1:A:317:GLU:OE2	2.42	0.50
1:B:170:SER:O	1:D:233:ARG:NH2	2.41	0.50
1:D:246:LEU:HD12	1:D:246:LEU:O	2.12	0.50
1:B:276:LYS:O	1:B:280:THR:HG23	2.12	0.50
1:C:217:GLY:HA2	1:C:269:SER:OG	2.12	0.49
1:B:216:VAL:HG12	1:B:265:VAL:HG11	1.94	0.49
1:A:252:HIS:HD2	1:B:300:ARG:CZ	2.25	0.48
1:C:296:ASP:O	1:C:300:ARG:HB3	2.13	0.48
1:B:216:VAL:O	1:B:220:VAL:HG23	2.15	0.47
1:D:216:VAL:HG12	1:D:265:VAL:HG11	1.97	0.47
1:B:309:GLN:O	1:B:314:GLY:HA3	2.16	0.46
1:A:173:GLU:HA	1:A:173:GLU:OE1	2.16	0.45
1:B:290:LEU:O	1:B:294:ILE:HG13	2.16	0.45
1:A:216:VAL:HG12	1:A:265:VAL:HG11	1.98	0.45
1:A:309:GLN:O	1:A:314:GLY:HA3	2.18	0.44
1:A:255:SER:HA	1:A:302:LYS:HE2	2.00	0.44
1:B:257:GLY:HA2	1:C:226:THR:HG21	2.01	0.43
1:A:217:GLY:HA2	1:A:269:SER:OG	2.19	0.42
1:A:300:ARG:HE	1:A:300:ARG:HB3	1.47	0.42
1:C:251:ILE:HA	1:C:255:SER:HG	1.83	0.42
1:D:184:ARG:CD	1:D:199:MET:CE	2.97	0.42
1:B:305:TRP:CH2	1:B:309:GLN:HG3	2.54	0.42
1:B:288:GLU:O	1:B:292:GLU:HG3	2.20	0.41
1:C:309:GLN:O	1:C:314:GLY:HA3	2.20	0.41
1:A:305:TRP:CH2	1:A:309:GLN:HG3	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:246:LEU:O	1:D:250:MET:HG3	2.21	0.41
1:B:208:LYS:HB3	1:B:316:VAL:HG11	2.02	0.41
1:D:305:TRP:CH2	1:D:309:GLN:HG3	2.57	0.40
1:C:305:TRP:CH2	1:C:309:GLN:HG3	2.57	0.40
1:D:309:GLN:O	1:D:314:GLY:HA3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	149/170 (88%)	140 (94%)	7 (5%)	2 (1%)	12 34
1	B	149/170 (88%)	146 (98%)	3 (2%)	0	100 100
1	C	149/170 (88%)	144 (97%)	5 (3%)	0	100 100
1	D	149/170 (88%)	143 (96%)	6 (4%)	0	100 100
All	All	596/680 (88%)	573 (96%)	21 (4%)	2 (0%)	41 70

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	198	PRO
1	A	199	MET

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	126/148 (85%)	115 (91%)	11 (9%)	10 28
1	B	125/148 (84%)	116 (93%)	9 (7%)	14 37
1	C	125/148 (84%)	113 (90%)	12 (10%)	8 24
1	D	126/148 (85%)	117 (93%)	9 (7%)	14 38
All	All	502/592 (85%)	461 (92%)	41 (8%)	11 31

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	194	LYS
1	A	195	ASP
1	A	198	PRO
1	A	202	SER
1	A	207	ARG
1	A	208	LYS
1	A	226	THR
1	A	238	LYS
1	A	269	SER
1	A	293	SER
1	A	300	ARG
1	B	171	GLU
1	B	184	ARG
1	B	195	ASP
1	B	196	THR
1	B	229	GLN
1	B	247	SER
1	B	293	SER
1	B	296	ASP
1	B	317	GLU
1	C	171	GLU
1	C	176	ARG
1	C	184	ARG
1	C	195	ASP
1	C	199	MET
1	C	239	ASN
1	C	245	SER
1	C	247	SER
1	C	255	SER
1	C	293	SER
1	C	296	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	317	GLU
1	D	171	GLU
1	D	176	ARG
1	D	184	ARG
1	D	195	ASP
1	D	196	THR
1	D	201	ARG
1	D	225	GLU
1	D	296	ASP
1	D	300	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	252	HIS
1	A	282	ASN
1	B	252	HIS
1	B	282	ASN
1	D	282	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	U6N	B	401	-	32,38,38	0.87	1 (3%)	36,53,53	1.13	3 (8%)
2	U6N	C	401	-	32,38,38	0.85	1 (3%)	36,53,53	1.17	4 (11%)
2	U6N	A	401	-	32,38,38	0.90	2 (6%)	36,53,53	1.04	2 (5%)
2	U6N	D	401	-	32,38,38	0.88	2 (6%)	36,53,53	1.12	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	U6N	B	401	-	-	3/17/22/22	0/4/4/4
2	U6N	C	401	-	-	3/17/22/22	0/4/4/4
2	U6N	A	401	-	-	3/17/22/22	0/4/4/4
2	U6N	D	401	-	-	2/17/22/22	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	U6N	C22-C4	-3.03	1.46	1.50
2	A	401	U6N	C6-C5	-2.72	1.41	1.44
2	A	401	U6N	C22-C4	-2.32	1.47	1.50
2	D	401	U6N	C6-C5	-2.31	1.41	1.44
2	C	401	U6N	C22-C4	-2.21	1.47	1.50
2	D	401	U6N	C22-C4	-2.19	1.47	1.50

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	U6N	C5-C6-N2	-3.77	118.24	121.35
2	B	401	U6N	C19-C18-N4	3.76	117.40	112.10
2	D	401	U6N	C5-C6-N2	-3.42	118.53	121.35
2	A	401	U6N	C5-C6-N2	-3.28	118.65	121.35
2	C	401	U6N	C5-C6-N2	-3.23	118.69	121.35
2	A	401	U6N	C19-C18-N4	3.15	116.54	112.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	U6N	C2-C3-C4	-2.79	125.58	130.34
2	C	401	U6N	C5-C6-N1	2.47	122.98	120.66
2	D	401	U6N	C19-C18-N4	2.45	115.56	112.10
2	B	401	U6N	C5-C6-N1	2.11	122.64	120.66
2	C	401	U6N	C18-N4-C3	2.05	128.10	123.97
2	D	401	U6N	C22-C4-C5	2.01	128.55	124.37

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	U6N	C28-C22-C4-C5
2	A	401	U6N	N4-C18-C19-C20
2	B	401	U6N	N4-C18-C19-C20
2	C	401	U6N	C28-C22-C4-C5
2	C	401	U6N	N4-C18-C19-C20
2	D	401	U6N	C28-C22-C4-C5
2	B	401	U6N	C28-C22-C4-C5
2	C	401	U6N	C28-C22-C4-C3
2	A	401	U6N	C28-C22-C4-C3
2	D	401	U6N	C28-C22-C4-C3
2	B	401	U6N	C28-C22-C4-C3


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	U6N	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/170 (88%)	0.05	5 (3%) 46 36	13, 29, 62, 107	1 (0%)
1	B	151/170 (88%)	0.37	11 (7%) 15 8	13, 35, 102, 149	7 (4%)
1	C	151/170 (88%)	0.30	6 (3%) 38 28	16, 40, 75, 119	7 (4%)
1	D	151/170 (88%)	0.51	12 (7%) 12 7	19, 45, 87, 120	7 (4%)
All	All	604/680 (88%)	0.31	34 (5%) 24 16	13, 38, 78, 149	22 (3%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	170	SER	9.6
1	C	170	SER	7.9
1	B	199	MET	7.0
1	A	170	SER	6.4
1	D	170	SER	6.1
1	C	171	GLU	5.9
1	D	171	GLU	5.4
1	B	198	PRO	5.3
1	B	203	GLY	4.6
1	B	171	GLU	4.6
1	B	202	SER	4.2
1	B	196	THR	4.0
1	B	200	GLY	3.9
1	B	201	ARG	3.6
1	C	193	ALA	3.5
1	A	171	GLU	3.4
1	D	193	ALA	3.4
1	D	191	THR	3.1
1	C	196	THR	2.8
1	D	196	THR	2.8
1	D	198	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	195	ASP	2.6
1	A	196	THR	2.5
1	D	204	ALA	2.5
1	D	236	ASP	2.4
1	D	203	GLY	2.3
1	D	252	HIS	2.2
1	D	233	ARG	2.1
1	B	193	ALA	2.1
1	D	282	ASN	2.1
1	A	252	HIS	2.1
1	C	203	GLY	2.0
1	A	202	SER	2.0
1	C	285	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	U6N	B	401	35/35	0.94	0.16	19,20,23,29	0
2	U6N	C	401	35/35	0.94	0.17	21,24,31,38	0
2	U6N	D	401	35/35	0.94	0.17	18,25,29,33	0
2	U6N	A	401	35/35	0.95	0.15	13,18,23,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.