



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2023 – 03:56 AM EDT

PDB ID : 1NEJ
Title : Crystalline Human Carbonmonoxy Hemoglobin S (Liganded Sickle Cell Hemoglobin) Exhibits The R2 Quaternary State At Neutral pH In The Presence Of Polyethylene Glycol: The 2.1 Angstrom Resolution Crystal Structure
Authors : Patskovska, L.N.; Patskovsky, Y.V.; Almo, S.C.; Hirsch, R.E.
Deposited on : 2002-12-11
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	141	 79% 18% .
1	C	141	 74% 22% .
2	B	146	 80% 17% .
2	D	146	 83% 14% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4793 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemoglobin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1069	685	187	194	3	0	0	0
1	C	141	1069	685	187	194	3	0	0	0

- Molecule 2 is a protein called Hemoglobin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	146	1121	724	195	199	3	0	0	0
2	D	146	1121	724	195	199	3	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	6	VAL	GLU	SEE REMARK 999	UNP P68871
D	6	VAL	GLU	SEE REMARK 999	UNP P68871

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 2 1 1	0	0
4	B	1	Total C O 2 1 1	0	0
4	C	1	Total C O 2 1 1	0	0
4	D	1	Total C O 2 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	56	Total O 56 56	0	0
5	B	53	Total O 53 53	0	0
5	C	57	Total O 57 57	0	0
5	D	67	Total O 67 67	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemoglobin alpha chain




- Molecule 1: Hemoglobin alpha chain



- Molecule 2: Hemoglobin beta chain



- Molecule 2: Hemoglobin beta chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.00Å 58.53Å 171.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.10 29.73 – 2.10	Depositor EDS
% Data completeness (in resolution range)	83.3 (8.00-2.10) 91.3 (29.73-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.10Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.235 , 0.268 0.295 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	40.7	Xtrriage
Anisotropy	0.455	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 86.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.044 for k,h,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4793	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/1097	0.85	0/1491
1	C	0.66	0/1097	0.81	0/1491
2	B	0.65	0/1151	0.77	0/1564
2	D	0.68	0/1151	0.79	2/1564 (0.1%)
All	All	0.65	0/4496	0.80	2/6110 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	104	ARG	NE-CZ-NH1	6.03	123.31	120.30
2	D	104	ARG	NE-CZ-NH2	-5.53	117.53	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	76	MET	Mainchain
2	D	104	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1073	18	0
1	C	1069	0	1073	19	0
2	B	1121	0	1121	19	0
2	D	1121	0	1121	15	0
3	A	43	0	30	2	0
3	B	43	0	30	1	0
3	C	43	0	30	4	0
3	D	43	0	30	2	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
5	A	56	0	0	1	0
5	B	53	0	0	0	0
5	C	57	0	0	3	0
5	D	67	0	0	2	0
All	All	4793	0	4508	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:16:LYS:HE3	5:C:323:HOH:O	1.60	1.02
2:B:47:ASP:O	2:B:48:LEU:HD12	1.67	0.94
3:A:142:HEM:HHA	3:A:142:HEM:HBD1	1.55	0.87
3:C:142:HEM:HBD2	5:C:354:HOH:O	1.76	0.85
3:C:142:HEM:HHA	3:C:142:HEM:HBD1	1.67	0.76
1:A:66:LEU:O	1:A:70:VAL:HG23	1.88	0.73
2:D:52:ASP:HB3	5:D:395:HOH:O	1.91	0.70
2:D:106:LEU:HD12	3:D:147:HEM:HAB	1.76	0.68
1:C:66:LEU:O	1:C:70:VAL:HG23	1.94	0.66
1:A:89:HIS:HB2	1:A:141:ARG:HH21	1.61	0.66
2:B:106:LEU:HD12	3:B:147:HEM:HAB	1.78	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:46:PHE:HA	1:C:54:GLN:OE1	1.98	0.63
1:C:28:ALA:HB1	1:C:105:LEU:HD22	1.80	0.62
2:D:82:LYS:HD3	2:D:82:LYS:H	1.65	0.61
1:A:28:ALA:HB1	1:A:105:LEU:HD22	1.82	0.60
2:B:47:ASP:C	2:B:48:LEU:HD12	2.23	0.58
1:C:89:HIS:HB2	1:C:141:ARG:HH21	1.70	0.57
1:C:89:HIS:HB2	1:C:141:ARG:NH2	2.21	0.56
1:C:70:VAL:HG22	1:C:128:PHE:CZ	2.41	0.56
2:B:50:THR:O	2:B:54:VAL:HG23	2.06	0.56
2:B:31:LEU:HD22	2:B:106:LEU:HD23	1.88	0.54
1:A:27:GLU:OE2	1:A:108:THR:HG23	2.08	0.53
1:A:42:TYR:HE2	1:A:97:ASN:ND2	2.07	0.52
2:D:3:LEU:HD11	2:D:78:LEU:HD13	1.92	0.52
1:A:42:TYR:CZ	1:A:93:VAL:HG13	2.46	0.51
2:D:3:LEU:HD12	2:D:3:LEU:H	1.75	0.50
1:A:42:TYR:CE2	1:A:97:ASN:ND2	2.80	0.50
1:C:72:HIS:ND1	1:C:72:HIS:N	2.60	0.50
1:A:72:HIS:N	1:A:72:HIS:ND1	2.60	0.49
1:C:83:LEU:HD21	3:C:142:HEM:HMA3	1.94	0.49
2:D:51:PRO:O	2:D:55:MET:HG2	2.13	0.49
2:B:82:LYS:HD3	2:B:82:LYS:H	1.78	0.48
2:B:3:LEU:HD11	2:B:78:LEU:HD13	1.94	0.48
1:C:43:PHE:HB3	1:C:46:PHE:HB2	1.95	0.48
3:C:142:HEM:HHA	3:C:142:HEM:CBD	2.42	0.48
1:A:29:LEU:HD22	1:A:101:LEU:HD11	1.95	0.48
2:D:29:GLY:O	2:D:33:VAL:HG23	2.14	0.47
2:D:137:VAL:O	2:D:141:LEU:HB2	2.14	0.47
2:B:35:TYR:HB3	2:B:37:TRP:CZ2	2.48	0.47
1:C:49:SER:HA	5:C:333:HOH:O	2.16	0.46
2:D:50:THR:O	2:D:54:VAL:HG23	2.15	0.46
1:A:43:PHE:HB3	1:A:46:PHE:HB2	1.96	0.46
2:D:1:VAL:CG2	5:D:306:HOH:O	2.63	0.46
3:A:142:HEM:HBD1	3:A:142:HEM:CHA	2.34	0.46
1:A:122:HIS:HD2	2:B:30:ARG:HH11	1.64	0.45
1:C:29:LEU:HD22	1:C:101:LEU:HD11	2.00	0.44
1:A:77:PRO:HG3	1:C:1:VAL:HG21	1.99	0.44
1:C:76:MET:SD	1:C:132:VAL:HG22	2.58	0.44
2:B:30:ARG:O	2:B:34:VAL:HG23	2.18	0.44
2:D:88:LEU:HD11	3:D:147:HEM:HMA1	1.99	0.44
1:A:28:ALA:CB	1:A:105:LEU:HD22	2.46	0.43
2:D:18:VAL:HG22	2:D:23:VAL:HG21	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:24:TYR:CE1	1:C:112:HIS:ND1	2.83	0.43
1:C:84:SER:O	1:C:88:ALA:HB3	2.18	0.43
2:B:129:ALA:O	2:B:133:VAL:HG23	2.18	0.43
1:A:78:ASN:HD22	1:A:78:ASN:C	2.21	0.43
1:A:118:THR:OG1	1:A:121:VAL:HG23	2.17	0.43
2:B:18:VAL:HG22	2:B:23:VAL:HG21	2.00	0.43
2:B:39:GLN:HG2	2:B:48:LEU:HD23	2.02	0.42
2:D:82:LYS:H	2:D:82:LYS:CD	2.26	0.42
1:A:30:GLU:HG3	5:A:224:HOH:O	2.18	0.42
1:A:42:TYR:OH	1:A:93:VAL:HA	2.20	0.42
1:C:107:VAL:HG13	2:D:115:ALA:HB2	2.02	0.42
2:B:111:VAL:HG11	2:B:131:GLN:OE1	2.20	0.41
1:C:98:PHE:HE2	1:C:136:LEU:HD12	1.84	0.41
2:B:51:PRO:O	2:B:55:MET:HG2	2.20	0.41
2:B:57:ASN:HA	2:B:58:PRO:HD2	1.96	0.41
1:C:117:PHE:CZ	1:C:122:HIS:CD2	3.09	0.41
2:D:92:HIS:HA	2:D:96:LEU:HB2	2.03	0.41
2:B:137:VAL:O	2:B:141:LEU:HB2	2.20	0.41
1:A:98:PHE:HE2	1:A:136:LEU:HD12	1.85	0.40
2:B:35:TYR:CZ	2:B:105:LEU:HD22	2.56	0.40
2:B:82:LYS:H	2:B:82:LYS:CD	2.32	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	129 (93%)	10 (7%)	0	100	100
1	C	139/141 (99%)	125 (90%)	14 (10%)	0	100	100
2	B	144/146 (99%)	133 (92%)	9 (6%)	2 (1%)	11	6
2	D	144/146 (99%)	135 (94%)	7 (5%)	2 (1%)	11	6

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	566/574 (99%)	522 (92%)	40 (7%)	4 (1%)	22 18

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	20	VAL
2	B	47	ASP
2	D	20	VAL
2	D	47	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	113/113 (100%)	102 (90%)	11 (10%)	8 5
1	C	113/113 (100%)	101 (89%)	12 (11%)	6 4
2	B	118/118 (100%)	114 (97%)	4 (3%)	37 39
2	D	118/118 (100%)	111 (94%)	7 (6%)	19 17
All	All	462/462 (100%)	428 (93%)	34 (7%)	13 10

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LEU
1	A	50	HIS
1	A	72	HIS
1	A	73	VAL
1	A	74	ASP
1	A	78	ASN
1	A	86	LEU
1	A	90	LYS
1	A	105	LEU
1	A	122	HIS
1	A	140	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	3	LEU
2	B	5	PRO
2	B	82	LYS
2	B	141	LEU
1	C	9	ASN
1	C	29	LEU
1	C	34	LEU
1	C	72	HIS
1	C	73	VAL
1	C	74	ASP
1	C	78	ASN
1	C	85	ASP
1	C	86	LEU
1	C	89	HIS
1	C	90	LYS
1	C	105	LEU
2	D	3	LEU
2	D	5	PRO
2	D	18	VAL
2	D	48	LEU
2	D	82	LYS
2	D	104	ARG
2	D	141	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	68	ASN
1	A	78	ASN
1	A	122	HIS
2	B	39	GLN
2	B	80	ASN
1	C	9	ASN
1	C	78	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	C	142	4,1	41,50,50	1.54	6 (14%)	45,82,82	1.56	6 (13%)
4	CMO	B	2148	3	0,1,1	-	-	-	-	-
3	HEM	D	147	2,4	41,50,50	1.53	6 (14%)	45,82,82	1.23	4 (8%)
4	CMO	D	4148	3	0,1,1	-	-	-	-	-
4	CMO	C	3143	3	0,1,1	-	-	-	-	-
3	HEM	A	142	4,1	41,50,50	1.50	7 (17%)	45,82,82	1.31	3 (6%)
4	CMO	A	1143	3	0,1,1	-	-	-	-	-
3	HEM	B	147	2,4	41,50,50	1.33	5 (12%)	45,82,82	1.21	5 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	142	4,1	-	10/12/54/54	-
3	HEM	D	147	2,4	-	7/12/54/54	-
3	HEM	B	147	2,4	-	7/12/54/54	-
3	HEM	C	142	4,1	-	8/12/54/54	-

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	142	HEM	C3C-CAC	-4.87	1.37	1.47
3	A	142	HEM	C3C-CAC	-4.74	1.38	1.47
3	D	147	HEM	C3C-CAC	-4.64	1.38	1.47
3	D	147	HEM	C3C-C2C	-3.84	1.35	1.40
3	B	147	HEM	C3C-CAC	-3.58	1.40	1.47
3	C	142	HEM	C3C-C2C	-3.29	1.35	1.40
3	C	142	HEM	CAB-C3B	-3.17	1.38	1.47
3	A	142	HEM	CAB-C3B	-3.11	1.39	1.47
3	A	142	HEM	C3C-C2C	-3.11	1.36	1.40
3	B	147	HEM	CAB-C3B	-3.10	1.39	1.47
3	D	147	HEM	CAB-C3B	-3.08	1.39	1.47
3	B	147	HEM	CBB-CAB	3.07	1.45	1.30
3	D	147	HEM	CBB-CAB	3.06	1.45	1.30
3	A	142	HEM	CBB-CAB	2.94	1.44	1.30
3	C	142	HEM	CBB-CAB	2.82	1.44	1.30
3	B	147	HEM	C3C-C2C	-2.79	1.36	1.40
3	C	142	HEM	C1B-NB	-2.58	1.35	1.40
3	B	147	HEM	CBC-CAC	2.58	1.46	1.29
3	A	142	HEM	C1B-NB	-2.33	1.36	1.40
3	D	147	HEM	CBC-CAC	2.33	1.44	1.29
3	A	142	HEM	CHC-C4B	-2.24	1.34	1.41
3	C	142	HEM	CBC-CAC	2.20	1.43	1.29
3	A	142	HEM	CBC-CAC	2.12	1.43	1.29
3	D	147	HEM	C1A-CHA	-2.09	1.35	1.41

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	142	HEM	CAD-C3D-C4D	4.52	132.56	124.66
3	A	142	HEM	CAD-C3D-C4D	3.65	131.03	124.66
3	D	147	HEM	C4C-CHD-C1D	3.57	127.27	122.56
3	C	142	HEM	CMA-C3A-C4A	-3.51	123.08	128.46
3	C	142	HEM	CAD-C3D-C2D	-3.21	121.90	127.88
3	C	142	HEM	CMA-C3A-C2A	3.16	130.90	124.94
3	B	147	HEM	C2C-C3C-C4C	-3.00	104.80	106.90
3	D	147	HEM	CBA-CAA-C2A	2.77	117.34	112.62
3	C	142	HEM	CMC-C2C-C3C	2.67	129.68	124.68
3	B	147	HEM	C4C-CHD-C1D	2.66	126.07	122.56
3	A	142	HEM	CMC-C2C-C3C	2.65	129.64	124.68
3	D	147	HEM	CMC-C2C-C3C	2.44	129.25	124.68
3	B	147	HEM	CMC-C2C-C3C	2.19	128.78	124.68
3	A	142	HEM	CBD-CAD-C3D	2.17	118.65	112.63

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	147	HEM	C3D-C4D-ND	2.12	112.52	110.17
3	B	147	HEM	CMA-C3A-C4A	-2.06	125.30	128.46
3	B	147	HEM	C4B-CHC-C1C	2.04	125.25	122.56
3	C	142	HEM	CHC-C4B-C3B	-2.03	121.46	124.57

There are no chirality outliers.

All (32) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	142	HEM	C2B-C3B-CAB-CBB
3	A	142	HEM	C2D-C3D-CAD-CBD
3	A	142	HEM	C4D-C3D-CAD-CBD
3	B	147	HEM	C2B-C3B-CAB-CBB
3	B	147	HEM	C4B-C3B-CAB-CBB
3	C	142	HEM	C2D-C3D-CAD-CBD
3	C	142	HEM	C4D-C3D-CAD-CBD
3	D	147	HEM	C2B-C3B-CAB-CBB
3	D	147	HEM	C4B-C3B-CAB-CBB
3	D	147	HEM	C3D-CAD-CBD-CGD
3	B	147	HEM	C3D-CAD-CBD-CGD
3	A	142	HEM	C2A-CAA-CBA-CGA
3	C	142	HEM	C2A-CAA-CBA-CGA
3	A	142	HEM	C3D-CAD-CBD-CGD
3	A	142	HEM	C4B-C3B-CAB-CBB
3	C	142	HEM	C3D-CAD-CBD-CGD
3	B	147	HEM	CAA-CBA-CGA-O1A
3	D	147	HEM	CAA-CBA-CGA-O1A
3	A	142	HEM	CAA-CBA-CGA-O2A
3	B	147	HEM	CAA-CBA-CGA-O2A
3	A	142	HEM	CAA-CBA-CGA-O1A
3	D	147	HEM	CAD-CBD-CGD-O2D
3	A	142	HEM	CAD-CBD-CGD-O2D
3	B	147	HEM	CAD-CBD-CGD-O2D
3	C	142	HEM	CAD-CBD-CGD-O2D
3	B	147	HEM	CAD-CBD-CGD-O1D
3	D	147	HEM	CAA-CBA-CGA-O2A
3	D	147	HEM	CAD-CBD-CGD-O1D
3	C	142	HEM	CAD-CBD-CGD-O1D
3	A	142	HEM	CAD-CBD-CGD-O1D
3	C	142	HEM	CAA-CBA-CGA-O1A
3	C	142	HEM	CAA-CBA-CGA-O2A


There are no ring outliers.


4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	142	HEM	4	0
3	D	147	HEM	2	0
3	A	142	HEM	2	0
3	B	147	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEM B 147:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM C 142:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

Unable to reproduce the depositor's R factor - this section is therefore empty.