

# Full wwPDB NMR Structure Validation Report (i)

### Aug 20, 2022 - 08:01 AM EDT

PDB ID	:	1NGL
Title	:	HUMAN NEUTROPHIL GELATINASE-ASSOCIATED LIPOCALIN (HN-
		GAL), REGULARISED AVERAGE NMR STRUCTURE
Authors	:	Coles, M.; Diercks, T.; Muehlenweg, B.; Bartsch, S.; Zoelzer, V.; Tschesche,
		H.; Kessler, H.
Deposited on	:	1999-02-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. $(2010)$
ShiftChecker	:	2.29
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment was not calculated.

There are no overall percentile quality scores available for this entry.

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	А	179	100%



## 2 Ensemble composition and analysis (i)

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.



## 3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2916 atoms, of which 1456 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called PROTEIN (NGAL).

Mol	Chain	Residues			Atom	IS			Trace
1	Δ	179	Total	С	Η	Ν	0	S	0
	Л	119	2916	942	1456	243	269	6	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	1	MET	-	SEE REMARK 999	UNP P80188



## 4 Residue-property plots (i)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

Cl	hε	ii	n	А	: '																					10	20	%																							
M1 CO	2 00	S4	Τ5	S6	D7	α ⊐⊢	- <del></del>		P12	P13	L14 2	S15	K16				120	N22	F23	Q24		N26	$\alpha$		67.h	າຕ		Y33	<b>m</b>	ო	1 27	A38	G39	N40	A41	142	541 R 44	E45	D46	K47	D48	P49	<b>q</b> 50	M52	Υ53	A54	T55	156 V57	E58	L59	K60
E61 D62	707 K63	S64	Y 65	N66	767 #00	1 00 2 69	V70	L71	F72	R73	K74	K75	K76		D/2		181 181	R82	T83	F84	V 85		G87		680	G91	ົດ	6	T94	L95	0690 M07	16N	K99	S100	Y101	P102	2 6	T105	S106			V109	R110	V112	듺	÷.	Ξ.	Y116 N117	井	H119	1
M121 V122	NO	F124	K125	K126	V127	8128 0129	N130	R131	E132	13	F134	13	I136	<u>n</u> 5	29	34	R141	4	4	4		T146	÷.	÷.	L149 К150	101	12	F153	19	ι n	H 7	K158	12	16	G161	L162 D162	2 4	9	H166	6	<u>ں</u> و	ω I	N 1		~	5	212	C176 1177	1	14	

• Molecule 1: PROTEIN (NGAL)



## 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.8
X-PLOR	structure solution	

No chemical shift data was provided.



## 6 Model quality (i)

## 6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	0	0	0	0
All	All	0	0	0	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

## 6.3 Torsion angles (i)

### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mo	l Chair	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	0	-	-	-	-
Al	All	0	-	-	-	-

There are no Ramachandran outliers.



#### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	0	-	-	-
All	All	0	-	-	-

There are no protein residues with a non-rotameric sidechain to report.

#### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.6 Ligand geometry (i)

There are no ligands in this entry.

### 6.7 Other polymers (i)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 7 Chemical shift validation (i)

No chemical shift data were provided

