



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 17, 2022 – 06:31 AM EST

PDB ID : 1NGR  
Title : DEATH DOMAIN OF P75 LOW AFFINITY NEUROTROPHIN RECEPTOR, RESIDUES 334-418, NMR, 20 STRUCTURES  
Authors : Otting, G.; Liepinsh, E.  
Deposited on : 1997-01-28

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.26  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.26


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	85	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:336-A:340, A:344-A:415 (77)	0.36	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters. No single-model clusters were found.

Cluster number	Models
1	2, 9, 11, 15, 20
2	3, 7, 10, 13, 19
3	6, 8, 16, 18
4	1, 4, 14
5	5, 12, 17

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1314 atoms, of which 651 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called P75 LOW AFFINITY NEUROTROPHIN RECEPTOR.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	85	1314	413	651	118	130	2	0

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




#### 4.2.2 Score per residue for model 2

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.3 Score per residue for model 3

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.4 Score per residue for model 4

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.5 Score per residue for model 5

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.6 Score per residue for model 6

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.7 Score per residue for model 7

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.8 Score per residue for model 8

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.9 Score per residue for model 9

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.10 Score per residue for model 10

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.11 Score per residue for model 11

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.12 Score per residue for model 12

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.13 Score per residue for model 13

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.14 Score per residue for model 14

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.15 Score per residue for model 15

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




### 4.2.16 Score per residue for model 16

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR



### 4.2.17 Score per residue for model 17


- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR






#### 4.2.18 Score per residue for model 18

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR




#### 4.2.19 Score per residue for model 19

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR



#### 4.2.20 Score per residue for model 20

- Molecule 1: P75 LOW AFFINITY NEUROTROPHIN RECEPTOR



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *VARIABLE TARGET FUNCTION (DIANA) APPROACH*.

Of the 50 calculated structures, 20 were deposited, based on the following criterion: *LOWEST TARGET FUNCTION AFTER DIANA*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
OPAL	refinement	
EASY	structure solution	
HABAS	structure solution	
DIANA	structure solution	
OPAL	structure solution	

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.59±0.01	0±0/619 ( 0.0± 0.0%)	1.16±0.04	2±1/842 ( 0.2± 0.2%)
All	All	0.59	0/12380 ( 0.0%)	1.16	31/16840 ( 0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.1±0.2	1.9±1.0
All	All	1	39

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	337	TYR	CB-CG-CD2	-9.74	115.16	121.00	17	8
1	A	413	GLU	CB-CA-C	8.61	127.62	110.40	2	6
1	A	345	ARG	CD-NE-CZ	8.18	135.04	123.60	2	1
1	A	404	ARG	NE-CZ-NH1	6.48	123.54	120.30	14	1
1	A	408	ARG	NE-CZ-NH2	-6.47	117.06	120.30	13	2
1	A	345	ARG	NE-CZ-NH2	-6.00	117.30	120.30	2	2
1	A	366	TYR	CB-CG-CD2	-5.87	117.48	121.00	13	3
1	A	345	ARG	NE-CZ-NH1	5.83	123.22	120.30	2	1
1	A	382	ARG	NE-CZ-NH1	5.82	123.21	120.30	8	2
1	A	345	ARG	CA-CB-CG	5.57	125.65	113.40	8	1
1	A	406	ILE	CA-CB-CG1	5.49	121.43	111.00	18	1
1	A	382	ARG	NE-CZ-NH2	-5.49	117.56	120.30	20	1
1	A	337	TYR	CB-CG-CD1	5.33	124.20	121.00	17	1
1	A	408	ARG	NE-CZ-NH1	5.13	122.87	120.30	13	1

All unique chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	413	GLU	CA	1

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	366	TYR	Sidechain	9
1	A	382	ARG	Sidechain	8
1	A	337	TYR	Sidechain	6
1	A	405	ARG	Sidechain	5
1	A	408	ARG	Sidechain	4
1	A	345	ARG	Sidechain	4
1	A	373	SER	Mainchain	2
1	A	404	ARG	Sidechain	1

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	607	599	599	3±2
All	All	12140	11980	11980	56

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:351:LEU:HB3	1:A:411:ILE:HG23	0.71	1.62	9	14
1:A:364:LEU:HD22	1:A:388:TRP:CZ2	0.52	2.39	15	3
1:A:396:LEU:HD11	1:A:415:LEU:HD22	0.52	1.82	1	3
1:A:337:TYR:OH	1:A:415:LEU:HD21	0.51	2.05	19	4
1:A:351:LEU:CB	1:A:411:ILE:HG23	0.50	2.37	3	1
1:A:364:LEU:HD21	1:A:399:LEU:HA	0.49	1.84	15	1
1:A:352:LEU:HD21	1:A:403:LEU:HD11	0.48	1.84	16	2
1:A:356:THR:HG21	1:A:411:ILE:HD12	0.48	1.85	18	2

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:352:LEU:HD23	1:A:411:ILE:HD13	0.46	1.88	16	1
1:A:388:TRP:CH2	1:A:394:ALA:HA	0.46	2.45	16	2
1:A:388:TRP:HH2	1:A:398:ALA:HB3	0.46	1.71	4	6
1:A:337:TYR:CE2	1:A:385:LEU:HD13	0.44	2.47	2	1
1:A:356:THR:HG21	1:A:411:ILE:CD1	0.44	2.42	18	1
1:A:337:TYR:CD1	1:A:385:LEU:HD22	0.43	2.47	1	1
1:A:337:TYR:CE1	1:A:385:LEU:HD22	0.43	2.49	18	2
1:A:400:LEU:HG	1:A:412:VAL:HG13	0.43	1.89	19	1
1:A:337:TYR:OH	1:A:348:VAL:HG21	0.42	2.14	10	1
1:A:344:LYS:O	1:A:348:VAL:HG23	0.42	2.14	10	1
1:A:337:TYR:CD2	1:A:399:LEU:HD22	0.41	2.50	8	1
1:A:352:LEU:CD2	1:A:403:LEU:HD21	0.41	2.45	12	2
1:A:364:LEU:HD22	1:A:388:TRP:CE2	0.41	2.50	15	1
1:A:352:LEU:HD11	1:A:399:LEU:HD21	0.41	1.90	14	1
1:A:352:LEU:HD23	1:A:411:ILE:CD1	0.41	2.46	16	1
1:A:357:TRP:CG	1:A:358:ARG:N	0.40	2.89	8	2
1:A:352:LEU:CD2	1:A:411:ILE:HG21	0.40	2.46	15	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	77/85 (91%)	62±2 (81±2%)	12±2 (15±2%)	3±1 (4±1%)	5	32
All	All	1540/1700 (91%)	1247 (81%)	233 (15%)	60 (4%)	5	32

All 8 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	380	PRO	20
1	A	355	ASP	18
1	A	393	SER	10
1	A	394	ALA	4
1	A	354	GLY	4

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	392	ASP	2
1	A	407	GLN	1
1	A	377	GLU	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	63/70 (90%)	54±2 (86±4%)	9±2 (14±4%)	<b>7</b> 47
All	All	1260/1400 (90%)	1088 (86%)	172 (14%)	<b>7</b> 47

All 30 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	396	LEU	16
1	A	406	ILE	15
1	A	372	ASP	13
1	A	397	ASP	13
1	A	336	LEU	12
1	A	340	LEU	10
1	A	345	ARG	10
1	A	407	GLN	10
1	A	400	LEU	9
1	A	353	ASN	8
1	A	413	GLU	7
1	A	337	TYR	6
1	A	404	ARG	5
1	A	347	GLU	4
1	A	350	LYS	4
1	A	382	ARG	4
1	A	367	GLN	4
1	A	405	ARG	3
1	A	408	ARG	3
1	A	374	PHE	2
1	A	359	HIS	2
1	A	376	HIS	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	363	GLU	2
1	A	358	ARG	2
1	A	355	ASP	1
1	A	373	SER	1
1	A	377	GLU	1
1	A	339	SER	1
1	A	415	LEU	1
1	A	370	HIS	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided