



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2020 – 07:32 PM BST

PDB ID : 4NGP
Title : Crystal Structure of Glutamate Carboxypeptidase II in a complex with urea-based inhibitor
Authors : Tykvar, J.; Pachl, P.
Deposited on : 2013-11-02
Resolution : 1.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	739	
2	B	2	
2	C	2	
2	F	2	
3	D	5	
4	E	3	

2 Entry composition i

There are 10 unique types of molecules in this entry. The entry contains 6474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	695	5593	3605	937	1032	19	0	13	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	ARG	-	expression tag	UNP Q04609
A	13	SER	-	expression tag	UNP Q04609
A	14	GLY	-	expression tag	UNP Q04609
A	15	LEU	-	expression tag	UNP Q04609
A	16	ASN	-	expression tag	UNP Q04609
A	17	ASP	-	expression tag	UNP Q04609
A	18	ILE	-	expression tag	UNP Q04609
A	19	PHE	-	expression tag	UNP Q04609
A	20	GLU	-	expression tag	UNP Q04609
A	21	ALA	-	expression tag	UNP Q04609
A	22	GLN	-	expression tag	UNP Q04609
A	23	LYS	-	expression tag	UNP Q04609
A	24	ILE	-	expression tag	UNP Q04609
A	25	GLU	-	expression tag	UNP Q04609
A	26	TRP	-	expression tag	UNP Q04609
A	27	HIS	-	expression tag	UNP Q04609
A	28	GLU	-	expression tag	UNP Q04609
A	29	GLY	-	expression tag	UNP Q04609
A	30	SER	-	expression tag	UNP Q04609
A	31	GLY	-	expression tag	UNP Q04609
A	32	SER	-	expression tag	UNP Q04609
A	33	GLY	-	expression tag	UNP Q04609
A	34	SER	-	expression tag	UNP Q04609
A	35	GLU	-	expression tag	UNP Q04609
A	36	ASN	-	expression tag	UNP Q04609
A	37	LEU	-	expression tag	UNP Q04609
A	38	TYR	-	expression tag	UNP Q04609

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	39	PHE	-	expression tag	UNP Q04609
A	40	GLN	-	expression tag	UNP Q04609
A	41	GLY	-	expression tag	UNP Q04609
A	42	ARG	-	expression tag	UNP Q04609
A	43	SER	-	expression tag	UNP Q04609

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0
2	C	2	28	16	2	10	0	0	0
2	F	2	28	16	2	10	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	5	61	34	2	25	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	E	3	39	22	2	15	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
5	A	1	14	8	1	5	0	0

- Molecule 6 is N-[[[(1S)-1-carboxy-5-[[2-[(4,4-dioxo-4H-(3aS,4S,6aR)-2-oxohexahydro-1H-tieno[3,4-d]imidazol-4-yl]-7,10,13,16,19,22,25,28,31,34,37,40-dodecaoxa-3,43-diazaoctatetracont-1-yl]oxy)benzoyl]amino}pentyl]carbonyl]-L-glutamic acid (three-letter code: J31) (formula: $C_{58}H_{97}N_7O_{24}S$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	42	26	4	12	0	0

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
7	A	2	2	2	0	0

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
8	A	1	1	1	0	0

- Molecule 9 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
9	A	1	1	1	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
10	A	620	623	623	0	3

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1
MAG2
BMA3
MAN4
MAN5

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2
BMA3

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	102.07Å 130.41Å 160.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.35 – 1.63 45.57 – 1.63	Depositor EDS
% Data completeness (in resolution range)	98.3 (47.35-1.63) 98.3 (45.57-1.63)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 1.63Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.178 , 0.205 0.185 , 0.208	Depositor DCC
R_{free} test set	6475 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtrriage
Anisotropy	0.050	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6474	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG, CL, CA, J31, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.02	4/5777 (0.1%)	1.01	11/7831 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	437	GLU	CD-OE2	-5.89	1.19	1.25
1	A	434	TRP	CE3-CZ3	5.77	1.48	1.38
1	A	183	GLU	CD-OE1	5.64	1.31	1.25
1	A	557	GLU	CD-OE2	-5.25	1.19	1.25

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	370	ARG	NE-CZ-NH1	9.82	125.21	120.30
1	A	370	ARG	NE-CZ-NH2	-7.97	116.32	120.30
1	A	387	ASP	CB-CG-OD1	-7.92	111.17	118.30
1	A	673	ARG	NE-CZ-NH2	-7.54	116.53	120.30
1	A	673	ARG	NE-CZ-NH1	7.36	123.98	120.30
1	A	465	ASP	CB-CG-OD1	7.17	124.75	118.30
1	A	309	MET	CG-SD-CE	-6.20	90.28	100.20
1	A	536	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	A	63	ASP	CB-CG-OD1	5.13	122.92	118.30
1	A	369	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	666	ASP	CB-CG-OD1	5.04	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5593	0	5460	28	0
2	B	28	0	25	0	0
2	C	28	0	25	0	0
2	F	28	0	25	0	0
3	D	61	0	52	0	0
4	E	39	0	34	0	0
5	A	28	0	26	0	0
6	A	42	0	34	0	0
7	A	2	0	0	0	0
8	A	1	0	0	0	0
9	A	1	0	0	0	0
10	A	623	0	0	10	0
All	All	6474	0	5681	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:443:GLN:HG2	10:A:1516:HOH:O	1.73	0.87
1:A:91:GLN:HG2	10:A:1499:HOH:O	1.77	0.85
1:A:58:MET:HE1	1:A:586:GLU:HG2	1.71	0.72
1:A:215:LYS:HE2	10:A:1437:HOH:O	1.93	0.67
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.30	0.66
1:A:688[B]:ARG:NH1	10:A:1139:HOH:O	2.29	0.65
1:A:651:GLN:O	1:A:655:LYS:NZ	2.33	0.60
1:A:659[B]:ILE:O	1:A:663[B]:MET:HG3	2.04	0.58
1:A:506:PHE:O	1:A:507:SER:O	2.22	0.57
1:A:58:MET:CE	1:A:586:GLU:HG2	2.35	0.55
1:A:506:PHE:CB	1:A:509:MET:HG3	2.37	0.54
1:A:718:LYS:HE3	10:A:995:HOH:O	2.08	0.53
1:A:169:MET:HA	1:A:344:MET:O	2.08	0.52
1:A:189:GLU:HG3	10:A:1299:HOH:O	2.10	0.51
1:A:325:VAL:HB	1:A:326:PRO:HD2	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:543:THR:HG22	1:A:543:THR:O	2.13	0.48
1:A:641:GLU:HG3	10:A:1418:HOH:O	2.13	0.48
1:A:169:MET:HE1	10:A:943:HOH:O	2.17	0.45
1:A:551:LEU:HD22	1:A:556:TYR:HB2	1.99	0.43
1:A:591[A]:ILE:HD12	10:A:931:HOH:O	2.18	0.43
1:A:507:SER:C	1:A:509:MET:H	2.23	0.42
1:A:657[A]:ASN:CG	1:A:660[A]:VAL:HG23	2.40	0.42
1:A:227:LEU:O	1:A:296:PRO:HA	2.20	0.41
1:A:539:LYS:HE2	10:A:1128:HOH:O	2.20	0.41
1:A:225:VAL:O	1:A:294:VAL:HA	2.21	0.41
1:A:449:TYR:O	1:A:532:SER:HA	2.21	0.40
1:A:387:ASP:HA	1:A:388:PRO:HA	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	703/739 (95%)	682 (97%)	18 (3%)	3 (0%)	34 15

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	335	GLY
1	A	507	SER
1	A	382	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	598/629 (95%)	587 (98%)	11 (2%)	59 34

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	188	LEU
1	A	189	GLU
1	A	303	GLN
1	A	443	GLN
1	A	499	LYS
1	A	507	SER
1	A	519	ASN
1	A	537	TYR
1	A	539	LYS
1	A	600	TYR
1	A	673	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	651	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	1.04	2 (14%)	17,19,21	0.97	1 (5%)
2	NAG	B	2	2	14,14,15	0.54	0	17,19,21	1.12	1 (5%)
2	NAG	C	1	1,2	14,14,15	0.75	0	17,19,21	1.36	3 (17%)
2	NAG	C	2	2	14,14,15	0.65	0	17,19,21	1.03	1 (5%)
3	NAG	D	1	1,3	14,14,15	0.86	0	17,19,21	1.50	3 (17%)
3	NAG	D	2	3	14,14,15	0.79	1 (7%)	17,19,21	1.20	1 (5%)
3	BMA	D	3	3	11,11,12	0.76	0	15,15,17	0.88	1 (6%)
3	MAN	D	4	3	11,11,12	0.78	0	15,15,17	1.21	3 (20%)
3	MAN	D	5	3	11,11,12	0.62	0	15,15,17	1.23	2 (13%)
4	NAG	E	1	1,4	14,14,15	0.95	0	17,19,21	2.38	8 (47%)
4	NAG	E	2	4	14,14,15	0.58	0	17,19,21	1.18	1 (5%)
4	BMA	E	3	4	11,11,12	0.52	0	15,15,17	1.15	1 (6%)
2	NAG	F	1	1,2	14,14,15	0.84	1 (7%)	17,19,21	1.78	5 (29%)
2	NAG	F	2	2	14,14,15	0.65	0	17,19,21	1.91	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	1/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	MAN	D	5	3	-	0/2/19/22	0/1/1/1
4	NAG	E	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	0/6/23/26	0/1/1/1
4	BMA	E	3	4	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	2/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	NAG	O5-C1	-2.36	1.39	1.43
2	B	1	NAG	C1-C2	2.20	1.55	1.52
2	F	1	NAG	C3-C2	2.18	1.57	1.52
2	B	1	NAG	C4-C5	2.05	1.57	1.53

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	1	NAG	O5-C1-C2	-4.99	103.41	111.29
2	F	2	NAG	C1-O5-C5	4.18	117.85	112.19
4	E	1	NAG	O5-C5-C4	-3.89	101.37	110.83
4	E	2	NAG	O5-C5-C6	3.43	112.57	107.20
4	E	1	NAG	O4-C4-C5	-3.38	100.90	109.30
3	D	1	NAG	O5-C1-C2	-3.35	105.99	111.29
4	E	1	NAG	O4-C4-C3	-3.32	102.68	110.35
2	F	1	NAG	O3-C3-C4	-3.26	102.81	110.35
2	F	2	NAG	C6-C5-C4	-3.04	105.88	113.00
2	F	1	NAG	C8-C7-N2	2.99	121.16	116.10
2	F	1	NAG	O3-C3-C2	2.93	115.53	109.47
2	F	2	NAG	O7-C7-C8	-2.91	116.66	122.06
3	D	5	MAN	O5-C5-C6	2.82	111.62	107.20
4	E	1	NAG	C8-C7-N2	2.62	120.53	116.10
4	E	3	BMA	C6-C5-C4	2.61	119.12	113.00
4	E	1	NAG	O6-C6-C5	-2.61	102.35	111.29
3	D	1	NAG	O5-C5-C6	2.60	111.28	107.20
2	F	1	NAG	O5-C1-C2	-2.58	107.21	111.29
2	F	1	NAG	O7-C7-C8	-2.58	117.27	122.06
4	E	1	NAG	C3-C4-C5	-2.55	105.69	110.24
3	D	4	MAN	O5-C5-C6	2.52	111.16	107.20
4	E	1	NAG	O7-C7-C8	-2.51	117.39	122.06
3	D	2	NAG	O4-C4-C5	-2.45	103.21	109.30
2	C	1	NAG	C2-N2-C7	-2.35	119.56	122.90
2	C	1	NAG	O4-C4-C3	-2.29	105.06	110.35
2	C	2	NAG	O3-C3-C2	-2.20	104.92	109.47
3	D	3	BMA	O3-C3-C2	-2.15	105.88	109.99
3	D	5	MAN	O2-C2-C1	-2.14	104.77	109.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	4	MAN	O3-C3-C4	-2.11	105.47	110.35
2	B	2	NAG	O5-C5-C6	2.10	110.50	107.20
2	F	2	NAG	O5-C5-C6	2.09	110.47	107.20
2	F	2	NAG	O5-C1-C2	-2.07	108.02	111.29
3	D	4	MAN	C1-O5-C5	2.06	114.99	112.19
3	D	1	NAG	C6-C5-C4	-2.06	108.19	113.00
2	B	1	NAG	O3-C3-C2	-2.05	105.22	109.47
2	C	1	NAG	C4-C3-C2	2.02	113.98	111.02

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	E	3	BMA	C4-C5-C6-O6
4	E	3	BMA	O5-C5-C6-O6
2	F	2	NAG	C4-C5-C6-O6
2	F	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	J31	A	817	7	33,42,92	0.91	2 (6%)	36,52,109	1.43	5 (13%)
5	NAG	A	801	1	14,14,15	0.60	0	17,19,21	1.73	1 (5%)
5	NAG	A	806	1	14,14,15	0.96	1 (7%)	17,19,21	1.88	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	J31	A	817	7	-	6/36/46/110	0/1/1/3
5	NAG	A	801	1	-	2/6/23/26	0/1/1/1
5	NAG	A	806	1	-	2/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	817	J31	C5-N1	-3.00	1.42	1.46
6	A	817	J31	C9-C8	2.19	1.56	1.53
5	A	806	NAG	C2-N2	-2.07	1.42	1.46

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	801	NAG	C1-O5-C5	5.21	119.25	112.19
5	A	806	NAG	O5-C5-C6	5.03	115.08	107.20
6	A	817	J31	C4-C5-N1	-4.82	103.18	110.19
6	A	817	J31	C18-C13-C	-3.09	120.53	126.16
5	A	806	NAG	C1-C2-N2	-2.72	105.84	110.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	806	NAG	O5-C5-C4	-2.44	104.88	110.83
6	A	817	J31	O-C-C13	2.33	125.27	121.01
5	A	806	NAG	C3-C4-C5	-2.31	106.11	110.24
6	A	817	J31	N2-C7-N1	2.30	118.48	115.25
5	A	806	NAG	O3-C3-C2	-2.23	104.86	109.47
6	A	817	J31	C9-C8-N2	2.07	113.22	110.19

There are no chirality outliers.


All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	817	J31	N-C-C13-C18
6	A	817	J31	C12-C8-N2-C7
6	A	817	J31	O-C-C13-C18
5	A	806	NAG	O5-C5-C6-O6
5	A	801	NAG	C4-C5-C6-O6
5	A	801	NAG	O5-C5-C6-O6
5	A	806	NAG	C4-C5-C6-O6
6	A	817	J31	O9-C24-C25-O10
6	A	817	J31	N-C-C13-C14
6	A	817	J31	C25-C24-O9-C23

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	695/739 (94%)	0.07	21 (3%) 50 48	11, 21, 41, 65	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	507	SER	4.6
1	A	506	PHE	3.9
1	A	336	ASN	3.5
1	A	487	GLY	3.3
1	A	337	PHE	3.3
1	A	508	GLY	3.1
1	A	334	THR	3.0
1	A	153	ASN	2.9
1	A	135	GLY	2.8
1	A	719	VAL	2.8
1	A	134	ASP	2.8
1	A	505	GLU	2.8
1	A	136	ASN	2.6
1	A	131	ILE	2.3
1	A	55	LYS	2.3
1	A	124	HIS	2.3
1	A	155	SER	2.3
1	A	490	GLY	2.2
1	A	521	PHE	2.2
1	A	335	GLY	2.1
1	A	333	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BMA	E	3	11/12	0.65	0.30	59,63,66,70	0
2	NAG	F	2	14/15	0.77	0.32	53,57,64,65	0
3	MAN	D	5	11/12	0.77	0.23	46,51,59,66	0
2	NAG	C	2	14/15	0.78	0.28	35,43,50,57	0
2	NAG	B	1	14/15	0.85	0.14	33,37,41,49	0
2	NAG	F	1	14/15	0.87	0.11	26,38,42,48	0
2	NAG	B	2	14/15	0.89	0.25	48,52,58,67	0
3	MAN	D	4	11/12	0.89	0.14	41,44,48,51	0
3	BMA	D	3	11/12	0.91	0.11	37,40,44,45	0
4	NAG	E	1	14/15	0.91	0.09	24,26,29,37	0
3	NAG	D	2	14/15	0.92	0.14	31,36,45,48	0
2	NAG	C	1	14/15	0.93	0.12	26,32,37,43	0
3	NAG	D	1	14/15	0.93	0.08	18,24,32,43	0
4	NAG	E	2	14/15	0.94	0.12	30,37,44,50	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain C:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	806	14/15	0.80	0.24	38,48,65,67	0
5	NAG	A	801	14/15	0.85	0.22	53,59,63,63	0
6	J31	A	817	42/90	0.95	0.14	12,28,54,60	0
8	CL	A	820	1/1	0.99	0.11	15,15,15,15	0
7	ZN	A	818	1/1	1.00	0.08	13,13,13,13	0
9	CA	A	821	1/1	1.00	0.08	12,12,12,12	0
7	ZN	A	819	1/1	1.00	0.08	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.