



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2021 – 12:10 pm BST

PDB ID : 7NIN  
Title : X-ray crystal structure of LsAA9A - CinnamtanninB1 soak  
Authors : Frandsen, K.E.H.; Tokin, R.; Skov, L.; Johansen, K.S.; Lo Leggio, L.  
Deposited on : 2021-02-12  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 2372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Auxiliary activity 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	235	1917	1207	328	379	3	0	19	0

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cu		
2	A	1	1	1	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	6	Total Cl 6 6	0	1

- Molecule 5 is Cinnamtannin B1 (three-letter code: UFK) (formula: C<sub>45</sub>H<sub>36</sub>O<sub>18</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 63 45 18	0	0

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	338	Total O 347 347	0	18

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Auxiliary activity 9



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	125.04Å 125.04Å 125.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.25 – 1.40 44.21 – 1.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (44.25-1.40) 100.0 (44.21-1.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.12 (at 1.40Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.174 , 0.192 0.178 , 0.196	Depositor DCC
$R_{free}$ test set	3292 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.7	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 47.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	2372	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CL, HIC, NAG, UFK, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.84	0/1969	0.97	3/2712 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	85	TYR	CB-CG-CD1	6.27	124.76	121.00
1	A	23	ARG	NE-CZ-NH1	5.33	122.97	120.30
1	A	191	TYR	CB-CG-CD1	5.15	124.09	121.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1917	0	1802	21	0
2	A	1	0	0	0	0
3	A	14	0	13	0	0
4	A	6	0	0	0	0
5	A	63	0	0	0	0
6	A	24	0	18	0	0
7	A	347	0	0	3	0
All	All	2372	0	1833	21	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177[B]:ASP:O	1:A:178[B]:SER:C	2.20	0.75
1:A:176[B]:ASP:OD1	1:A:178[B]:SER:N	2.20	0.74
1:A:91[B]:SER:O	1:A:92[B]:ASN:HB2	1.88	0.73
1:A:96[B]:ASN:HB3	1:A:184[B]:GLY:O	1.94	0.68
1:A:91[B]:SER:O	1:A:92[B]:ASN:CB	2.44	0.64
1:A:177[B]:ASP:C	1:A:179[B]:THR:N	2.57	0.54
1:A:91[B]:SER:OG	1:A:97[B]:VAL:HG11	2.11	0.51
1:A:178[B]:SER:HB3	1:A:179[B]:THR:HG23	1.93	0.50
1:A:131:HIS:CD2	1:A:222:LEU:HD11	2.47	0.50
1:A:182[A]:PRO:HG3	1:A:218:TRP:NE1	2.27	0.49
1:A:96[B]:ASN:ND2	7:A:407:HOH:O	2.44	0.49
1:A:177[B]:ASP:O	1:A:179[B]:THR:N	2.45	0.49
1:A:183[B]:ALA:HA	7:A:668[B]:HOH:O	2.14	0.47
1:A:177[B]:ASP:C	1:A:179[B]:THR:H	2.20	0.44
1:A:50[B]:LYS:HE2	1:A:50[B]:LYS:HB3	1.60	0.44
1:A:48:VAL:HB	1:A:49:PRO:HD2	2.00	0.43
1:A:8:TRP:CZ3	1:A:230:GLY:HA3	2.54	0.42
1:A:80:GLY:HA3	1:A:144:ILE:O	2.19	0.42
1:A:176[B]:ASP:OD1	1:A:178[B]:SER:HB2	2.20	0.41
1:A:182[A]:PRO:HD3	1:A:218:TRP:CZ2	2.56	0.41
1:A:117:ARG:HD2	7:A:571:HOH:O	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	252/235 (107%)	229 (91%)	22 (9%)	1 (0%)	34 12

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	42	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	210/194 (108%)	208 (99%)	2 (1%)	<a href="#">76</a> <a href="#">53</a>

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	ARG
1	A	209	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	HIC	A	1	1,2	8,11,12	0.97	1 (12%)	6,14,16	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HIC	A	1	1,2	-	0/5/6/8	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	HIC	CE1-ND1	-2.56	1.30	1.35

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 7 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	UFK	A	309	-	71,72,72	2.07	19 (26%)	94,114,114	1.63	16 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	ACT	A	311	-	1,3,3	3.43	1 (100%)	0,3,3	0.00	-
6	ACT	A	315	-	1,3,3	3.27	1 (100%)	0,3,3	0.00	-
6	ACT	A	310	-	1,3,3	4.97	1 (100%)	0,3,3	0.00	-
6	ACT	A	312	-	1,3,3	2.19	1 (100%)	0,3,3	0.00	-
6	ACT	A	313	-	1,3,3	3.41	1 (100%)	0,3,3	0.00	-
3	NAG	A	302	1	14,14,15	0.69	0	17,19,21	1.36	2 (11%)
6	ACT	A	314	-	1,3,3	3.63	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	UFK	A	309	-	-	0/18/74/74	0/11/10/10
3	NAG	A	302	1	-	0/6/23/26	0/1/1/1

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	309	UFK	CBU-CBW	-6.83	1.42	1.51
5	A	309	UFK	CBG-CAW	-5.91	1.41	1.51
5	A	309	UFK	CBR-CBS	-5.86	1.43	1.52
5	A	309	UFK	CAP-CAS	-5.59	1.46	1.53
6	A	310	ACT	CH3-C	4.97	1.55	1.48
5	A	309	UFK	CBB-CBS	-4.30	1.45	1.52
6	A	314	ACT	CH3-C	3.63	1.53	1.48
6	A	311	ACT	CH3-C	3.43	1.53	1.48
6	A	313	ACT	CH3-C	3.41	1.53	1.48
5	A	309	UFK	CAE-CAG	-3.28	1.45	1.51
6	A	315	ACT	CH3-C	3.27	1.52	1.48
5	A	309	UFK	CAT-CAS	-3.23	1.47	1.52
5	A	309	UFK	CAH-CAG	3.13	1.58	1.52
5	A	309	UFK	CAX-CAS	-2.98	1.50	1.55
5	A	309	UFK	OCC-CBZ	-2.73	1.30	1.36
5	A	309	UFK	CAI-CAJ	-2.40	1.47	1.51
5	A	309	UFK	OCD-CBY	-2.38	1.31	1.36
5	A	309	UFK	CBX-CBY	2.36	1.42	1.38
5	A	309	UFK	CCA-CBZ	-2.34	1.35	1.39
5	A	309	UFK	CBX-CBW	-2.28	1.35	1.39
5	A	309	UFK	OCK-CBT	2.21	1.47	1.42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	312	ACT	CH3-C	2.19	1.51	1.48
5	A	309	UFK	OAL-CAG	2.17	1.48	1.45
5	A	309	UFK	CAK-CAP	2.07	1.43	1.39
5	A	309	UFK	OAQ-CAH	2.02	1.47	1.43

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	309	UFK	CBB-CBS-CBR	5.59	118.83	110.14
5	A	309	UFK	CAD-CAC-CAB	-4.49	115.89	120.50
5	A	309	UFK	OAV-CAW-CAX	4.10	118.72	110.26
5	A	309	UFK	CAT-CAU-CBB	3.70	126.16	119.45
3	A	302	NAG	O5-C1-C2	-3.44	105.86	111.29
5	A	309	UFK	CAI-CAH-CAG	3.36	113.59	109.09
5	A	309	UFK	CBU-OBP-CBA	3.29	124.67	116.62
5	A	309	UFK	CBY-CBX-CBW	-2.83	117.57	120.76
5	A	309	UFK	CBH-CBG-CAW	-2.80	113.93	119.83
5	A	309	UFK	CAK-OAL-CAG	2.75	122.32	115.39
5	A	309	UFK	CAF-CAA-CAB	2.73	122.27	119.86
3	A	302	NAG	O3-C3-C4	-2.71	104.08	110.35
5	A	309	UFK	CAY-CAT-CAU	-2.53	113.46	116.43
5	A	309	UFK	CBL-CBG-CAW	2.40	125.11	120.64
5	A	309	UFK	CAC-CAD-CAE	2.28	123.50	121.20
5	A	309	UFK	CAN-CAO-CAP	2.13	124.10	121.48
5	A	309	UFK	CBK-CBL-CBG	-2.01	119.17	121.20
5	A	309	UFK	CBU-CBT-CBS	-2.01	105.65	110.72

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	234/235 (99%)	0.24	11 (4%) 31 31	18, 23, 38, 57	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	108	VAL	4.6
1	A	109	THR	3.7
1	A	110	ASN	3.6
1	A	111	SER	3.2
1	A	203	TYR	3.1
1	A	235	GLU	2.6
1	A	208	THR	2.5
1	A	211	VAL	2.2
1	A	112	THR	2.1
1	A	107	ASN	2.0
1	A	96[A]	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	HIC	A	1	11/12	0.93	0.07	19,21,23,28	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	ACT	A	314	4/4	0.54	0.25	38,40,41,47	4
6	ACT	A	311	4/4	0.72	0.20	65,66,68,70	0
6	ACT	A	315	4/4	0.75	0.22	48,49,54,55	0
4	CL	A	304	1/1	0.77	0.14	68,68,68,68	1
6	ACT	A	312	4/4	0.77	0.17	36,37,38,45	4
6	ACT	A	313	4/4	0.81	0.18	49,50,52,62	0
3	NAG	A	302	14/15	0.84	0.13	26,31,35,36	0
4	CL	A	308	1/1	0.85	0.14	52,52,52,52	0
5	UFK	A	309	63/63	0.86	0.12	21,28,48,54	0
4	CL	A	303[A]	1/1	0.88	0.14	33,33,33,33	1
6	ACT	A	310	4/4	0.92	0.16	42,44,50,56	0
4	CL	A	305	1/1	0.98	0.05	30,30,30,30	0
4	CL	A	306	1/1	0.99	0.04	26,26,26,26	0
4	CL	A	307	1/1	0.99	0.05	31,31,31,31	0
2	CU	A	301	1/1	1.00	0.04	22,22,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.