



Full wwPDB X-ray Structure Validation Report i

Mar 24, 2022 – 12:36 pm GMT

PDB ID : 5NLM
Title : Complex between a UDP-glucosyltransferase from *Polygonum tinctorium* capable of glucosylating indoxyl and indoxyl sulfate
Authors : Welner, D.H.; Hsu, T.; Dueber, J.; Adams, P.D.
Deposited on : 2017-04-04
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.27
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)


There are 4 unique types of molecules in this entry. The entry contains 14195 atoms, of which 7064 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called indoxyl UDP-glucosyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	459	7111	2268	3567	607	655	14	0	1	0
1	B	448	6935	2210	3485	593	634	13	0	0	0

- Molecule 2 is 3-SULFOOXY-1H-INDOLE (three-letter code: IOS) (formula: C₈H₇NO₄S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	S		
2	A	1	20	8	6	1	4	1	0	0
2	B	1	20	8	6	1	4	1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	4	Total Mg 4 4	0	0
3	B	1	Total Mg 1 1	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	90	Total O 90 90	0	0
4	B	14	Total O 14 14	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: indoxyl UDP-glucosyltransferase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	121.00 Å 172.82 Å 48.41 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.41 – 2.14 49.56 – 2.14	Depositor EDS
% Data completeness (in resolution range)	98.0 (48.41-2.14) 98.0 (49.56-2.14)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.31 (at 2.14 Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R , R_{free}	0.225 , 0.252 0.225 , 0.252	Depositor DCC
R_{free} test set	2780 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	42.5	Xtriage
Anisotropy	0.486	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14195	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOS, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3632	0.41	0/4933
1	B	0.24	0/3533	0.41	0/4800
All	All	0.24	0/7165	0.41	0/9733

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3544	3567	3569	5	0
1	B	3450	3485	3485	22	0
2	A	14	6	7	0	0
2	B	14	6	7	1	0
3	A	4	0	0	0	0
3	B	1	0	0	0	0
4	A	90	0	0	1	0
4	B	14	0	0	0	0
All	All	7131	7064	7068	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:293:GLU:OE2	1:B:421:ARG:NH1	2.20	0.74
1:B:40:ARG:NH2	1:B:474:GLU:OE1	2.24	0.69
1:B:451:ALA:O	1:B:455:VAL:HG23	1.94	0.68
1:B:460:GLY:O	1:B:462:SER:N	2.33	0.62
1:B:40:ARG:NH1	1:B:467:GLU:OE1	2.35	0.60
1:B:189:ASP:N	1:B:189:ASP:OD1	2.34	0.60
1:B:457:SER:O	1:B:463:THR:OG1	2.12	0.59
1:A:417:ASP:OD2	1:A:421:ARG:NH2	2.38	0.55
1:B:396:LYS:O	1:B:397:MET:HB2	2.08	0.54
1:B:396:LYS:O	1:B:397:MET:CB	2.57	0.52
1:B:145:SER:HB2	2:B:1000:IOS:H5	1.97	0.47
1:B:153:PHE:O	1:B:206:SER:OG	2.28	0.47
1:B:341:GLN:O	1:B:342:THR:OG1	2.33	0.45
1:A:77:ASP:O	1:A:97:ARG:NH1	2.50	0.45
1:B:188:ILE:HD11	1:B:202:LEU:HD21	1.98	0.45
1:B:262:GLU:HA	1:B:262:GLU:OE1	2.16	0.45
1:A:260:ARG:N	1:A:261:PRO:CD	2.80	0.45
1:A:310:ARG:NH1	4:A:1104:HOH:O	2.50	0.44
1:B:182:VAL:HG22	1:B:186:ASP:HB2	1.99	0.44
1:B:270:GLN:OE1	1:B:275:VAL:HG22	2.18	0.44
1:B:455:VAL:O	1:B:460:GLY:O	2.34	0.43
1:B:459:ASP:O	1:B:463:THR:OG1	2.35	0.43
1:B:261:PRO:HA	1:B:265:LYS:HE2	2.00	0.43
1:B:417:ASP:OD2	1:B:421:ARG:NH2	2.49	0.43
1:B:455:VAL:HG13	1:B:461:SER:HB3	2.02	0.42
1:B:436:GLU:OE1	1:B:436:GLU:N	2.54	0.41
1:A:188:ILE:O	1:A:191:VAL:HG22	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	454/478 (95%)	436 (96%)	18 (4%)	0	100 100
1	B	440/478 (92%)	414 (94%)	24 (6%)	2 (0%)	29 22
All	All	894/956 (94%)	850 (95%)	42 (5%)	2 (0%)	47 45

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	397	MET
1	B	262	GLU

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	389/399 (98%)	385 (99%)	4 (1%)	76 79
1	B	378/399 (95%)	372 (98%)	6 (2%)	62 65
All	All	767/798 (96%)	757 (99%)	10 (1%)	71 73

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103[A]	ARG
1	A	103[B]	ARG
1	A	321	TYR
1	A	476	LYS
1	B	189	ASP
1	B	260	ARG
1	B	268	ASP
1	B	272	ARG
1	B	345	ARG
1	B	411	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	IOS	B	1000	-	15,15,15	1.69	2 (13%)	14,22,22	1.05	2 (14%)
2	IOS	A	1000	-	15,15,15	1.72	2 (13%)	14,22,22	1.15	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IOS	B	1000	-	-	3/3/5/5	0/2/2/2
2	IOS	A	1000	-	-	0/3/5/5	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	IOS	O1-S	5.04	1.66	1.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1000	IOS	O1-S	4.92	1.65	1.58
2	A	1000	IOS	C1-N1	-2.07	1.32	1.36
2	B	1000	IOS	C1-N1	-2.01	1.32	1.36

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	IOS	O3-S-O2	-2.82	100.91	112.22
2	B	1000	IOS	O4-S-O1	2.26	111.27	105.83
2	B	1000	IOS	O4-S-O2	-2.10	101.21	108.49

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1000	IOS	C2-O1-S-O2
2	B	1000	IOS	C2-O1-S-O4
2	B	1000	IOS	C2-O1-S-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1000	IOS	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	459/478 (96%)	0.77	25 (5%) 25 32	29, 51, 95, 128	0
1	B	448/478 (93%)	1.33	107 (23%) 0 0	48, 89, 132, 175	0
All	All	907/956 (94%)	1.04	132 (14%) 2 3	29, 70, 122, 175	0

All (132) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	260	ARG	8.6
1	A	323	SER	8.5
1	A	328	ILE	8.3
1	B	167	PHE	8.1
1	B	78	LEU	7.5
1	A	324	VAL	7.1
1	A	322	PHE	6.5
1	B	393	ALA	6.2
1	B	191	VAL	5.9
1	B	333	LEU	5.8
1	B	266	TRP	5.7
1	A	325	ASP	5.4
1	B	433	MET	5.4
1	B	436	GLU	5.3
1	A	321	TYR	5.3
1	B	262	GLU	5.2
1	B	182	VAL	5.1
1	B	418	GLY	4.9
1	A	464	ARG	4.6
1	B	163	VAL	4.5
1	B	406	LEU	4.5
1	B	159	LEU	4.4
1	B	269	GLN	4.3
1	A	327	GLU	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	267	LEU	4.2
1	B	174	VAL	4.2
1	B	184	GLY	4.2
1	B	392	TYR	4.1
1	B	180	ILE	4.1
1	B	442	ARG	4.1
1	B	152	PHE	4.0
1	A	330	PRO	4.0
1	B	458	LYS	4.0
1	A	326	GLY	3.9
1	B	432	LEU	3.9
1	B	162	THR	3.9
1	B	358	VAL	3.8
1	B	223	LEU	3.7
1	A	256	GLY	3.7
1	B	263	CYS	3.7
1	B	441	ILE	3.6
1	B	410	LEU	3.6
1	B	464	ARG	3.6
1	B	414	VAL	3.5
1	B	183	HIS	3.5
1	B	455	VAL	3.5
1	B	434	GLU	3.4
1	B	456	LEU	3.3
1	B	409	GLY	3.3
1	B	176	ILE	3.3
1	B	448	LEU	3.2
1	B	195	LYS	3.2
1	B	158	LYS	3.2
1	B	270	GLN	3.2
1	B	299	ALA	3.1
1	B	202	LEU	3.1
1	B	209	TYR	3.1
1	B	317	ALA	3.0
1	B	215	VAL	3.0
1	B	304	ARG	3.0
1	B	459	ASP	3.0
1	B	155	HIS	3.0
1	B	294	LEU	3.0
1	B	330	PRO	3.0
1	B	111	ALA	3.0
1	A	180	ILE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	420	ILE	3.0
1	B	305	PHE	2.9
1	B	408	VAL	2.8
1	B	361	HIS	2.8
1	B	431	GLU	2.8
1	A	333	LEU	2.8
1	A	202	LEU	2.7
1	A	11	ALA	2.7
1	B	460	GLY	2.7
1	B	175	GLN	2.7
1	B	279	ASN	2.7
1	B	359	LEU	2.7
1	B	429	ILE	2.6
1	B	58	ARG	2.6
1	A	260	ARG	2.6
1	A	255	LYS	2.6
1	B	307	TRP	2.6
1	B	316	ILE	2.6
1	B	407	ARG	2.6
1	A	468	GLU	2.6
1	B	161	GLU	2.6
1	B	303	GLN	2.6
1	B	315	GLY	2.5
1	B	185	LYS	2.5
1	A	336	GLU	2.5
1	B	109	TYR	2.5
1	B	430	GLY	2.5
1	B	466	LEU	2.5
1	B	259	ALA	2.5
1	B	275	VAL	2.4
1	B	343	ALA	2.4
1	B	156	LEU	2.4
1	B	164	SER	2.4
1	B	250	ALA	2.4
1	A	283	GLY	2.4
1	B	293	GLU	2.4
1	B	200	LYS	2.4
1	B	468	GLU	2.3
1	B	318	ASN	2.3
1	B	187	LEU	2.3
1	B	280	PHE	2.3
1	A	348	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	452	ALA	2.3
1	B	283	GLY	2.2
1	A	96	VAL	2.2
1	B	38	LEU	2.2
1	B	348	VAL	2.2
1	B	425	ILE	2.2
1	B	278	VAL	2.2
1	A	329	ASP	2.1
1	B	413	SER	2.1
1	B	81	ALA	2.1
1	B	123	LEU	2.1
1	B	264	LEU	2.1
1	B	154	LEU	2.1
1	B	306	LEU	2.1
1	B	240	ARG	2.1
1	B	302	GLN	2.1
1	B	394	GLU	2.1
1	B	403	THR	2.1
1	B	389	TRP	2.1
1	B	165	CYS	2.1
1	B	286	LEU	2.0
1	B	397	MET	2.0
1	B	364	THR	2.0
1	A	298	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	A	1004	1/1	-0.02	0.32	100,100,100,100	0
2	IOS	B	1000	14/14	0.63	0.27	153,160,195,195	0
2	IOS	A	1000	14/14	0.65	0.21	113,124,140,141	0
3	MG	A	1003	1/1	0.68	0.13	70,70,70,70	0
3	MG	B	1001	1/1	0.73	0.23	52,52,52,52	1
3	MG	A	1001	1/1	0.74	0.21	88,88,88,88	0
3	MG	A	1002	1/1	0.97	0.15	29,29,29,29	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.