

Full wwPDB X-ray Structure Validation Report (i)

May 14, 2020 – 11:17 am BST

PDB ID : 6NL4

Title : Crystal structure of Mycobacterium tuberculosis dethiobiotin synthetase in

complex with gemcitabine

Authors: Thompson, A.P.; Polyak, S.W.; Wegener, K.L.; Bruning, J.B.

Deposited on : 2019-01-07

Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	233	91%	7% •
1	В	233	93%	
1	С	233	90%	8% •
1	D	233	8%	10% • •



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7269 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ATP-dependent dethiobiotin synthetase BioD.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	227	Total	С	N	О	S	0	6	0
1	A	221	1608	1002	292	307	7	0	0	
1	В	226	Total	С	N	О	S	0	3	0
1	Б	220	1589	990	287	305	7	0	3	
1	С	228	Total	С	N	О	S	0	3	0
1		220	1600	997	288	308	7	0	3	
1	D	227	Total	С	N	О	S	0	9	0
1	ש	221	1584	988	284	305	7	U	2	U

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	HIS	-	expression tag	UNP P9WPQ5
A	-5	HIS	-	expression tag	UNP P9WPQ5
A	-4	HIS	-	expression tag	UNP P9WPQ5
A	-3	HIS	_	expression tag	UNP P9WPQ5
A	-2	HIS	-	expression tag	UNP P9WPQ5
A	-1	HIS	-	expression tag	UNP P9WPQ5
A	0	GLY	_	expression tag	UNP P9WPQ5
A	1	GLY	-	expression tag	UNP P9WPQ5
В	-6	HIS	-	expression tag	UNP P9WPQ5
В	-5	HIS	-	expression tag	UNP P9WPQ5
В	-4	HIS	-	expression tag	UNP P9WPQ5
В	-3	HIS	-	expression tag	UNP P9WPQ5
В	-2	HIS	-	expression tag	UNP P9WPQ5
В	-1	HIS	-	expression tag	UNP P9WPQ5
В	0	GLY	-	expression tag	UNP P9WPQ5
В	1	GLY	=	expression tag	UNP P9WPQ5
С	-6	HIS	-	expression tag	UNP P9WPQ5
С	-5	HIS	-	expression tag	UNP P9WPQ5
С	-4	HIS	-	expression tag	UNP P9WPQ5
С	-3	HIS	-	expression tag	UNP P9WPQ5
С	-2	HIS	-	expression tag	UNP P9WPQ5



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	-1	HIS	-	expression tag	UNP P9WPQ5
С	0	GLY	-	expression tag	UNP P9WPQ5
С	1	GLY	-	expression tag	UNP P9WPQ5
D	-6	HIS	-	expression tag	UNP P9WPQ5
D	-5	HIS	-	expression tag	UNP P9WPQ5
D	-4	HIS	-	expression tag	UNP P9WPQ5
D	-3	HIS	-	expression tag	UNP P9WPQ5
D	-2	HIS	-	expression tag	UNP P9WPQ5
D	-1	HIS	-	expression tag	UNP P9WPQ5
D	0	GLY	-	expression tag	UNP P9WPQ5
D	1	GLY	_	expression tag	UNP P9WPQ5


• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	В	1	Total O S 5 4 1	0	0
2	С	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

• Molecule 3 is GEMCITABINE (three-letter code: GEO) (formula: $C_9H_{11}F_2N_3O_4$) (labeled as "Ligand of Interest" by author).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	Λ	1	Total	С	F	N	О	0	0
'	Λ	1	18	9	2	3	4	0	0
2	D	1	Total	С	F	N	О	0	0
)	Ъ	1	18	9	2	3	4	0	0
3	D	1	Total	С	F	N	О	0	0
J	ש	1	18	9	2	3	4		U

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	225	Total O 225 225	0	0
4	В	232	Total O 232 232	0	0
4	С	211	Total O 211 211	0	0
4	D	146	Total O 146 146	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP-dependent dethiobiotin synthetase BioD





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	55.51Å 106.08Å 151.95Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.35 - 1.99	Depositor
resolution (A)	38.35 - 1.99	EDS
% Data completeness	99.4 (38.35-1.99)	Depositor
(in resolution range)	99.5 (38.35-1.99)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.23 (at 2.00Å)	Xtriage
Refinement program	PHENIX	Depositor
P. P.	0.197 , 0.252	Depositor
R, R_{free}	0.197 , 0.252	DCC
R_{free} test set	3092 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 55.2	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7269	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GEO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ # Z > 5		RMSZ	# Z > 5	
1	A	0.38	0/1627	0.57	0/2225	
1	В	0.38	0/1608	0.59	0/2199	
1	С	0.36	0/1619	0.56	0/2215	
1	D	0.36	0/1603	0.57	0/2195	
All	All	0.37	0/6457	0.57	0/8834	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1608	0	1649	8	0
1	В	1589	0	1633	8	0
1	С	1600	0	1635	13	0
1	D	1584	0	1620	15	0
2	A	5	0	0	0	0
2	В	5	0	0	0	0
2	С	5	0	0	0	0
2	D	5	0	0	0	0
3	A	18	0	11	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	18	0	10	1	0
3	D	18	0	10	3	0
4	A	225	0	0	3	0
4	В	232	0	0	2	0
4	С	211	0	0	3	0
4	D	146	0	0	0	0
All	All	7269	0	6568	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		$\operatorname{distance}\ (ext{\AA})$	overlap $(ext{Å})$
3:D:302:GEO:H6	3:D:302:GEO:H5'2	1.78	0.65
1:C:219:ASN:ND2	4:C:405:HOH:O	2.31	0.61
1:B:175:PRO:O	4:B:401:HOH:O	2.16	0.61
1:C:31:ILE:HD11	1:C:222:ALA:HA	1.83	0.61
3:A:302:GEO:H6	3:A:302:GEO:O5'	2.04	0.58
1:A:150:LYS:HE2	1:A:189[B]:ILE:HG21	1.88	0.54
1:C:11:THR:HG21	1:D:143:LEU:HD13	1.90	0.54
1:D:88:ARG:HB2	1:D:122:VAL:HG21	1.89	0.54
1:C:88:ARG:NH1	1:C:130[B]:ASP:OD1	2.40	0.54
1:A:15:LYS:NZ	4:A:405:HOH:O	2.28	0.53
1:C:29:ALA:HB3	1:C:31:ILE:HD12	1.92	0.50
1:D:169:GLY:O	3:D:302:GEO:H5	2.12	0.49
1:B:177:LEU:N	4:B:409:HOH:O	2.46	0.49
1:A:27:ARG:NH1	1:A:57:ALA:O	2.47	0.48
1:D:35:VAL:HG21	1:D:53:VAL:HG11	1.97	0.47
1:D:112:GLY:H	1:D:115:VAL:CG2	2.27	0.46
1:C:65:LEU:HB2	1:C:94:LEU:HD22	1.96	0.46
1:C:43:THR:OG1	1:C:69:PRO:HA	2.16	0.46
1:B:15:LYS:HB3	1:B:15:LYS:HE2	1.70	0.45
1:A:188:ARG:NH1	4:A:410:HOH:O	2.43	0.45
1:D:78:ALA:HB1	1:D:83:MET:O	2.17	0.45
1:D:43:THR:HG21	1:D:67:ARG:HB3	2.00	0.44
1:B:13:VAL:HG13	1:B:138:VAL:HG12	2.00	0.44
1:C:226:GLY:O	4:C:402:HOH:O	2.21	0.43
1:A:44:ALA:HB2	1:A:69:PRO:HB3	2.00	0.43
1:D:52:GLU:HG3	1:D:55:ARG:HH11	1.83	0.43
1:C:68:TYR:HA	1:C:69:PRO:HD3	1.83	0.43



Continued from previous page...

A tom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:D:70:GLN:HG3	1:D:80:HIS:CD2	2.54	0.43
1:B:169:GLY:O	3:B:302:GEO:H5	2.18	0.43
1:B:91:ILE:HD12	1:B:118:ALA:HB2	2.01	0.43
1:D:13:VAL:HG23	1:D:140:THR:HG22	2.00	0.43
1:D:43:THR:OG1	1:D:69:PRO:HA	2.20	0.42
1:D:13:VAL:HG23	1:D:140:THR:CG2	2.49	0.42
1:A:173:ASP:HA	1:A:174:PRO:HA	1.87	0.42
1:A:31:ILE:O	1:A:33:VAL:HG23	2.20	0.42
1:D:119:GLU:HA	1:D:120:PRO:HA	1.85	0.42
1:C:37:LYS:HE3	1:C:39:VAL:O	2.20	0.41
4:A:576:HOH:O	1:B:80:HIS:HD2	2.03	0.41
1:D:112:GLY:H	1:D:115:VAL:HG21	1.86	0.41
3:D:302:GEO:H6	3:D:302:GEO:C5'	2.50	0.41
1:C:174:PRO:HA	1:C:175:PRO:HD3	1.90	0.41
1:C:173:ASP:HA	1:C:174:PRO:HA	1.96	0.40
1:A:39:VAL:HG11	1:A:74:PRO:HB3	2.02	0.40
1:C:97:ASP:OD2	4:C:401:HOH:O	2.21	0.40
1:B:172:PRO:HG2	1:B:175:PRO:HA	2.03	0.40
1:D:173[B]:ASP:HA	1:D:174:PRO:HA	1.80	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	231/233 (99%)	227 (98%)	4 (2%)	0	100	100
1	В	227/233 (97%)	225 (99%)	2 (1%)	0	100	100
1	С	229/233 (98%)	223 (97%)	6 (3%)	0	100	100
1	D	227/233 (97%)	225 (99%)	1 (0%)	1 (0%)	34	30
All	All	914/932 (98%)	900 (98%)	13 (1%)	1 (0%)	51	49



All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	120	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percen	tiles
1	A	155/157~(99%)	153 (99%)	2 (1%)	69	74
1	В	154/157 (98%)	152 (99%)	2 (1%)	69	74
1	С	153/157 (98%)	153 (100%)	0	100	100
1	D	152/157 (97%)	146 (96%)	6 (4%)	32	30
All	All	614/628 (98%)	604 (98%)	10 (2%)	60	67

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	VAL
1	A	218	ARG
1	В	13	VAL
1	В	177	LEU
1	D	11	THR
1	D	71	PRO
1	D	119	GLU
1	D	143	LEU
1	D	153	LEU
1	D	177	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	T	Chain	Dag	Link	Bond lengths			Bond angles		
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GEO	D	302	-	15,19,19	3.85	6 (40%)	16,29,29	2.03	5 (31%)
3	GEO	В	302	-	15,19,19	3.76	6 (40%)	16,29,29	2.07	3 (18%)
2	SO4	В	301	-	4,4,4	0.16	0	6,6,6	0.12	0
2	SO4	D	301	-	4,4,4	0.16	0	6,6,6	0.10	0
2	SO4	A	301	-	4,4,4	0.29	0	6,6,6	0.26	0
2	SO4	С	301	-	4,4,4	0.18	0	6,6,6	0.21	0
3	GEO	A	302	-	15,19,19	3.91	6 (40%)	16,29,29	2.25	5 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GEO	D	302	_	-	2/3/26/26	0/2/2/2
3	GEO	В	302	-	-	1/3/26/26	0/2/2/2
3	GEO	A	302	_	-	1/3/26/26	0/2/2/2

All (18) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	${ m Observed}(m \AA)$	$\operatorname{Ideal}(\text{\AA})$
3	A	302	GEO	O4'-C1'	11.82	1.62	1.42
3	D	302	GEO	O4'-C1'	11.48	1.62	1.42
3	В	302	GEO	O4'-C1'	10.92	1.61	1.42
3	В	302	GEO	O4'-C4'	-7.13	1.29	1.45
3	D	302	GEO	O4'-C4'	-6.95	1.29	1.45
3	A	302	GEO	O4'-C4'	-6.85	1.29	1.45
3	A	302	GEO	C4-N4	3.67	1.46	1.35
3	D	302	GEO	C4-N4	3.52	1.45	1.35
3	В	302	GEO	C4-N4	3.25	1.44	1.35
3	В	302	GEO	C2-N3	-3.20	1.31	1.38
3	В	302	GEO	O3'-C3'	-3.13	1.36	1.42
3	D	302	GEO	O3'-C3'	-3.07	1.36	1.42
3	A	302	GEO	O3'-C3'	-2.80	1.37	1.42
3	D	302	GEO	C4'-C3'	2.78	1.58	1.53
3	A	302	GEO	C4'-C3'	2.77	1.58	1.53
3	A	302	GEO	C2-N3	-2.66	1.32	1.38
3	D	302	GEO	C2-N3	-2.42	1.33	1.38
3	В	302	GEO	C4'-C3'	2.30	1.57	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
3	A	302	GEO	F2-C2'-F1	5.45	111.42	105.20
3	В	302	GEO	C2-N3-C4	4.78	121.19	116.34
3	A	302	GEO	C2-N3-C4	4.59	120.99	116.34
3	D	302	GEO	C2-N3-C4	4.53	120.93	116.34
3	В	302	GEO	F2-C2'-F1	4.52	110.37	105.20
3	D	302	GEO	F2-C2'-F1	4.20	110.00	105.20
3	A	302	GEO	C5'-C4'-C3'	-3.44	108.27	114.90
3	В	302	GEO	C5'-C4'-C3'	-3.39	108.36	114.90
3	D	302	GEO	N4-C4-N3	3.06	121.32	116.49
3	A	302	GEO	N4-C4-N3	2.88	121.04	116.49
3	D	302	GEO	C5'-C4'-C3'	-2.48	110.11	114.90
3	A	302	GEO	C5-C4-N3	-2.36	119.00	121.72
3	D	302	GEO	C5-C4-N3	-2.00	119.41	121.72

There are no chirality outliers.

All (4) torsion outliers are listed below:

	Mol	Chain	Res	Type	Atoms
	3	D	302	GEO	O4'-C4'-C5'-O5'
ľ	3	D	302	GEO	C3'-C4'-C5'-O5'



Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	302	GEO	O4'-C4'-C5'-O5'
3	В	302	GEO	C3'-C4'-C5'-O5'


There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	302	GEO	3	0
3	В	302	GEO	1	0
3	A	302	GEO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	227/233 (97%)	0.09	3 (1%) 77 76	17, 28, 43, 62	0
1	В	$226/233 \ (96\%)$	0.02	0 100 100	22, 31, 45, 84	0
1	С	228/233 (97%)	0.34	12 (5%) 26 25	23, 35, 57, 95	0
1	D	227/233 (97%)	0.60	18 (7%) 12 11	22, 37, 64, 83	2 (0%)
All	All	908/932 (97%)	0.26	33 (3%) 42 42	17, 32, 56, 95	2 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	-1	HIS	3.9
1	D	214	ALA	3.8
1	С	174	PRO	3.5
1	D	44	ALA	3.5
1	С	178	VAL	3.4
1	A	218	ARG	3.4
1	С	176	GLY	3.2
1	С	209	ALA	3.2
1	D	225	VAL	3.2
1	С	218	ARG	3.0
1	D	171	TRP	3.0
1	С	173	ASP	2.7
1	D	205	ALA	2.7
1	D	177	LEU	2.6
1	D	180	ALA	2.6
1	D	219	ASN	2.5
1	D	174	PRO	2.5
1	D	210	ALA	2.5
1	D	0	GLY	2.4
1	D	181	SER	2.3
1	D	223	GLY	2.3



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	177	LEU	2.3
1	D	217	ASP	2.2
1	A	111	GLY	2.2
1	С	184	SER	2.2
1	A	213	ALA	2.2
1	С	171	TRP	2.1
1	С	208	PHE	2.1
1	С	216	PHE	2.1
1	D	172	PRO	2.1
1	С	179	ALA	2.0
1	D	30	GLY	2.0
1	D	221	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	\mathbf{Res}	Atoms	RSCC	RSR	${f B\text{-factors}}({f A}^2)$	Q<0.9
3	GEO	В	302	18/18	0.88	0.21	19,38,41,45	18
3	GEO	D	302	18/18	0.91	0.33	48,76,85,86	0
3	GEO	A	302	18/18	0.91	0.15	23,37,49,51	18
2	SO4	D	301	5/5	0.98	0.12	43,47,52,78	0
2	SO4	A	301	5/5	0.99	0.07	29,31,34,43	0
2	SO4	С	301	5/5	0.99	0.09	34,39,43,45	0
2	SO4	В	301	5/5	0.99	0.08	30,32,35,36	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GEO B 302:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













Electron density around GEO A 302: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)












Electron density around SO4 C 301: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)





6.5 Other polymers (i)

There are no such residues in this entry.

