



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 28, 2022 – 04:37 PM EDT

PDB ID : 6NMG  
Title : Crystal Structure of Rat Ric-8A G alpha binding domain  
Authors : Zeng, B.; Mou, T.C.; Sprang, S.R.  
Deposited on : 2019-01-10  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.29  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	 5% 72% 18% • 9%
1	B	453	 5% 80% 14% • 5%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6873 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Resistance to inhibitors of cholinesterase 8 homolog A (C. elegans).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	413	Total 3271	C 2070	N 587	O 596	S 18	0	0	0
1	B	430	Total 3400	C 2150	N 612	O 619	S 19	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP B1H241
A	232	PHE	TYR	engineered mutation	UNP B1H241
B	0	GLY	-	expression tag	UNP B1H241
B	232	PHE	TYR	engineered mutation	UNP B1H241

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	70	Total O 70 70	0	0
3	B	112	Total O 112 112	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.05Å 103.64Å 141.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.05 – 2.20 31.29 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (31.05-2.20) 99.4 (31.29-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.34 (at 2.20Å)	Xtrriage
Refinement program	PHENIX (1.13rc2_2975: ???)	Depositor
R, $R_{free}$	0.227 , 0.275 0.227 , 0.275	Depositor DCC
$R_{free}$ test set	1505 reflections (2.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.8	Xtrriage
Anisotropy	0.261	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6873	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/3321	0.43	0/4487
1	B	0.29	0/3456	0.45	0/4676
All	All	0.28	0/6777	0.44	0/9163

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3271	0	3352	58	0
1	B	3400	0	3490	39	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
3	A	70	0	0	4	0
3	B	112	0	0	1	0
All	All	6873	0	6842	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:295:HIS:CD2	1:A:298:SER:HB3	2.14	0.82
1:A:283:LEU:HB2	1:A:339:GLU:HG2	1.62	0.81
1:A:295:HIS:HD2	1:A:298:SER:HB3	1.43	0.81
1:A:2:GLU:OE2	1:A:39:GLN:NE2	2.20	0.74
1:B:284:LYS:HA	1:B:343:MET:HE1	1.72	0.71
1:A:325:ARG:HB3	1:B:342:ARG:HD2	1.73	0.70
1:B:419:LEU:HD22	1:B:423:GLY:HA3	1.74	0.69
1:A:115:LEU:HB2	1:A:160:VAL:HG12	1.74	0.69
1:A:44:LYS:HE3	1:A:45:ARG:HH21	1.58	0.68
1:B:36:ASP:O	1:B:43:ARG:NH2	2.27	0.68
1:A:367:GLU:N	1:A:367:GLU:OE1	2.28	0.65
1:A:237:ARG:HD2	1:A:283:LEU:HG	1.77	0.65
1:B:424:LEU:O	1:B:424:LEU:HD12	1.96	0.64
1:A:129:PRO:HB3	1:B:299:LEU:HD13	1.80	0.64
1:A:294:LEU:CD2	1:A:300:GLU:OE1	2.46	0.63
1:B:412:TYR:HD2	1:B:429:ARG:HB3	1.66	0.61
1:A:300:GLU:OE2	1:A:303:GLY:CA	2.49	0.60
1:A:300:GLU:OE2	1:A:303:GLY:HA2	2.01	0.60
1:A:352:LYS:NZ	1:A:356:LEU:O	2.34	0.59
1:A:289:LEU:O	1:A:308:VAL:HG21	2.03	0.58
1:A:360:ARG:NH1	3:A:609:HOH:O	2.36	0.58
1:B:177:ASP:N	1:B:177:ASP:OD1	2.38	0.56
1:A:43:ARG:NH1	1:A:76:ASP:OD2	2.39	0.56
1:A:389:ARG:NH1	3:A:607:HOH:O	2.33	0.56
1:A:344:HIS:HB3	1:A:347:ALA:HB3	1.87	0.55
1:A:300:GLU:OE2	1:A:303:GLY:N	2.40	0.55
1:B:101:SER:OG	1:B:103:GLU:O	2.24	0.55
1:A:294:LEU:HD21	1:A:300:GLU:OE1	2.06	0.55
1:B:406:PHE:O	1:B:410:THR:HG22	2.07	0.54
1:A:295:HIS:NE2	1:A:307:ASP:OD2	2.39	0.54
1:B:150:LEU:HD22	1:B:153:LYS:HE3	1.91	0.53
1:A:326:LEU:HD12	1:B:401:GLU:HG3	1.91	0.53
1:A:294:LEU:HD22	1:A:300:GLU:OE1	2.09	0.52
1:B:154:ARG:NH2	3:B:609:HOH:O	2.33	0.51
1:B:203:ALA:O	1:B:205:LYS:N	2.41	0.51
1:B:217:THR:O	1:B:221:MET:HG3	2.10	0.51
1:A:319:ARG:O	1:A:323:THR:HG22	2.11	0.50
1:A:386:ASP:OD1	1:A:389:ARG:NH2	2.36	0.50
1:A:158:HIS:CE1	1:A:215:GLN:HB2	2.46	0.50
1:B:423:GLY:O	1:B:424:LEU:HG	2.11	0.50
1:B:262:ALA:HB3	1:B:267:THR:HA	1.93	0.50

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2:GLU:HB2	1:B:3:PRO:HD2	1.94	0.49
1:B:43:ARG:NH1	1:B:76:ASP:OD2	2.46	0.49
1:B:103:GLU:HG2	1:B:104:PRO:HD2	1.94	0.48
1:A:41:GLU:O	1:A:45:ARG:HG2	2.14	0.48
1:A:86:ARG:NH1	1:A:137:GLU:OE1	2.46	0.48
1:A:384:ASP:OD1	1:A:387:VAL:HG23	2.14	0.47
1:B:373:ARG:NH1	1:B:409:TYR:O	2.47	0.47
1:B:239:VAL:HG21	1:B:282:PRO:HG3	1.97	0.47
1:A:240:ASP:OD1	1:A:241:GLU:N	2.47	0.47
1:A:327:LYS:HD2	1:A:379:LEU:HD21	1.96	0.47
1:A:182:LEU:HD23	1:A:230:ILE:HG21	1.97	0.47
1:A:299:LEU:HG	1:A:306:MET:HB2	1.96	0.47
1:B:403:VAL:HG13	1:B:419:LEU:HD11	1.96	0.47
1:B:105:ILE:HD12	1:B:105:ILE:H	1.81	0.46
1:B:403:VAL:HG22	1:B:419:LEU:HD21	1.98	0.46
1:A:237:ARG:HH22	1:A:284:LYS:HB3	1.81	0.46
1:A:12:THR:HG22	1:A:14:GLU:HG2	1.98	0.46
1:B:98:ILE:HG21	1:B:144:LEU:HD13	1.97	0.46
1:B:218:GLU:HG2	1:B:266:ARG:NH2	2.31	0.46
1:A:366:PRO:O	1:A:374:ASN:ND2	2.48	0.46
1:A:320:LEU:HD13	1:A:330:VAL:HG21	1.98	0.45
1:A:348:ARG:HH21	1:A:349:LYS:HE3	1.82	0.45
1:A:0:GLY:N	3:A:614:HOH:O	2.50	0.45
1:B:341:ALA:O	1:B:348:ARG:HB2	2.17	0.45
1:B:153:LYS:HD2	1:B:154:ARG:HG2	1.99	0.45
1:A:151:TYR:CZ	1:A:161:GLN:HG2	2.53	0.44
1:A:342:ARG:NH2	1:A:393:GLU:OE2	2.51	0.44
1:B:407:ILE:HD11	1:B:412:TYR:HA	2.00	0.43
1:A:71:ARG:NH1	1:A:116:GLU:OE2	2.51	0.43
1:B:144:LEU:HD12	1:B:144:LEU:HA	1.89	0.43
1:A:237:ARG:NH2	1:A:284:LYS:H	2.15	0.43
1:A:290:LEU:HD22	1:A:344:HIS:CD2	2.54	0.43
1:A:340:CYS:HA	1:A:343:MET:HE2	2.00	0.43
1:A:202:VAL:HG11	1:A:256:HIS:CD2	2.54	0.42
1:B:1:MET:HE2	1:B:6:VAL:HG22	2.00	0.42
1:B:299:LEU:HD12	1:B:299:LEU:HA	1.81	0.42
1:A:89:LEU:HD21	1:A:124:LEU:HD13	2.02	0.42
1:A:98:ILE:HG21	1:A:144:LEU:HD13	2.01	0.42
1:A:190:ARG:NH1	3:A:617:HOH:O	2.51	0.42
1:A:376:LEU:HD12	1:A:379:LEU:HD12	2.01	0.42
1:B:237:ARG:HA	1:B:237:ARG:HD2	1.69	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:115:LEU:HB2	1:B:160:VAL:HG12	2.01	0.42
1:A:15:GLU:HA	1:A:18:VAL:HG22	2.01	0.42
1:A:61:LYS:HA	1:A:61:LYS:HD3	1.87	0.42
1:B:10:LEU:HD13	1:B:49:LEU:HB3	2.01	0.42
1:A:299:LEU:HD21	1:A:350:PHE:CZ	2.55	0.41
1:B:356:LEU:HD21	1:B:395:LEU:HD22	2.03	0.41
1:B:179:ARG:HD2	1:B:229:ASN:O	2.20	0.41
1:B:404:PRO:O	1:B:408:LYS:HG2	2.20	0.41
1:A:80:LEU:HG	1:A:124:LEU:HD23	2.03	0.41
1:A:230:ILE:O	1:A:234:SER:OG	2.35	0.40
1:A:64:VAL:HG22	1:A:113:VAL:HG23	2.04	0.40
1:A:320:LEU:HD12	1:A:320:LEU:HA	1.88	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/453 (89%)	392 (97%)	13 (3%)	0	100	100
1	B	428/453 (94%)	415 (97%)	13 (3%)	0	100	100
All	All	833/906 (92%)	807 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/390 (91%)	346 (97%)	10 (3%)	43	56
1	B	370/390 (95%)	356 (96%)	14 (4%)	33	42
All	All	726/780 (93%)	702 (97%)	24 (3%)	38	49

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ASP
1	A	45	ARG
1	A	52	SER
1	A	77	ARG
1	A	87	GLN
1	A	237	ARG
1	A	295	HIS
1	A	319	ARG
1	A	365	ARG
1	A	376	LEU
1	B	74	SER
1	B	85	SER
1	B	101	SER
1	B	103	GLU
1	B	153	LYS
1	B	154	ARG
1	B	191	LEU
1	B	324	HIS
1	B	325	ARG
1	B	328	GLU
1	B	363	ARG
1	B	388	LYS
1	B	402	SER
1	B	422	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	501	-	4,4,4	0.14	0	6,6,6	0.14	0
2	SO4	A	502	-	4,4,4	0.14	0	6,6,6	0.14	0
2	SO4	B	502	-	4,4,4	0.14	0	6,6,6	0.10	0
2	SO4	B	501	-	4,4,4	0.15	0	6,6,6	0.09	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	413/453 (91%)	0.31	24 (5%) 23 22	20, 50, 81, 105	0
1	B	430/453 (94%)	0.11	24 (5%) 24 23	19, 39, 87, 121	0
All	All	843/906 (93%)	0.21	48 (5%) 23 22	19, 43, 81, 121	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	426	ALA	9.3
1	A	0	GLY	7.9
1	A	265	ASP	6.9
1	B	425	MET	6.9
1	A	103	GLU	6.0
1	B	360	ARG	5.7
1	B	420	ALA	5.3
1	A	102	GLU	5.2
1	B	361	ASP	5.0
1	A	101	SER	4.8
1	A	297	GLY	4.8
1	B	424	LEU	4.7
1	B	427	GLY	4.7
1	B	204	PRO	4.6
1	B	429	ARG	4.5
1	B	428	GLY	4.5
1	A	295	HIS	3.8
1	B	412	TYR	3.8
1	A	203	ALA	3.7
1	A	33	PHE	3.7
1	B	238	GLU	3.7
1	B	325	ARG	3.7
1	B	411	GLY	3.5
1	A	293	GLU	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	421	ALA	3.2
1	A	383	LEU	3.1
1	B	324	HIS	2.9
1	B	363	ARG	2.9
1	A	296	GLU	2.9
1	B	416	ALA	2.8
1	B	236	LYS	2.8
1	A	108	PRO	2.7
1	A	237	ARG	2.8
1	B	237	ARG	2.7
1	A	300	GLU	2.7
1	B	265	ASP	2.6
1	A	294	LEU	2.6
1	B	322	GLN	2.5
1	B	383	LEU	2.4
1	A	345	ARG	2.3
1	A	303	GLY	2.3
1	A	238	GLU	2.2
1	A	359	LEU	2.2
1	A	20	GLU	2.2
1	A	304	VAL	2.2
1	B	362	VAL	2.2
1	A	241	GLU	2.1
1	A	301	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	501	5/5	0.85	0.16	81,81,99,102	0
2	SO4	B	502	5/5	0.94	0.12	66,70,81,83	0
2	SO4	A	502	5/5	0.97	0.11	63,71,76,84	0
2	SO4	B	501	5/5	0.98	0.08	52,58,67,70	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.