



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 30, 2021 – 01:12 pm GMT

PDB ID : 7NMJ  
Title : Crystal structure of Polyphosphate Kinase 2 from *Deinococcus radiodurans* in complex with ADP  
Authors : Silva, S.T.N.; Romao, C.  
Deposited on : 2021-02-23  
Resolution : 2.08 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4 (270009), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	95% 5%
1	B	266	97% .
1	C	266	93% 7%
1	D	266	93% 7%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 17757 atoms, of which 8609 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PPK2 domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	266	Total	C	H	N	O	S	0	0	0
			4282	1368	2124	382	405	3			
1	B	266	Total	C	H	N	O	S	0	1	0
			4303	1373	2137	383	406	4			
1	C	266	Total	C	H	N	O	S	0	1	0
			4304	1373	2138	383	406	4			
1	D	266	Total	C	H	N	O	S	0	0	0
			4284	1368	2126	382	405	3			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
2	A	1	Total	C	H	N	O	P	0	0
			38	10	11	5	10	2		
2	A	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
2	B	1	Total	C	H	N	O	P	0	0
			38	10	11	5	10	2		
2	C	1	Total	C	H	N	O	P	0	0
			38	10	11	5	10	2		
2	C	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	D	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			13	3	7	3		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	1	Total O P 5 4 1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total Mg 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	84	Total O 88 88	0	4
6	B	82	Total O 86 86	0	4
6	C	70	Total O 72 72	0	2
6	D	72	Total O 74 74	0	2

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PPK2 domain-containing protein

Chain A:  95% 5%




- Molecule 1: PPK2 domain-containing protein

Chain B:  % 97% .



- Molecule 1: PPK2 domain-containing protein

Chain C:  % 93% 7%



- Molecule 1: PPK2 domain-containing protein

Chain D:  93% 7%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.78Å 78.85Å 112.14Å 90.00° 109.39° 90.00°	Depositor
Resolution (Å)	63.22 – 2.08 63.22 – 2.08	Depositor EDS
% Data completeness (in resolution range)	39.9 (63.22-2.08) 39.9 (63.22-2.08)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.33 (at 2.08Å)	Xtriage
Refinement program	PHENIX 1.19_4092	Depositor
R, $R_{free}$	0.187 , 0.248 0.187 , 0.247	Depositor DCC
$R_{free}$ test set	1708 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.5	Xtriage
Anisotropy	0.347	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	17757	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ADP, PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/2209	0.49	0/2990
1	B	0.25	0/2217	0.49	0/3000
1	C	0.26	0/2217	0.48	0/3000
1	D	0.26	0/2209	0.48	0/2990
All	All	0.26	0/8852	0.49	0/11980

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2158	2124	2125	6	0
1	B	2166	2137	2135	5	0
1	C	2166	2138	2135	14	0
1	D	2158	2126	2125	10	0
2	A	54	23	24	0	0
2	B	27	11	12	0	0
2	C	54	23	24	1	0
2	D	27	12	12	0	0
3	B	12	15	16	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	5	0	0	0	0
5	D	1	0	0	0	0
6	A	88	0	0	0	0
6	B	86	0	0	0	0
6	C	72	0	0	2	0
6	D	74	0	0	1	0
All	All	9148	8609	8608	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:182:ARG:NH2	2:C:302:ADP:O1A	2.22	0.71
1:C:14:VAL:HG22	1:C:19:TRP:CH2	2.30	0.67
1:A:13:ARG:NH2	1:A:212:ASP:OD1	2.30	0.63
1:C:175:GLN:OE1	1:C:200:ARG:NH1	2.40	0.53
1:A:183:LEU:HD21	1:A:192:PHE:CE2	2.44	0.52
1:C:144:ARG:NH1	1:C:147:GLU:OE1	2.41	0.52
1:C:14:VAL:HG13	1:C:19:TRP:CZ3	2.45	0.51
1:D:65:ARG:NH2	1:D:207:ASN:OD1	2.38	0.51
1:C:65:ARG:NH2	1:C:207:ASN:OD1	2.45	0.50
1:B:65:ARG:NH2	1:B:207:ASN:OD1	2.44	0.49
1:C:94:PRO:HG2	1:C:99:LEU:HD21	1.95	0.49
1:D:63:GLN:NE2	1:D:126:GLU:OE2	2.46	0.48
1:C:206:PHE:HA	1:C:209:VAL:HG22	1.93	0.48
1:B:265:ILE:O	1:B:266:HIS:CB	2.62	0.47
1:D:102:ASP:OD1	1:D:105:TRP:HB2	2.16	0.46
1:B:265:ILE:O	1:B:266:HIS:HB2	2.16	0.46
1:D:154:GLU:OE2	6:D:401:HOH:O	2.20	0.46
1:C:94:PRO:CG	1:C:99:LEU:HD21	2.45	0.46
1:B:206:PHE:HA	1:B:209:VAL:HG22	1.98	0.46
1:D:206:PHE:HA	1:D:209:VAL:HG22	1.98	0.46
1:C:148:HIS:ND1	6:C:404:HOH:O	2.36	0.45
1:D:159:ASN:O	1:D:160:ALA:HB3	2.16	0.45
1:C:159:ASN:O	1:C:160:ALA:HB3	2.18	0.43
1:A:158:ASP:OD2	1:C:148:HIS:NE2	2.51	0.43
1:C:257:ASP:OD1	1:C:257:ASP:N	2.50	0.43
1:A:159:ASN:O	1:A:160:ALA:HB3	2.18	0.42
1:C:105:TRP:O	6:C:401:HOH:O	2.22	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:ASN:O	1:B:160:ALA:HB3	2.20	0.42
1:A:189:HIS:O	1:D:258:TYR:OH	2.37	0.41
1:D:182:ARG:NH1	1:D:191:LYS:O	2.53	0.41
1:D:14:VAL:HG12	1:D:19:TRP:HH2	1.86	0.41
1:A:206:PHE:HA	1:A:209:VAL:HG22	2.03	0.41
1:D:63:GLN:O	1:D:167:TYR:HA	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/266 (99%)	257 (97%)	5 (2%)	2 (1%)	19	14
1	B	264/266 (99%)	258 (98%)	6 (2%)	0	100	100
1	C	264/266 (99%)	256 (97%)	7 (3%)	1 (0%)	34	31
1	D	264/266 (99%)	256 (97%)	7 (3%)	1 (0%)	34	31
All	All	1056/1064 (99%)	1027 (97%)	25 (2%)	4 (0%)	34	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	ASP
1	A	122	ARG
1	C	186	PRO
1	D	195	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/228 (100%)	228 (100%)	0	100	100
1	B	229/228 (100%)	229 (100%)	0	100	100
1	C	229/228 (100%)	229 (100%)	0	100	100
1	D	228/228 (100%)	228 (100%)	0	100	100
All	All	914/912 (100%)	914 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.



There are no torsion outliers.



There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	266/266 (100%)	-0.70	0 <a href="#">100</a>   <a href="#">100</a>	11, 20, 48, 83	0
1	B	266/266 (100%)	-0.68	2 (0%) <a href="#">86</a>   <a href="#">87</a>	10, 20, 46, 101	0
1	C	266/266 (100%)	-0.59	3 (1%) <a href="#">80</a>   <a href="#">83</a>	12, 25, 66, 108	0
1	D	266/266 (100%)	-0.63	1 (0%) <a href="#">92</a>   <a href="#">93</a>	11, 25, 65, 95	0
All	All	1064/1064 (100%)	-0.65	6 (0%) <a href="#">89</a>   <a href="#">91</a>	10, 22, 61, 108	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	191	LYS	3.4
1	B	257	ASP	3.3
1	C	194	PRO	3.2
1	D	194	PRO	3.2
1	B	1[A]	MET	2.1
1	C	187	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	303	6/6	0.93	0.10	22,37,47,47	0
2	ADP	C	302	27/27	0.95	0.14	35,71,95,98	0
2	ADP	A	302	27/27	0.95	0.13	31,55,85,86	0
2	ADP	D	301	27/27	0.96	0.09	22,38,73,98	0
2	ADP	C	301	27/27	0.96	0.08	24,38,66,81	0
3	GOL	B	302	6/6	0.97	0.10	13,28,43,43	0
2	ADP	A	301	27/27	0.97	0.08	17,35,62,76	0
4	PO4	D	302	5/5	0.97	0.08	30,36,49,56	0
2	ADP	B	301	27/27	0.98	0.08	9,28,58,71	0
5	MG	D	303	1/1	0.99	0.03	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around ADP A 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ADP D 301:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PO4 D 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.