



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2020 – 08:33 PM BST

PDB ID : 5NNO  
Title : Structure of TbALDH3 complexed with NAD and AN3057 aldehyde  
Authors : Zoltner, M.; Zhang, N.; Horn, D.; Field, M.C.  
Deposited on : 2017-04-10  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	546	 9% 85% 11%
1	B	546	 9% 85% 11%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7941 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	484	3801	2413	662	709	17	0	5	0
1	B	484	3814	2425	661	711	17	0	6	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP C9ZQX6
A	2	SER	-	expression tag	UNP C9ZQX6
A	3	MET	-	expression tag	UNP C9ZQX6
A	4	ALA	-	expression tag	UNP C9ZQX6
A	259	SER	CYS	conflict	UNP C9ZQX6
A	394	ALA	SER	conflict	UNP C9ZQX6
B	1	GLY	-	expression tag	UNP C9ZQX6
B	2	SER	-	expression tag	UNP C9ZQX6
B	3	MET	-	expression tag	UNP C9ZQX6
B	4	ALA	-	expression tag	UNP C9ZQX6
B	259	SER	CYS	conflict	UNP C9ZQX6
B	394	ALA	SER	conflict	UNP C9ZQX6

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	44	21	7	14	2	0	0
2	B	1	44	21	7	14	2	0	0

- Molecule 3 is 4-[(1-oxidanyl-3 {H}-2,1-benzoxaborol-5-yl)oxy]benzaldehyde (three-letter code: 92N) (formula: C<sub>14</sub>H<sub>11</sub>BO<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	B	C			O
3	A	1	19	1	14	4	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	B	C	O		
3	B	1	19	1	14	4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	106	106	106	0	0
4	B	94	94	94	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldehyde dehydrogenase



- Molecule 1: Aldehyde dehydrogenase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.17Å 62.52Å 92.97Å 90.00° 100.08° 90.00°	Depositor
Resolution (Å)	46.56 – 2.50 46.56 – 2.40	Depositor EDS
% Data completeness (in resolution range)	85.9 (46.56-2.50) 89.5 (46.56-2.40)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.97 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.234 , 0.278 0.271 , 0.332	Depositor DCC
$R_{free}$ test set	1820 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.8	Xtriage
Anisotropy	1.998	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 42.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h	Xtriage
Reported twinning fraction	0.862 for H, K, L 0.138 for L, -K, H	Depositor
Outliers	0 of 35943 reflections	Xtriage
$F_o, F_c$ correlation	0.83	EDS
Total number of atoms	7941	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, 92N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/3870	0.80	5/5246 (0.1%)
1	B	0.57	0/3885	0.75	2/5269 (0.0%)
All	All	0.58	0/7755	0.77	7/10515 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	ARG	NE-CZ-NH1	9.82	125.21	120.30
1	A	66	ARG	NE-CZ-NH2	-9.30	115.65	120.30
1	A	283	ARG	NE-CZ-NH1	7.55	124.07	120.30
1	A	283	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	B	283	ARG	NE-CZ-NH2	5.89	123.25	120.30
1	A	44	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	B	283	ARG	NE-CZ-NH1	-5.11	117.75	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3801	0	3880	6	1
1	B	3814	0	3886	10	1

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	44	0	26	1	0
2	B	44	0	26	0	0
3	A	19	0	0	0	0
3	B	19	0	0	0	0
4	A	106	0	0	0	0
4	B	94	0	0	0	0
All	All	7941	0	7818	15	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:67:ARG:NH1	1:B:128:ASN:O	2.33	0.62
1:A:127:TRP:NE1	2:A:601:NAD:O2N	2.32	0.57
1:B:171[B]:VAL:O	1:B:171[B]:VAL:CG2	2.55	0.55
1:A:47:LEU:HB2	1:A:84:VAL:HG11	1.92	0.52
1:B:66:ARG:NH2	1:B:344:THR:O	2.42	0.52
1:B:47:LEU:HB2	1:B:84:VAL:HG11	1.92	0.51
1:B:85[A]:TRP:CE3	1:B:85[A]:TRP:HA	2.48	0.48
1:B:136:GLN:HB3	1:B:137:PRO:HD3	1.97	0.47
1:A:193[B]:ARG:NH1	1:A:193[B]:ARG:HG2	2.31	0.46
1:B:171[B]:VAL:O	1:B:171[B]:VAL:HG23	2.15	0.46
1:B:283:ARG:HG2	1:B:287:LEU:HD12	1.98	0.45
1:B:65:ARG:CZ	1:B:65:ARG:HA	2.47	0.45
1:A:136:GLN:HB3	1:A:137:PRO:HD3	1.98	0.44
1:A:65[A]:ARG:HA	1:A:65[A]:ARG:CZ	2.47	0.43
1:A:419:VAL:HB	1:B:461:ARG:HA	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:GLN:NE2	1:B:41:GLN:NE2[1_545]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	487/546 (89%)	476 (98%)	10 (2%)	1 (0%)	47	68
1	B	488/546 (89%)	475 (97%)	12 (2%)	1 (0%)	47	68
All	All	975/1092 (89%)	951 (98%)	22 (2%)	2 (0%)	47	68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	445	HIS
1	B	445	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	417/459 (91%)	410 (98%)	7 (2%)	60	82
1	B	418/459 (91%)	415 (99%)	3 (1%)	84	94
All	All	835/918 (91%)	825 (99%)	10 (1%)	71	88

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	113	LEU
1	A	151	LYS
1	A	174	ASP
1	A	300	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	445	HIS
1	A	465	LEU
1	A	483	ARG
1	B	151	LYS
1	B	184	SER
1	B	445	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	136	GLN
1	A	239	ASN
1	A	257	GLN
1	A	305	GLN
1	A	306	GLN
1	A	400	ASN
1	B	16	ASN
1	B	136	GLN
1	B	180	ASN
1	B	239	ASN
1	B	257	GLN
1	B	305	GLN
1	B	306	GLN
1	B	359	HIS
1	B	400	ASN
1	B	455	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAD	A	601	-	42,48,48	0.80	1 (2%)	50,73,73	1.44	10 (20%)
2	NAD	B	601	-	42,48,48	0.80	1 (2%)	50,73,73	1.43	7 (14%)
3	92N	A	602	-	21,21,21	0.38	0	27,29,29	1.15	2 (7%)
3	92N	B	602	-	21,21,21	0.47	0	27,29,29	1.10	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	601	-	-	4/26/62/62	0/5/5/5
2	NAD	B	601	-	-	10/26/62/62	0/5/5/5
3	92N	A	602	-	-	0/6/15/15	0/3/3/3
3	92N	B	602	-	-	0/6/15/15	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	NAD	C5A-C4A	2.50	1.47	1.40
2	A	601	NAD	C5A-C4A	2.20	1.46	1.40

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	NAD	PN-O3-PA	-5.04	115.54	132.83
3	A	602	92N	C11-C10-C8	-4.39	119.25	122.09
3	B	602	92N	C11-C10-C8	-4.28	119.31	122.09
2	A	601	NAD	PN-O3-PA	-4.20	118.40	132.83

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	NAD	N3A-C2A-N1A	-4.04	122.37	128.68
2	B	601	NAD	N3A-C2A-N1A	-4.00	122.43	128.68
2	B	601	NAD	C4A-C5A-N7A	-2.92	106.36	109.40
2	A	601	NAD	C1B-N9A-C4A	-2.71	121.88	126.64
2	B	601	NAD	C3N-C7N-N7N	2.57	120.84	117.75
3	A	602	92N	O1-C5-C11	2.52	127.01	119.10
2	A	601	NAD	C3N-C7N-N7N	2.44	120.67	117.75
2	A	601	NAD	C3D-C2D-C1D	2.42	104.62	100.98
2	B	601	NAD	C1B-N9A-C4A	-2.32	122.56	126.64
2	B	601	NAD	C3D-C2D-C1D	2.29	104.43	100.98
2	A	601	NAD	O4D-C4D-C5D	2.25	116.79	109.37
2	A	601	NAD	C4A-C5A-N7A	-2.23	107.08	109.40
3	B	602	92N	O1-C5-C11	2.16	125.87	119.10
2	A	601	NAD	O4D-C1D-C2D	-2.15	103.78	106.93
2	B	601	NAD	C2A-N1A-C6A	2.04	122.24	118.75
2	A	601	NAD	O2A-PA-O1A	2.03	122.28	112.24
2	A	601	NAD	O5D-C5D-C4D	2.00	115.88	108.99

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	601	NAD	C5D-O5D-PN-O3
2	B	601	NAD	C5D-O5D-PN-O1N
2	B	601	NAD	C5D-O5D-PN-O2N
2	B	601	NAD	O4D-C1D-N1N-C6N
2	B	601	NAD	C3B-C4B-C5B-O5B
2	A	601	NAD	O4B-C4B-C5B-O5B
2	A	601	NAD	C3B-C4B-C5B-O5B
2	B	601	NAD	O4B-C4B-C5B-O5B
2	A	601	NAD	O4D-C4D-C5D-O5D
2	B	601	NAD	O4D-C4D-C5D-O5D
2	B	601	NAD	C4D-C5D-O5D-PN
2	A	601	NAD	C3D-C4D-C5D-O5D
2	B	601	NAD	C4B-C5B-O5B-PA
2	B	601	NAD	C3D-C4D-C5D-O5D


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	NAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	484/546 (88%)	0.95	47 (9%) <b>7</b> <b>7</b>	17, 25, 42, 56	0
1	B	484/546 (88%)	0.91	48 (9%) <b>7</b> <b>7</b>	17, 24, 51, 76	0
All	All	968/1092 (88%)	0.93	95 (9%) <b>7</b> <b>7</b>	17, 24, 45, 76	0

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	355	ILE	4.8
1	A	358	THR	4.6
1	B	466	PHE	4.4
1	A	480	ALA	4.0
1	B	480	ALA	4.0
1	B	358	THR	4.0
1	A	471	THR	3.9
1	B	474	PHE	3.8
1	B	486	ASN	3.7
1	B	6	ALA	3.7
1	B	378	ASN	3.6
1	B	377	GLU	3.6
1	A	377	GLU	3.5
1	B	244	ALA	3.5
1	A	6	ALA	3.5
1	B	359	HIS	3.4
1	A	238	ASP	3.4
1	A	26[A]	GLU	3.4
1	A	321	HIS	3.4
1	B	85[A]	TRP	3.4
1	B	478	THR	3.3
1	B	479	THR	3.3
1	B	488	LEU	3.3
1	B	471	THR	3.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	476	PRO	3.2
1	A	297	GLY	3.2
1	A	466	PHE	3.2
1	A	235	THR	3.1
1	B	472	VAL	3.1
1	A	331	GLY	3.1
1	B	376	TYR	3.1
1	A	155	LEU	3.1
1	A	354	ASN	3.0
1	A	305	GLN	3.0
1	B	402	LYS	2.9
1	B	487	SER	2.9
1	B	323	GLN	2.8
1	A	330	LYS	2.8
1	A	332	GLY	2.8
1	B	41	GLN	2.7
1	A	359	HIS	2.7
1	B	327	GLU	2.7
1	B	467	SER	2.7
1	B	356	ASP	2.7
1	B	237	ALA	2.6
1	B	280	ALA	2.6
1	A	234	ASP	2.6
1	B	8	VAL	2.6
1	B	482	SER	2.6
1	B	475	PRO	2.6
1	B	274	VAL	2.6
1	B	375	THR	2.6
1	A	356	ASP	2.6
1	A	188	ALA	2.5
1	A	484	VAL	2.5
1	A	101	THR	2.5
1	B	301	VAL	2.4
1	A	403	ARG	2.4
1	A	363	GLN	2.4
1	B	354	ASN	2.4
1	B	277	ASP	2.4
1	B	7	GLY	2.3
1	B	469	ILE	2.3
1	A	288	GLY	2.3
1	A	474	PHE	2.3
1	A	353	THR	2.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	478	THR	2.3
1	B	406	ARG	2.3
1	B	464	PHE	2.3
1	B	383	LEU	2.3
1	B	156	ALA	2.3
1	B	305	GLN	2.3
1	A	85	TRP	2.3
1	B	484	VAL	2.3
1	A	27	ALA	2.3
1	B	465	LEU	2.2
1	A	104	GLY	2.2
1	B	481	LYS	2.2
1	A	333	LYS	2.2
1	A	376	TYR	2.1
1	A	31	ASP	2.1
1	A	273	SER	2.1
1	A	308	LEU	2.1
1	A	243	VAL	2.1
1	A	475	PRO	2.1
1	A	88	ILE	2.1
1	A	322	PHE	2.1
1	B	273	SER	2.1
1	A	470	ASP	2.0
1	A	102	MET	2.0
1	A	410	SER	2.0
1	A	467	SER	2.0
1	B	403	ARG	2.0
1	A	345	LEU	2.0
1	B	477	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	92N	B	602	19/19	0.74	0.33	21,25,27,27	0
3	92N	A	602	19/19	0.81	0.31	22,28,29,30	0
2	NAD	B	601	44/44	0.85	0.22	13,21,28,32	0
2	NAD	A	601	44/44	0.86	0.23	11,22,27,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around 92N A 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAD B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.