Full wwPDB X-ray Structure Validation Report (i)

Jun 24, 2021 – 01:22 PM BST

PDB ID	:	7NNV
Title	:	Crystal structure of Mycobacterium tuberculosis ArgF in complex with carbamoyl phosphate.
Authors	:	Mendes, V.; Gupta, P.; Burgess, A.; Sebastian-Perez, V.; Cattermole, E.; Meghir, C.; Blundell, T.L.
Deposited on	:	2021-02-25
Resolution	:	1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.20
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.20

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	2% 97% •
1	B	309	3% 92% 5% •
1	C	309	% 95% • •
1	D	309	2% 91% 6% •
1	E	309	% 91% 6% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	309	3% 94%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 15612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ornithine carbamoyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	308	2327	1452	421	448	6	0	1	0
			Total	C	N	O	S			
1	B	298	2248	1406	409	428	5	0	1	0
			Total	C	N	O	S			
1	C	307	2317	1443	421	448	5	0	0	0
			Total	C	N	O	S			
1	D	300	2264	1416	410	433	5	0	1	0
			Total	C	N	O	S			
1	E	299	2261	1416	407	433	5	0	1	0
			Total	C	N	O	S			
1	F	302	2275	1421	412	437	5	0	1	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P9WIT9
A	0	SER	-	expression tag	UNP P9WIT9
A	1	VAL	-	expression tag	UNP P9WIT9
B	-1	GLY	-	expression tag	UNP P9WIT9
B	0	SER	-	expression tag	UNP P9WIT9
B	1	VAL	-	expression tag	UNP P9WIT9
C	-1	GLY	-	expression tag	UNP P9WIT9
C	0	SER	-	expression tag	UNP P9WIT9
C	1	VAL	-	expression tag	UNP P9WIT9
D	-1	GLY	-	expression tag	UNP P9WIT9
D	0	SER	-	expression tag	UNP P9WIT9
D	1	VAL	-	expression tag	UNP P9WIT9
E	-1	GLY	-	expression tag	UNP P9WIT9
E	0	SER	-	expression tag	UNP P9WIT9
E	1	VAL	-	expression tag	UNP P9WIT9
F	-1	GLY	-	expression tag	UNP P9WIT9
F	0	SER	-	expression tag	UNP P9WIT9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	1	VAL	-	expression tag	UNP P9WIT9

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	O	P			
2	A	1	5	4	1		0	0
2	A	1	5	4	1		0	0
2	B	1	5	4	1		0	0
2	C	1	5	4	1		0	0
2	D	1	5	4	1		0	0
2	D	1	5	4	1		0	0
2	E	1	5	4	1		0	0
2	E	1	5	4	1		0	0
2	E	1	5	4	1		0	0

- Molecule 3 is PHOSPHORIC ACID MONO(FORMAMIDE)ESTER (three-letter code: CP) (formula: CH_4NO_5P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	8	1	1	5	1	0	0
3	B	1	8	1	1	5	1	0	0
3	C	1	8	1	1	5	1	0	0
3	D	1	8	1	1	5	1	0	0
3	E	1	8	1	1	5	1	0	0
3	F	1	8	1	1	5	1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	4	2	2	0	0
			Total	C	O		
4	A	1	4	2	2	0	0
			Total	C	O		
4	C	1	4	2	2	0	0
			Total	C	O		
4	D	1	4	2	2	0	0
			Total	C	O		
4	E	1	4	2	2	0	0
			Total	C	O		
4	F	1	4	2	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	308	308	308	0	0
			Total	O		
5	B	260	260	260	0	0
			Total	O		
5	C	299	299	299	0	0
			Total	O		
5	D	311	311	311	0	0
			Total	O		
5	E	311	311	311	0	0
			Total	O		
5	F	314	314	314	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ornithine carbamoyltransferase

Chain A:





- Molecule 1: Ornithine carbamoyltransferase

Chain B:





- Molecule 1: Ornithine carbamoyltransferase

Chain C:





- Molecule 1: Ornithine carbamoyltransferase

Chain D:





- Molecule 1: Ornithine carbamoyltransferase

Chain E:





- Molecule 1: Ornithine carbamoyltransferase

Chain F:


3%
94%
GLY
S0
D8
D72
S73
G74
S75
T76
Q77
L78
D81
Q85
K89
V90
L123
D187
P188
S189
S228
M229
GLY
GLN
GLU
ASN
ASP
GLY
L236
L247
R250
S307

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	91.44Å 142.37Å 99.02Å 90.00° 117.17° 90.00°	Depositor
Resolution (Å)	49.55 – 1.67 88.10 – 1.67	Depositor EDS
% Data completeness (in resolution range)	99.2 (49.55-1.67) 99.2 (88.10-1.67)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.21 (at 1.67Å)	Xtriage
Refinement program	PHENIX 1.18.2	Depositor
R, R_{free}	0.161 , 0.180 0.163 , 0.179	Depositor DCC
R_{free} test set	13179 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å^2)	27.9	Xtriage
Anisotropy	0.271	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 41.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.016 for h,-k,-h-l	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	15612	wwPDB-VP
Average B, all atoms (Å^2)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.81% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CP, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.43	0/2372	0.65	0/3229
1	B	0.39	0/2292	0.62	0/3120
1	C	0.41	0/2358	0.63	0/3209
1	D	0.42	0/2308	0.64	0/3142
1	E	0.44	0/2305	0.62	0/3138
1	F	0.42	0/2319	0.61	0/3158
All	All	0.42	0/13954	0.63	0/18996

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2327	0	2295	3	0
1	B	2248	0	2220	8	0
1	C	2317	0	2286	6	0
1	D	2264	0	2233	9	0
1	E	2261	0	2237	9	0
1	F	2275	0	2235	4	0
2	A	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	10	0	0	1	0
2	E	15	0	0	1	0
3	A	8	0	2	0	0
3	B	8	0	2	0	0
3	C	8	0	2	0	0
3	D	8	0	2	0	0
3	E	8	0	2	0	0
3	F	8	0	2	0	0
4	A	8	0	12	0	0
4	C	4	0	6	1	0
4	D	4	0	6	0	0
4	E	4	0	6	0	0
4	F	4	0	6	1	0
5	A	308	0	0	1	0
5	B	260	0	0	1	0
5	C	299	0	0	0	0
5	D	311	0	0	0	0
5	E	311	0	0	1	0
5	F	314	0	0	0	0
All	All	15612	0	13554	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:88:ALA:HB2	1:C:111:MET:HG2	1.78	0.66
1:B:78:LEU:HD11	1:B:87:THR:HG21	1.79	0.63
1:D:212:HIS:NE2	2:D:503:PO4:O2	2.24	0.63
1:D:191:ARG:O	1:D:195:GLU:HG2	2.02	0.60
1:E:1:VAL:HG12	1:E:2:ILE:HD13	1.87	0.57
1:B:220:VAL:HG13	1:B:260:ILE:HG23	1.86	0.56
1:B:222:VAL:HG22	1:B:262:LEU:HB2	1.89	0.54
1:C:72:ASP:O	4:C:503:EDO:H22	2.08	0.54
1:D:154:PHE:HE1	1:D:221:LEU:HD22	1.73	0.54
1:F:72:ASP:O	4:F:402:EDO:H22	2.08	0.53
1:B:5:PHE:CE2	1:B:129:PRO:HB2	2.45	0.52
1:E:4[A]:HIS:CD2	5:E:610:HOH:O	2.65	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:88:ALA:HB3	1:E:114:VAL:HG12	1.94	0.49
1:B:1:VAL:O	1:B:2:ILE:HD13	2.14	0.47
1:D:5:PHE:CE2	1:D:129:PRO:HB2	2.50	0.47
1:F:187:ASP:OD1	1:F:189:SER:OG	2.31	0.47
1:D:154:PHE:CE1	1:D:221:LEU:HD22	2.50	0.46
1:F:85:GLN:O	1:F:89:LYS:HG3	2.15	0.46
1:B:123:LEU:HD12	1:B:128:HIS:CD2	2.50	0.45
1:D:187:ASP:OD2	1:D:189:SER:HB3	2.16	0.45
1:E:5:PHE:CE2	1:E:129:PRO:HB2	2.52	0.45
1:D:77:GLN:HB3	1:D:80:ARG:HE	1.81	0.45
1:D:22:LEU:HB3	1:D:298:ALA:HB1	2.00	0.44
1:A:5:PHE:CE2	1:A:129:PRO:HB2	2.53	0.44
1:C:5:PHE:CE2	1:C:129:PRO:HB2	2.52	0.44
1:E:286:TRP:HZ3	1:F:90:VAL:HG23	1.82	0.43
1:C:85:GLN:O	1:C:89:LYS:HG3	2.18	0.43
1:C:226:TRP:CE2	1:C:242:PHE:HB3	2.54	0.43
1:A:4[B]:HIS:CD2	5:A:557:HOH:O	2.72	0.42
1:D:179:ALA:HA	1:D:208:THR:O	2.19	0.42
1:E:195:GLU:O	1:E:199:GLN:HG3	2.19	0.42
1:E:222:VAL:HG22	1:E:262:LEU:HB2	2.02	0.41
1:E:212:HIS:HE2	2:E:503:PO4:P	2.43	0.41
1:B:89:LYS:HD3	5:B:751:HOH:O	2.20	0.41
1:C:91:LEU:O	1:C:95:VAL:HG22	2.20	0.41
1:A:53:THR:HG21	1:A:122:ALA:HB1	2.02	0.41
1:B:78:LEU:HD23	1:B:100:TRP:CZ2	2.55	0.41
1:E:45:ILE:HD13	1:E:78:LEU:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/309 (99%)	300 (98%)	6 (2%)	1 (0%)	41	23
1	B	295/309 (96%)	288 (98%)	6 (2%)	1 (0%)	41	23
1	C	305/309 (99%)	299 (98%)	5 (2%)	1 (0%)	41	23
1	D	297/309 (96%)	287 (97%)	9 (3%)	1 (0%)	41	23
1	E	296/309 (96%)	287 (97%)	8 (3%)	1 (0%)	41	23
1	F	299/309 (97%)	292 (98%)	6 (2%)	1 (0%)	41	23
All	All	1799/1854 (97%)	1753 (97%)	40 (2%)	6 (0%)	41	23

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	LEU
1	B	123	LEU
1	C	123	LEU
1	D	123	LEU
1	E	123	LEU
1	F	123	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/243 (99%)	238 (99%)	3 (1%)	71	57
1	B	231/243 (95%)	230 (100%)	1 (0%)	91	86
1	C	240/243 (99%)	240 (100%)	0	100	100
1	D	233/243 (96%)	231 (99%)	2 (1%)	78	69
1	E	234/243 (96%)	231 (99%)	3 (1%)	69	54
1	F	234/243 (96%)	231 (99%)	3 (1%)	69	54
All	All	1413/1458 (97%)	1401 (99%)	12 (1%)	78	72

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	VAL
1	A	128	HIS
1	A	306	ARG
1	B	8	ASP
1	D	8	ASP
1	D	247	LEU
1	E	8	ASP
1	E	114	VAL
1	E	247	LEU
1	F	8	ASP
1	F	247	LEU
1	F	250	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	199	GLN
1	C	212	HIS
1	E	49	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	EDO	E	504	-	3,3,3	0.50	0	2,2,2	0.31	0
3	CP	E	502	-	6,7,7	7.38	3 (50%)	7,10,10	2.45	2 (28%)
4	EDO	D	504	-	3,3,3	0.51	0	2,2,2	0.30	0
4	EDO	C	503	-	3,3,3	0.50	0	2,2,2	0.17	0
3	CP	D	502	-	6,7,7	6.24	3 (50%)	7,10,10	2.25	2 (28%)
2	PO4	C	501	-	4,4,4	0.84	0	6,6,6	0.57	0
4	EDO	A	404	-	3,3,3	0.40	0	2,2,2	0.31	0
2	PO4	E	505	-	4,4,4	0.87	0	6,6,6	0.57	0
2	PO4	E	501	-	4,4,4	1.00	0	6,6,6	0.84	0
3	CP	C	502	-	6,7,7	6.15	3 (50%)	7,10,10	2.32	2 (28%)
3	CP	F	401	-	6,7,7	6.93	3 (50%)	7,10,10	2.07	1 (14%)
2	PO4	D	503	-	4,4,4	0.78	0	6,6,6	0.59	0
2	PO4	A	401	-	4,4,4	0.88	0	6,6,6	0.52	0
4	EDO	F	402	-	3,3,3	0.39	0	2,2,2	0.47	0
2	PO4	E	503	-	4,4,4	0.84	0	6,6,6	0.49	0
4	EDO	A	405	-	3,3,3	0.52	0	2,2,2	0.24	0
2	PO4	B	401	-	4,4,4	0.84	0	6,6,6	0.53	0
2	PO4	A	403	-	4,4,4	0.74	0	6,6,6	0.45	0
3	CP	A	402	-	6,7,7	7.39	3 (50%)	7,10,10	2.20	2 (28%)
3	CP	B	402	-	6,7,7	7.69	3 (50%)	7,10,10	2.32	2 (28%)
2	PO4	D	501	-	4,4,4	1.02	0	6,6,6	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	E	504	-	-	0/1/1/1	-
3	CP	F	401	-	-	0/3/5/5	-
3	CP	C	502	-	-	0/3/5/5	-
4	EDO	F	402	-	-	0/1/1/1	-
3	CP	A	402	-	-	0/3/5/5	-
3	CP	E	502	-	-	0/3/5/5	-
4	EDO	A	404	-	-	0/1/1/1	-
4	EDO	D	504	-	-	0/1/1/1	-
4	EDO	A	405	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	503	-	-	0/1/1/1	-
3	CP	D	502	-	-	0/3/5/5	-
3	CP	B	402	-	-	0/3/5/5	-

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	CP	P-O4P	17.98	1.86	1.59
3	E	502	CP	P-O4P	17.23	1.85	1.59
3	A	402	CP	P-O4P	17.21	1.85	1.59
3	F	401	CP	P-O4P	15.48	1.82	1.59
3	C	502	CP	P-O4P	14.52	1.81	1.59
3	D	502	CP	P-O4P	14.04	1.80	1.59
3	F	401	CP	C-N	5.45	1.43	1.33
3	D	502	CP	C-N	4.71	1.42	1.33
3	A	402	CP	C-N	4.54	1.42	1.33
3	B	402	CP	O-C	4.45	1.27	1.21
3	E	502	CP	C-N	4.27	1.41	1.33
3	F	401	CP	O-C	4.07	1.27	1.21
3	D	502	CP	O-C	3.70	1.26	1.21
3	B	402	CP	C-N	3.23	1.39	1.33
3	E	502	CP	O-C	3.07	1.26	1.21
3	A	402	CP	O-C	3.03	1.26	1.21
3	C	502	CP	C-N	2.72	1.38	1.33
3	C	502	CP	O-C	2.64	1.25	1.21

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	E	502	CP	O-C-N	-5.27	116.81	125.51
3	B	402	CP	O-C-N	-5.02	117.23	125.51
3	C	502	CP	O-C-N	-4.80	117.59	125.51
3	F	401	CP	O-C-N	-4.69	117.78	125.51
3	A	402	CP	O-C-N	-4.53	118.04	125.51
3	D	502	CP	O-C-N	-4.49	118.11	125.51
3	B	402	CP	O3P-P-O2P	2.69	117.91	107.64
3	A	402	CP	O4P-P-O1P	-2.67	99.84	109.32
3	D	502	CP	O3P-P-O2P	2.60	117.59	107.64
3	C	502	CP	O3P-P-O2P	2.58	117.50	107.64
3	E	502	CP	O3P-P-O2P	2.02	115.35	107.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	503	EDO	1	0
2	D	503	PO4	1	0
4	F	402	EDO	1	0
2	E	503	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand CP E 502



Ligand CP D 502



Ligand CP C 502



Ligand CP F 401



Ligand CP A 402



Ligand CP B 402



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	308/309 (99%)	-0.43	5 (1%)	72	75	22, 30, 46, 71	0
1	B	298/309 (96%)	-0.24	10 (3%)	45	48	22, 36, 59, 81	0
1	C	307/309 (99%)	-0.50	2 (0%)	87	89	24, 34, 50, 68	0
1	D	300/309 (97%)	-0.49	5 (1%)	70	74	22, 31, 48, 90	0
1	E	299/309 (96%)	-0.46	3 (1%)	82	85	23, 31, 50, 63	0
1	F	302/309 (97%)	-0.36	9 (2%)	50	53	22, 31, 55, 79	0
All	All	1814/1854 (97%)	-0.42	34 (1%)	66	70	22, 32, 53, 90	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	74	GLY	7.1
1	B	228	SER	6.3
1	B	75	SER	5.8
1	D	237	ASP	5.7
1	F	228	SER	5.4
1	F	73	SER	5.4
1	A	74	GLY	5.2
1	B	76	THR	5.2
1	F	74	GLY	4.7
1	B	227	THR	4.6
1	A	75	SER	4.1
1	B	1	VAL	4.0
1	E	236	LEU	3.8
1	F	229	MET	3.8
1	A	73	SER	3.7
1	A	76	THR	3.5
1	A	0	SER	3.5
1	D	74	GLY	3.4
1	E	73	SER	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	73	SER	3.0
1	F	0	SER	3.0
1	F	75	SER	2.8
1	B	73	SER	2.7
1	B	239	VAL	2.7
1	D	0	SER	2.6
1	C	74	GLY	2.3
1	F	76	THR	2.2
1	B	268	HIS	2.2
1	F	81	ASP	2.2
1	D	236	LEU	2.2
1	B	77	GLN	2.1
1	D	75	SER	2.1
1	E	1	VAL	2.0
1	F	78	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	EDO	C	503	4/4	0.80	0.12	43,53,56,57	0
2	PO4	D	503	5/5	0.86	0.33	60,65,68,68	0
4	EDO	F	402	4/4	0.87	0.17	50,53,54,57	0
4	EDO	A	404	4/4	0.88	0.15	49,51,53,57	0
2	PO4	A	403	5/5	0.90	0.20	54,62,65,73	0
4	EDO	E	504	4/4	0.91	0.08	42,47,49,54	0
4	EDO	A	405	4/4	0.91	0.21	43,43,46,52	0
4	EDO	D	504	4/4	0.93	0.13	42,43,44,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	CP	B	402	8/8	0.94	0.10	31,37,38,39	0
3	CP	A	402	8/8	0.95	0.09	27,30,34,35	0
2	PO4	B	401	5/5	0.96	0.17	52,60,65,65	0
3	CP	D	502	8/8	0.96	0.08	29,30,32,33	0
3	CP	E	502	8/8	0.96	0.10	30,32,33,36	0
2	PO4	E	503	5/5	0.97	0.21	44,51,55,59	0
2	PO4	D	501	5/5	0.98	0.06	40,40,43,44	0
2	PO4	E	505	5/5	0.98	0.14	50,50,56,62	0
3	CP	F	401	8/8	0.98	0.07	26,29,31,35	0
2	PO4	E	501	5/5	0.99	0.07	30,36,40,42	0
2	PO4	A	401	5/5	0.99	0.06	36,40,42,45	0
3	CP	C	502	8/8	0.99	0.06	26,27,29,31	0
2	PO4	C	501	5/5	0.99	0.06	42,45,47,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CP B 402:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around CP A 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CP D 502:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CP E 502:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CP F 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CP C 502:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.