



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 13, 2023 – 02:33 AM EDT

PDB ID : 4NOG
Title : Crystal structure of a putative ornithine aminotransferase from *Toxoplasma gondii* ME49 in complex with pyrodoxal-5'-phosphate
Authors : Filippova, E.V.; Halavaty, A.; Ruan, J.; Shuvalova, L.; Flores, K.; Dubrovskaya, I.; Ngo, H.; Shanmugam, D.; Roos, D.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2013-11-19
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	442	
1	B	442	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

Validation Pipeline (wwPDB-VP) : 2.35.1

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BME	A	502	-	-	X	-
6	PEG	B	504	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative ornithine aminotransferase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	422	3401	2147	603	630	21	0	16	0
1	B	424	3384	2133	602	629	20	0	12	0

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	16	MET	-	expression tag	UNP S8EY38
A	442	GLY	-	expression tag	UNP S8EY38
A	443	GLU	-	expression tag	UNP S8EY38
A	444	ASN	-	expression tag	UNP S8EY38
A	445	LEU	-	expression tag	UNP S8EY38
A	446	TYR	-	expression tag	UNP S8EY38
A	447	PHE	-	expression tag	UNP S8EY38
A	448	GLN	-	expression tag	UNP S8EY38
A	449	SER	-	expression tag	UNP S8EY38
A	450	ALA	-	expression tag	UNP S8EY38
A	451	GLY	-	expression tag	UNP S8EY38
A	452	HIS	-	expression tag	UNP S8EY38
A	453	HIS	-	expression tag	UNP S8EY38
A	454	HIS	-	expression tag	UNP S8EY38
A	455	HIS	-	expression tag	UNP S8EY38
A	456	HIS	-	expression tag	UNP S8EY38
A	457	HIS	-	expression tag	UNP S8EY38
B	16	MET	-	expression tag	UNP S8EY38
B	442	GLY	-	expression tag	UNP S8EY38
B	443	GLU	-	expression tag	UNP S8EY38
B	444	ASN	-	expression tag	UNP S8EY38
B	445	LEU	-	expression tag	UNP S8EY38
B	446	TYR	-	expression tag	UNP S8EY38
B	447	PHE	-	expression tag	UNP S8EY38
B	448	GLN	-	expression tag	UNP S8EY38

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	449	SER	-	expression tag	UNP S8EY38
B	450	ALA	-	expression tag	UNP S8EY38
B	451	GLY	-	expression tag	UNP S8EY38
B	452	HIS	-	expression tag	UNP S8EY38
B	453	HIS	-	expression tag	UNP S8EY38
B	454	HIS	-	expression tag	UNP S8EY38
B	455	HIS	-	expression tag	UNP S8EY38
B	456	HIS	-	expression tag	UNP S8EY38
B	457	HIS	-	expression tag	UNP S8EY38

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	B	1	Total	C	N	O	P	0	0
			16	8	1	6	1		

- Molecule 3 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
3	A	1	4	2	1	1	0	0
3	B	1	4	2	1	1	0	0


- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2^-$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	B	1	4	2	2	0	0


- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN

E-1,3-DIOL (three-letter code: BTB) (formula: $C_8H_{19}NO_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
5	B	1	14	8	1	5	0	0

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
			Total	C			O
6	B	1	7	4	3	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	549	Total 552	O 552	0	3
7	B	492	Total 495	O 495	0	3

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative ornithine aminotransferase, mitochondrial

Chain A:  88% 7% 5%



- Molecule 1: Putative ornithine aminotransferase, mitochondrial

Chain B:  86% 9% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	56.19Å 61.33Å 63.68Å 100.58° 93.23° 107.74°	Depositor
Resolution (Å)	62.14 – 1.20 25.17 – 1.20	Depositor EDS
% Data completeness (in resolution range)	90.8 (62.14-1.20) 90.9 (25.17-1.20)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.20Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.133 , 0.165 0.133 , 0.166	Depositor DCC
R_{free} test set	11260 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	9.8	Xtrriage
Anisotropy	0.580	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7897	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BME, ACT, PEG, PLP, BTB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.89	3/3464 (0.1%)	0.96	7/4688 (0.1%)
1	B	0.94	3/3447 (0.1%)	0.97	9/4665 (0.2%)
All	All	0.92	6/6911 (0.1%)	0.97	16/9353 (0.2%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	274	HIS	C-O	5.90	1.34	1.23
1	B	210	GLU	CD-OE2	-5.67	1.19	1.25
1	B	300	GLY	C-O	-5.39	1.15	1.23
1	A	276	GLU	CD-OE1	5.13	1.31	1.25
1	A	41	ASP	CB-CG	-5.10	1.41	1.51
1	B	199	GLU	CD-OE2	5.01	1.31	1.25

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	63	ARG	NE-CZ-NH2	-12.07	114.26	120.30
1	A	166	LEU	CA-CB-CG	6.56	130.39	115.30
1	A	97	ARG	NE-CZ-NH1	-5.72	117.44	120.30
1	A	422	ARG	NE-CZ-NH2	-5.60	117.50	120.30
1	B	66	ASP	CB-CG-OD2	-5.60	113.26	118.30
1	A	42	ARG	NE-CZ-NH2	-5.48	117.56	120.30
1	B	222	PHE	CB-CG-CD1	5.47	124.63	120.80
1	B	355	ARG	NE-CZ-NH2	-5.41	117.60	120.30
1	B	427	ARG	NE-CZ-NH2	-5.39	117.60	120.30
1	A	342[A]	ARG	CG-CD-NE	-5.34	100.59	111.80
1	A	342[B]	ARG	CG-CD-NE	-5.34	100.59	111.80
1	B	38	PHE	CB-CG-CD2	-5.33	117.07	120.80
1	A	129	ARG	NE-CZ-NH2	-5.28	117.66	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	63	ARG	CG-CD-NE	-5.23	100.81	111.80
1	B	26	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	B	101	ARG	NE-CZ-NH1	5.11	122.85	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3401	0	3418	35	0
1	B	3384	0	3396	36	0
2	A	16	0	8	0	0
2	B	16	0	7	1	0
3	A	4	0	5	8	0
3	B	4	0	5	2	0
4	B	4	0	3	0	0
5	B	14	0	19	0	0
6	B	7	0	10	5	0
7	A	552	0	0	9	0
7	B	495	0	0	17	0
All	All	7897	0	6871	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1148:HOH:O	1:B:92[B]:ILE:HD11	0.95	1.13
1:A:92[A]:ILE:HD11	7:B:1084:HOH:O	0.83	1.00
1:B:425:THR:HG21	3:B:505:BME:H22	1.45	0.97
1:A:353[B]:CYS:HG	3:A:502:BME:C1	1.85	0.89
1:A:353[B]:CYS:SG	1:A:429[B]:ILE:HD11	2.14	0.87
1:B:352:ASP:HB3	7:B:993:HOH:O	1.75	0.87

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:25:TYR:HB2	7:B:945:HOH:O	1.75	0.86
1:A:274:HIS:CD2	7:A:1016:HOH:O	2.29	0.85
1:B:35:GLU:HG3	7:B:912:HOH:O	1.78	0.83
1:A:353[B]:CYS:SG	1:A:429[B]:ILE:CD1	2.68	0.81
1:B:293[B]:VAL:CG1	7:B:825:HOH:O	2.27	0.81
1:B:352:ASP:CB	7:B:993:HOH:O	2.26	0.81
1:B:436:GLU:OE2	1:B:440:LYS:HE3	1.80	0.81
1:A:79[A]:VAL:HG23	1:A:286[A]:LYS:HZ3	1.47	0.80
1:A:79[A]:VAL:HG23	1:A:286[A]:LYS:NZ	1.99	0.78
1:A:353[B]:CYS:SG	3:A:502:BME:C1	2.74	0.76
1:A:353[B]:CYS:HG	3:A:502:BME:HO1	1.29	0.75
1:A:353[B]:CYS:SG	3:A:502:BME:S2	2.86	0.73
1:B:366:LYS:NZ	7:B:1076:HOH:O	2.23	0.72
1:B:268:ARG:NH1	7:B:1082:HOH:O	2.23	0.71
1:A:353[B]:CYS:SG	3:A:502:BME:C2	2.80	0.69
1:B:366:LYS:HE2	7:B:1049:HOH:O	1.93	0.68
1:A:353[B]:CYS:HG	3:A:502:BME:C2	2.09	0.65
1:B:425:THR:CG2	3:B:505:BME:H22	2.25	0.64
1:A:353[B]:CYS:HB2	3:A:502:BME:S2	2.38	0.63
1:B:51:PRO:HD3	7:B:850:HOH:O	1.98	0.62
1:B:352:ASP:CG	7:B:993:HOH:O	2.38	0.61
1:A:101[A]:ARG:NE	7:A:1095:HOH:O	2.33	0.61
1:B:35:GLU:CG	7:B:912:HOH:O	2.42	0.60
1:A:206[A]:VAL:HG12	7:A:765:HOH:O	2.01	0.59
1:A:353[B]:CYS:CB	3:A:502:BME:S2	2.92	0.58
1:A:281[A]:ILE:HD11	1:A:304:VAL:HG11	1.86	0.58
1:B:269:LEU:H	6:B:504:PEG:H42	1.68	0.57
1:B:356[A]:ARG:O	1:B:356[A]:ARG:HG2	2.06	0.54
1:A:22:ILE:HD13	1:B:328:VAL:HG21	1.90	0.52
1:B:33:THR:HA	7:B:769[B]:HOH:O	2.10	0.51
1:B:365[A]:ILE:HD12	1:B:379:VAL:HG12	1.91	0.51
1:A:410:PHE:CE1	1:A:428[A]:ILE:HD13	2.46	0.51
1:B:90:ARG:HH22	1:B:332:THR:HG22	1.77	0.50
1:B:293[A]:VAL:CG2	1:B:294:PRO:HD2	2.42	0.50
1:B:269:LEU:H	6:B:504:PEG:C4	2.25	0.49
1:A:280:ASP:O	1:A:281[A]:ILE:HD13	2.13	0.48
1:B:273:ASP:OD2	6:B:504:PEG:H42	2.13	0.48
1:A:79[A]:VAL:CG2	1:A:286[A]:LYS:NZ	2.75	0.48
7:A:1148:HOH:O	1:B:92[B]:ILE:CD1	1.82	0.48
1:A:92[A]:ILE:CD1	7:B:1084:HOH:O	1.71	0.48
1:A:274:HIS:HD2	7:A:1016:HOH:O	1.83	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:183:THR:HG22	1:B:200[B]:LEU:HD23	1.96	0.47
1:A:32:LYS:HE2	1:A:32:LYS:HB2	1.61	0.47
1:A:76:LEU:HD11	1:A:79[A]:VAL:HG12	1.95	0.47
1:A:342[B]:ARG:NH2	7:A:1024:HOH:O	2.26	0.47
1:A:101[B]:ARG:HH21	1:A:101[B]:ARG:HG3	1.80	0.46
6:B:504:PEG:H11	7:B:1050:HOH:O	2.16	0.46
1:B:183:THR:CG2	1:B:200[A]:LEU:HD13	2.46	0.46
1:B:293[A]:VAL:HG22	1:B:294:PRO:HD2	1.99	0.44
1:A:76:LEU:HD11	1:A:79[A]:VAL:CG1	2.48	0.44
1:A:87:CYS:HA	7:B:1084:HOH:O	2.18	0.44
1:B:347:GLY:HA3	1:B:372:GLY:O	2.19	0.43
1:B:183:THR:CG2	1:B:200[B]:LEU:HD23	2.48	0.43
1:B:301:ARG:NE	1:B:304:VAL:HG23	2.33	0.43
1:B:301:ARG:HE	1:B:304:VAL:HG23	1.83	0.43
1:A:396:ASN:O	1:A:427:ARG:HD3	2.19	0.42
1:A:245:GLU:CD	7:A:1084:HOH:O	2.57	0.42
1:B:172:TRP:CZ2	1:B:178:ALA:HA	2.55	0.42
1:B:25:TYR:CB	7:B:945:HOH:O	2.49	0.41
1:A:335:LYS:HE2	7:A:1114:HOH:O	2.20	0.41
1:A:172:TRP:CZ2	1:A:178:ALA:HA	2.55	0.41
1:A:410:PHE:CE1	1:A:428[A]:ILE:CD1	3.03	0.41
1:B:269:LEU:N	6:B:504:PEG:H42	2.35	0.41
1:B:286[B]:LYS:NZ	2:B:501:PLP:O4A	2.46	0.41
1:A:347:GLY:HA3	1:A:372:GLY:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	436/442 (99%)	422 (97%)	14 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	434/442 (98%)	419 (96%)	15 (4%)	0	100	100
All	All	870/884 (98%)	841 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/365 (100%)	360 (99%)	5 (1%)	67	32
1	B	362/365 (99%)	358 (99%)	4 (1%)	73	41
All	All	727/730 (100%)	718 (99%)	9 (1%)	76	37

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	LYS
1	A	294	PRO
1	A	428[A]	ILE
1	A	428[B]	ILE
1	A	437	GLU
1	B	26	ARG
1	B	275	ASP
1	B	365[A]	ILE
1	B	365[B]	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	274	HIS
1	B	47	GLN
1	B	170	ASN
1	B	396	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLP	A	501	-	16,16,16	1.24	3 (18%)	20,23,23	1.18	1 (5%)
4	ACT	B	502	-	3,3,3	0.73	0	3,3,3	0.71	0
2	PLP	B	501	-	16,16,16	1.34	2 (12%)	20,23,23	1.61	4 (20%)
5	BTB	B	503	-	13,13,13	1.07	2 (15%)	7,16,16	1.50	1 (14%)
3	BME	A	502	-	3,3,3	0.51	0	1,2,2	0.75	0
6	PEG	B	504	-	6,6,6	0.56	0	5,5,5	1.09	0
3	BME	B	505	1	3,3,3	0.18	0	1,2,2	1.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	501	-	-	0/8/8/8	0/1/1/1
2	PLP	B	501	-	-	0/8/8/8	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BTB	B	503	-	-	6/21/21/21	-
3	BME	A	502	-	-	1/1/1/1	-
6	PEG	B	504	-	-	3/4/4/4	-
3	BME	B	505	1	-	0/1/1/1	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	PLP	C3-C2	2.81	1.43	1.40
2	B	501	PLP	C4-C5	2.62	1.45	1.42
2	B	501	PLP	O4A-C4A	2.61	1.30	1.21
2	A	501	PLP	C4-C3	2.55	1.44	1.40
5	B	503	BTB	C5-N	2.31	1.51	1.48
2	A	501	PLP	C4-C5	2.21	1.44	1.42
5	B	503	BTB	O3-C3	2.02	1.49	1.42

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	PLP	C4-C3-C2	-3.74	117.88	120.19
2	A	501	PLP	C3-C4-C5	-3.18	115.82	118.26
2	B	501	PLP	C5-C6-N1	-2.64	119.42	123.82
2	B	501	PLP	O2P-P-O4P	-2.61	99.78	106.73
2	B	501	PLP	O4A-C4A-C4	-2.42	119.64	124.91
5	B	503	BTB	C6-C5-N	2.19	120.13	111.59

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	503	BTB	C1-C2-N-C5
5	B	503	BTB	C3-C2-N-C5
5	B	503	BTB	C4-C2-N-C5
3	A	502	BME	O1-C1-C2-S2
6	B	504	PEG	O1-C1-C2-O2
6	B	504	PEG	C1-C2-O2-C3
5	B	503	BTB	C8-C7-N-C5
6	B	504	PEG	C4-C3-O2-C2
5	B	503	BTB	N-C2-C4-O4
5	B	503	BTB	C6-C5-N-C2

There are no ring outliers.

4 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	PLP	1	0
3	A	502	BME	8	0
6	B	504	PEG	5	0
3	B	505	BME	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	422/442 (95%)	-0.54	2 (0%) 91 91	8, 13, 33, 92	0
1	B	424/442 (95%)	-0.42	5 (1%) 79 79	8, 13, 36, 98	0
All	All	846/884 (95%)	-0.48	7 (0%) 86 86	8, 13, 35, 98	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	18	ARG	4.1
1	B	31	LEU	2.9
1	B	33	THR	2.8
1	B	20	THR	2.5
1	B	17	ALA	2.3
1	A	32	LYS	2.1
1	B	24	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	BME	B	505	4/4	0.88	0.17	34,34,34,36	3
6	PEG	B	504	7/7	0.88	0.18	27,32,43,44	0
5	BTB	B	503	14/14	0.91	0.15	19,22,31,37	0
3	BME	A	502	4/4	0.93	0.13	27,35,39,42	1
4	ACT	B	502	4/4	0.95	0.14	28,32,33,38	0
2	PLP	B	501	16/16	0.99	0.04	10,11,14,53	0
2	PLP	A	501	16/16	0.99	0.04	9,11,14,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.