

# Full wwPDB X-ray Structure Validation Report (i)

Dec 16, 2023 – 11:48 PM EST

PDB ID : 2NPI

Title : Clp1-ATP-Pcf11 complex

Authors: Noble, C.G.; Beuth, B.; Taylor, I.A.

Deposited on : 2006-10-27

Resolution : 2.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to X-ray structures of similar resolution

Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length		Quality of chain					
1	A	460	5%		74%		17%	•	7%
1	В	460	3%		73%		18%	•	7%
2	С	110	17%	•		80%			
2	D	110	18%	5%		77%			_



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7437 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protein CLP1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	A	428	Total 3412	C 2193	N 571	_	S 6	Se 6	0	0	0
1	В	428	Total 3412	C 2193	N 571	_	S 6	Se 6	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	MSE	-	modified residue	UNP Q08685
A	-13	GLY	-	cloning artifact	UNP Q08685
A	-12	SER	-	cloning artifact	UNP Q08685
A	-11	SER	-	cloning artifact	UNP Q08685
A	-10	HIS	-	expression tag	UNP Q08685
A	-9	HIS	-	expression tag	UNP Q08685
A	-8	HIS	-	expression tag	UNP Q08685
A	-7	HIS	-	expression tag	UNP Q08685
A	-6	HIS	-	expression tag	UNP Q08685
A	-5	HIS	-	expression tag	UNP Q08685
A	-4	SER	-	cloning artifact	UNP Q08685
A	-3	GLN	-	cloning artifact	UNP Q08685
A	-2	ASP	-	cloning artifact	UNP Q08685
A	-1	PRO	_	cloning artifact	UNP Q08685
A	0	ASN	-	cloning artifact	UNP Q08685
A	1	SER	_	cloning artifact	UNP Q08685
A	103	MSE	MET	modified residue	UNP Q08685
A	112	MSE	MET	modified residue	UNP Q08685
A	118	MSE	MET	modified residue	UNP Q08685
A	206	MSE	MET	modified residue	UNP Q08685
A	277	MSE	MET	modified residue	UNP Q08685
A	436	MSE	MET	modified residue	UNP Q08685
В	-14	MSE	-	modified residue	UNP Q08685
В	-13	GLY	-	cloning artifact	UNP Q08685
В	-12	SER	-	cloning artifact	UNP Q08685
				Continued	on nert nage



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	-11	SER	-	cloning artifact	UNP Q08685
В	-10	HIS	-	expression tag	UNP Q08685
В	-9	HIS	-	expression tag	UNP Q08685
В	-8	HIS	-	expression tag	UNP Q08685
В	-7	HIS	-	expression tag	UNP Q08685
В	-6	HIS	-	expression tag	UNP Q08685
В	-5	HIS	-	expression tag	UNP Q08685
В	-4	SER	-	cloning artifact	UNP Q08685
В	-3	GLN	-	cloning artifact	UNP Q08685
В	-2	ASP	-	cloning artifact	UNP Q08685
В	-1	PRO	-	cloning artifact	UNP Q08685
В	0	ASN	-	cloning artifact	UNP Q08685
В	1	SER	-	cloning artifact	UNP Q08685
В	103	MSE	MET	modified residue	UNP Q08685
В	112	MSE	MET	modified residue	UNP Q08685
В	118	MSE	MET	modified residue	UNP Q08685
В	206	MSE	MET	modified residue	UNP Q08685
В	277	MSE	MET	modified residue	UNP Q08685
В	436	MSE	MET	modified residue	UNP Q08685

• Molecule 2 is a protein called Protein PCF11.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
2	С	22	Total 189	C 120			0	0	0
2	D	25	Total 212		N 36	O 43	0	0	0

There are 2 discrepancies between the modelled and reference sequences:  $\[$ 


Chain	Residue	Modelled	Actual	Comment	Reference
С	560	MSE	MET	modified residue	UNP P39081
D	560	MSE	MET	modified residue	UNP P39081

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0



 $\bullet$  Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3).$ 



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
1	Λ	1	Total	С	N	О	Р	0	0
4	4   A	1	31	10	5	13	3	0	0
1	D	1	Total	С	N	О	Р	0	0
4	Б	1	31	10	5	13	3	U	U


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	57	Total O 57 57	0	0
5	В	77	Total O 77 77	0	0
5	С	7	Total O 7 7	0	0
5	D	7	Total O 7 7	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	88.97Å 94.99Å 181.40Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 - 2.95	Depositor
rtesolution (A)	14.95 - 2.95	EDS
% Data completeness	90.0 (15.00-2.95)	Depositor
(in resolution range)	90.0 (14.95-2.95)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$< I/\sigma(I) > 1$	1.72 (at 2.96Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
P. P.	0.248 , 0.303	Depositor
$R, R_{free}$	0.257 , $0.301$	DCC
$R_{free}$ test set	1493 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.1	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.26, 53.7	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.52, < L^2>=0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	7437	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	$RMSZ \mid \# Z  > 5$		# Z  > 5	
1	A	0.33	0/3483	0.54	$1/4729 \ (0.0\%)$	
1	В	0.33	0/3483	0.56	0/4729	
2	С	0.34	0/194	0.43	0/263	
2	D	0.35	0/217	0.43	0/293	
All	All	0.33	0/7377	0.55	1/10014 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
1	A	199	LEU	CA-CB-CG	5.24	127.35	115.30

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3412	0	3451	44	0
1	В	3412	0	3451	47	0
2	С	189	0	168	2	0
2	D	212	0	192	6	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	31	0	12	0	0
4	В	31	0	12	0	0
5	A	57	0	0	0	0
5	В	77	0	0	7	0
5	С	7	0	0	0	0
5	D	7	0	0	0	0
All	All	7437	0	7286	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 9	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:148:LEU:HD11	1:A:180:ASP:HA	1.46	0.95
1:B:109:LEU:HD21	1:B:278:LEU:HD11	1.55	0.89
1:B:148:LEU:HD11	1:B:180:ASP:HA	1.56	0.85
1:B:125:ARG:H	1:B:275:ASN:HD21	1.26	0.84
1:B:376:ILE:HG23	1:B:381:LEU:HD12	1.63	0.78
1:A:172:CYS:HB3	1:A:209:ASN:HA	1.69	0.73
1:A:168:THR:HG22	1:A:169:VAL:H	1.53	0.73
1:B:127:VAL:HG22	1:B:274:VAL:HG11	1.73	0.69
1:B:180:ASP:HB2	5:B:2508:HOH:O	1.93	0.67
1:B:406:LEU:HD22	2:D:482:TRP:HB3	1.82	0.60
1:A:278:LEU:HD22	1:A:307:ILE:HD11	1.84	0.59
1:A:360:ASN:HB2	1:A:363:ASP:HB2	1.84	0.59
1:A:168:THR:HG23	2:C:483:TYR:OH	2.01	0.59
1:A:53:VAL:HG13	1:A:81:THR:HG21	1.84	0.58
1:A:376:ILE:HG23	1:A:381:LEU:HD12	1.86	0.57
1:B:360:ASN:HB2	1:B:363:ASP:HB2	1.87	0.56
1:B:214:ARG:HD3	5:B:2527:HOH:O	2.05	0.56
1:A:127:VAL:HG22	1:A:274:VAL:HG11	1.87	0.56
1:A:279:VAL:HG21	1:A:291:VAL:HG11	1.87	0.55
1:A:34:TRP:O	1:A:96:THR:O	2.25	0.55
1:A:278:LEU:HD23	1:A:305:PHE:HB2	1.87	0.55
1:A:279:VAL:HG12	1:A:281:CYS:HB2	1.87	0.55
1:A:353:LEU:HD23	1:A:355:ILE:HD11	1.87	0.55
1:B:168:THR:HG23	2:D:483:TYR:OH	2.06	0.55
1:A:110:HIS:HE1	1:A:147:ALA:HA	1.73	0.54
1:B:427:VAL:HG22	2:D:480:ARG:O	2.07	0.53
2:C:484:LEU:HD22	2:C:488:GLN:HE21	1.74	0.53



Continued from previous page...

Continued from previous		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	$overlap(\AA)$
1:B:382:GLN:HE21	1:B:413:GLU:HA	1.73	0.52
1:A:183:ASP:OD1	1:A:185:GLN:HG2	2.10	0.52
1:B:125:ARG:N	1:B:275:ASN:HD21	2.02	0.52
1:B:263:ALA:HA	1:B:266:HIS:HD2	1.75	0.51
1:B:129:VAL:HA	5:B:2509:HOH:O	2.11	0.50
1:B:208:LYS:HD2	1:B:228:GLN:HE21	1.76	0.50
1:B:360:ASN:HB3	1:B:363:ASP:H	1.75	0.50
1:B:46:VAL:HG13	1:B:83:LEU:HD13	1.94	0.50
1:A:336:SER:H	1:A:339:THR:HB	1.78	0.49
1:A:56:PHE:CZ	1:A:74:PRO:HD2	2.49	0.48
1:A:125:ARG:HG2	1:A:248:CYS:HB3	1.95	0.48
1:A:46:VAL:HG13	1:A:83:LEU:HD13	1.95	0.48
1:A:224:GLU:HG3	1:A:369:VAL:HG11	1.96	0.48
1:A:20:ASP:OD2	1:A:22:GLU:HG2	2.14	0.48
1:A:410:LEU:CD1	1:A:443:TYR:HD2	2.26	0.47
1:B:208:LYS:HD2	1:B:228:GLN:HG3	1.95	0.47
1:B:148:LEU:HD11	1:B:180:ASP:CA	2.36	0.47
1:B:166:ILE:HG22	1:B:167:PHE:CD1	2.49	0.47
1:B:56:PHE:CE2	1:B:138:SER:HB3	2.50	0.46
1:A:208:LYS:HD2	1:A:228:GLN:HG3	1.98	0.46
1:B:439:THR:HG21	5:B:2517:HOH:O	2.14	0.46
1:B:406:LEU:CD2	2:D:482:TRP:HB3	2.46	0.46
1:B:204:GLN:HG3	2:D:492:PHE:CE2	2.51	0.46
1:A:348:VAL:O	1:A:421:LEU:N	2.48	0.45
1:B:102:THR:HG21	1:B:310:LEU:HA	1.99	0.45
1:A:110:HIS:CE1	1:A:147:ALA:HA	2.52	0.45
1:B:206:MSE:HB2	2:D:489:TRP:CE2	2.52	0.45
1:B:381:LEU:HD22	1:B:386:ILE:HD11	1.98	0.45
1:B:102:THR:HB	5:B:2544:HOH:O	2.18	0.44
1:B:252:THR:HG22	1:B:253:PRO:HD2	1.99	0.44
1:B:262:LEU:HB3	1:B:265:LEU:HD12	1.99	0.44
1:B:109:LEU:HD22	1:B:307:ILE:HD11	2.00	0.44
1:A:146:TYR:CB	1:A:149:LYS:HB3	2.48	0.44
1:B:87:CYS:HA	1:B:88:PRO:HD3	1.89	0.44
1:B:126:VAL:HG13	1:B:276:ILE:HG12	2.01	0.43
1:B:168:THR:HG22	1:B:169:VAL:HG22	2.00	0.43
1:A:141:ARG:O	1:A:145:SER:HB2	2.19	0.43
1:B:28:ILE:HD11	1:B:32:SER:O	2.18	0.42
1:B:360:ASN:H	1:B:364:ASN:HD22	1.65	0.42
1:B:188:THR:O	1:B:189:TRP:HB2	2.18	0.42
1:B:183:ASP:OD1	1:B:185:GLN:HG2	2.19	0.42



Continued from previous page...

A + 1		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)
1:A:110:HIS:CE1	1:A:146:TYR:O	2.73	0.42
1:B:148:LEU:HD13	1:B:178:ILE:HG22	2.02	0.42
1:A:79:GLU:HG3	1:A:80:GLU:H	1.85	0.42
1:A:148:LEU:CD1	1:A:180:ASP:HA	2.32	0.42
1:B:125:ARG:HA	1:B:248:CYS:HB3	2.02	0.42
1:A:146:TYR:HB3	1:A:149:LYS:HB3	2.01	0.42
1:A:127:VAL:HG23	1:A:277:MSE:HG3	2.02	0.41
1:B:386:ILE:CG2	1:B:436:MSE:HB3	2.50	0.41
1:A:181:ILE:H	1:A:181:ILE:HG13	1.58	0.41
1:A:55:ILE:HD12	1:A:184:ALA:HB1	2.03	0.41
1:B:148:LEU:CD1	1:B:180:ASP:HA	2.41	0.41
1:A:126:VAL:HA	1:A:276:ILE:O	2.21	0.41
1:A:199:LEU:H	1:A:199:LEU:HD13	1.86	0.41
1:A:348:VAL:HG13	1:A:352:ASP:HB2	2.03	0.41
1:A:288:TRP:CZ3	1:A:306:PHE:HB3	2.56	0.41
1:A:414:VAL:HA	1:A:421:LEU:HD23	2.04	0.40
1:B:258:LEU:HD22	1:B:264:GLU:HG2	2.04	0.40
1:A:193:LEU:HD23	1:A:202:ASN:HB2	2.02	0.40
1:B:171:GLY:HA3	5:B:2517:HOH:O	2.20	0.40
1:B:245:ARG:HD3	5:B:2513:HOH:O	2.20	0.40
1:A:125:ARG:HG2	1:A:248:CYS:CB	2.51	0.40
1:A:126:VAL:HG13	1:A:276:ILE:HG12	2.03	0.40
1:B:265:LEU:HB2	1:B:295:PHE:CZ	2.57	0.40
1:A:158:ILE:HD13	1:A:272:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	426/460 (93%)	380 (89%)	42 (10%)	4 (1%)	17 51



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	В	426/460 (93%)	394 (92%)	24 (6%)	8 (2%)	8	32
2	С	20/110 (18%)	20 (100%)	0	0	100	100
2	D	23/110 (21%)	23 (100%)	0	0	100	100
All	All	895/1140 (78%)	817 (91%)	66 (7%)	12 (1%)	12	41

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	419	ARG
1	В	42	GLY
1	В	62	VAL
1	В	261	ASN
1	A	96	THR
1	A	340	ALA
1	A	81	THR
1	В	41	GLU
1	В	209	ASN
1	В	365	GLU
1	В	286	PRO
1	A	366	VAL

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	385/407 (95%)	356 (92%)	29 (8%)	13	40
1	В	385/407 (95%)	353 (92%)	32 (8%)	11	35
2	С	20/97~(21%)	20 (100%)	0	100	100
2	D	23/97~(24%)	23 (100%)	0	100	100
All	All	813/1008 (81%)	752 (92%)	61 (8%)	13	40

All (61) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	18	THR.
1	A	26	LEU
1	A A	44	LEU
1	A	53	VAL
1	A A A	112	MSE
1	A	113	LEU
1	A	115	LYS
1	A	127	VAL
1	A	132	SER
1	A	139	LEU
1	A	145	SER
1	A	146	TYR
1	A	168	THR
1	A A A A A A A A A A A A A A A A A A A	181	ILE
1	A	199	LEU
1	A	203	LYS
1	A	206	MSE
1	A	214	ARG
1	A	219	LYS LEU
1	A	221	
1	A	227	SER
1	A	245	ARG
1	A	255	ILE
1	A	281	CYS
1	A	330	ARG
1	A	348	VAL
1	A	361	VAL
1	A	371	LEU
1	A	438	LEU
1	В	22	GLU
1	В	44	LEU
1	В	49	ASN
1	В	53	VAL
1	В	94	THR
1	В	112	MSE
1	В	113	LEU
1	В	115	LYS
1	В	127	VAL
1	В	132	SER
1	В	136	LYS
1	В	179	SER
1	В	181	ILE
1	В	203	LYS



Continued from previous page...

Mol	Chain	Res	Type
1	В	219	LYS
1	В	221	LEU
1	В	232	VAL
1	В	245	ARG
1	В	252	THR
1	В	259	ASP
1	В	275	ASN
1	В	276	ILE
1	В	278	LEU
1	В	321	LYS
1	В	365	GLU
1	В	368	ARG
1	В	371	LEU
1	В	377	THR
1	В	406	LEU
1	В	410	LEU
1	В	419	ARG
1	В	444	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such side chains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	110	HIS
1	A	120	ASN
1	A	133	GLN
1	A	154	GLN
1	A	242	GLN
1	A	273	ASN
1	A	302	ASN
1	A	364	ASN
1	В	49	ASN
1	В	120	ASN
1	В	228	GLN
1	В	266	HIS
1	В	267	HIS
1	В	273	ASN
1	В	275	ASN
1	В	364	ASN
1	В	382	GLN
2	С	488	GLN
2	D	476	ASN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain		ain Res	Res Link	Во	Bond lengths			Bond angles		
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
4	ATP	В	1600	3	26,33,33	1.01	2 (7%)	31,52,52	1.37	4 (12%)	
4	ATP	A	600	3	26,33,33	0.98	1 (3%)	31,52,52	1.55	4 (12%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

$\mathbf{Mol}$	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	В	1600	3	-	3/18/38/38	0/3/3/3
4	ATP	A	600	3	-	0/18/38/38	0/3/3/3

#### All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}( ext{\AA})$
4	В	1600	ATP	C5-C4	2.64	1.47	1.40



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(A)
4	A	600	ATP	C5-C4	2.48	1.47	1.40
4	В	1600	ATP	C2-N3	2.15	1.35	1.32

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
4	A	600	ATP	PA-O3A-PB	-3.90	119.43	132.83
4	A	600	ATP	N3-C2-N1	-3.58	123.08	128.68
4	В	1600	ATP	PA-O3A-PB	-3.40	121.16	132.83
4	A	600	ATP	PB-O3B-PG	-3.39	121.19	132.83
4	В	1600	ATP	N3-C2-N1	-3.20	123.67	128.68
4	A	600	ATP	C4-C5-N7	-2.60	106.69	109.40
4	В	1600	ATP	C4-C5-N7	-2.52	106.78	109.40
4	В	1600	ATP	PB-O3B-PG	-2.28	124.99	132.83

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	В	1600	ATP	PB-O3A-PA-O1A
4	В	1600	ATP	PB-O3A-PA-O2A
4	В	1600	ATP	C5'-O5'-PA-O1A

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	422/460 (91%)	0.26	25 (5%) 22 13	26, 55, 90, 133	0
1	В	422/460 (91%)	-0.03	12 (2%) 53 36	25, 45, 69, 98	0
2	С	22/110 (20%)	0.01	0 100 100	31, 42, 62, 77	0
2	D	25/110 (22%)	0.28	1 (4%) 38 25	27, 39, 71, 90	0
All	All	891/1140 (78%)	0.11	38 (4%) 35 22	25, 49, 84, 133	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	499	SER	6.7
1	A	284	THR	5.6
1	A	18	THR	5.3
1	В	18	THR	3.4
1	A	420	LYS	3.3
1	A	197	ALA	3.3
1	В	62	VAL	3.2
1	В	366	VAL	3.1
1	A	419	ARG	3.1
1	A	366	VAL	3.0
1	A	417	LYS	3.0
1	A	100	ASN	2.9
1	A	63	ASP	2.7
1	В	445	GLU	2.6
1	В	197	ALA	2.6
1	В	41	GLU	2.5
1	A	360	ASN	2.4
1	В	284	THR	2.4
1	A	41	GLU	2.3
1	В	21	ASN	2.3
1	В	394	ARG	2.3



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	19	GLY	2.3
1	A	359	SER	2.2
1	A	421	LEU	2.2
1	A	362	PHE	2.2
1	В	283	GLU	2.2
1	A	80	GLU	2.1
1	A	94	THR	2.1
1	A	367	GLY	2.1
1	В	306	PHE	2.1
1	A	351	GLU	2.1
1	A	40	ALA	2.1
1	A	444	LEU	2.0
1	A	413	GLU	2.0
1	В	365	GLU	2.0
1	A	196	GLY	2.0
1	A	414	VAL	2.0
1	A	93	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
3	MG	В	2500	1/1	0.94	0.10	15,15,15,15	0
3	MG	A	1500	1/1	0.95	0.12	27,27,27,27	0
4	ATP	A	600	31/31	0.96	0.19	41,44,45,45	0
4	ATP	В	1600	31/31	0.97	0.16	37,39,40,41	0


The following is a graphical depiction of the model fit to experimental electron density of all



instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







# 6.5 Other polymers (i)

There are no such residues in this entry.

