Full wwPDB X-ray Structure Validation Report (i)

Jan 7, 2024 – 09:08 am GMT

PDB ID	:	5NQZ
Title	:	Structure of a fHbp(V1.1):PorA(P1.16) chimera. Fusion at fHbp position 309.
Authors	:	Johnson, S.; Hollingshead, S.; Lea, S.M.; Tang, C.M.
Deposited on	:	2017-04-21
Resolution	:	1.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	2%, 92%, 6%
1	B	269	4%, 91%, 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	406	-	-	X	-
3	ACT	A	408	-	X	X	-
3	ACT	A	412	-	-	X	-

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4631 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Factor H binding protein,Major outer membrane protein P.IA,Factor H binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	267	2036	1265	369	401	1	0	0	0
1	B	263	2010	1249	365	395	1	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	72	MET	-	initiating methionine	UNP Q9JXV4
A	309	VAL	ASN	conflict	UNP Q9JXV4
A	321	LEU	-	expression tag	UNP Q9JXV4
A	322	GLU	-	expression tag	UNP Q9JXV4
A	323	HIS	-	expression tag	UNP Q9JXV4
A	324	HIS	-	expression tag	UNP Q9JXV4
A	325	HIS	-	expression tag	UNP Q9JXV4
A	326	HIS	-	expression tag	UNP Q9JXV4
A	327	HIS	-	expression tag	UNP Q9JXV4
A	328	HIS	-	expression tag	UNP Q9JXV4
B	72	MET	-	initiating methionine	UNP Q9JXV4
B	309	VAL	ASN	conflict	UNP Q9JXV4
B	321	LEU	-	expression tag	UNP Q9JXV4
B	322	GLU	-	expression tag	UNP Q9JXV4
B	323	HIS	-	expression tag	UNP Q9JXV4
B	324	HIS	-	expression tag	UNP Q9JXV4
B	325	HIS	-	expression tag	UNP Q9JXV4
B	326	HIS	-	expression tag	UNP Q9JXV4
B	327	HIS	-	expression tag	UNP Q9JXV4
B	328	HIS	-	expression tag	UNP Q9JXV4

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	A	6	6	6	0	0
			Total	Zn		
2	B	3	3	3	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	4	2	2	0	0
			Total	C	O		
3	A	1	4	2	2	0	0
			Total	C	O		
3	A	1	4	2	2	0	0
			Total	C	O		
3	A	1	4	2	2	0	0
			Total	C	O		
3	A	1	4	2	2	0	0
			Total	C	O		
3	B	1	4	2	2	0	0
			Total	C	O		
3	B	1	4	2	2	0	0
			Total	C	O		
3	B	1	4	2	2	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	4	2	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	286	286	286	0	0
			Total	O		
5	B	254	254	254	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Factor H binding protein,Major outer membrane protein P.IA,Factor H binding protein

Chain A:





• Molecule 1: Factor H binding protein,Major outer membrane protein P.IA,Factor H binding protein

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	91.37Å 76.27Å 83.90Å 90.00° 91.26° 90.00°	Depositor
Resolution (Å)	58.55 – 1.63 58.55 – 1.63	Depositor EDS
% Data completeness (in resolution range)	98.5 (58.55-1.63) 98.5 (58.55-1.63)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.54 (at 1.63Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.162 , 0.194 0.174 , 0.204	Depositor DCC
R_{free} test set	3445 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å^2)	18.6	Xtriage
Anisotropy	0.340	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 35.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.188 for -h,-k,l	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	4631	wwPDB-VP
Average B, all atoms (Å^2)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.07	0/2069	1.07	11/2782 (0.4%)
1	B	1.09	0/2043	1.03	7/2746 (0.3%)
All	All	1.08	0/4112	1.05	18/5528 (0.3%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	105	VAL	CG1-CB-CG2	7.14	122.33	110.90
1	A	304	GLU	OE1-CD-OE2	7.13	131.86	123.30
1	A	90	ASP	CB-CG-OD1	7.08	124.67	118.30
1	B	90	ASP	CB-CG-OD1	6.97	124.58	118.30
1	B	105	VAL	CB-CA-C	-6.52	99.02	111.40
1	A	105	VAL	CB-CA-C	-6.47	99.10	111.40
1	A	181	ASP	CB-CG-OD1	-6.44	112.50	118.30
1	A	105	VAL	CG1-CB-CG2	6.41	121.16	110.90
1	A	105	VAL	CA-CB-CG2	5.79	119.58	110.90
1	B	195	ARG	NE-CZ-NH1	5.76	123.18	120.30
1	A	142	ASP	CB-CG-OD2	-5.72	113.15	118.30
1	A	269	ARG	NE-CZ-NH1	5.43	123.01	120.30
1	B	105	VAL	CA-CB-CG2	5.30	118.86	110.90
1	B	277	VAL	CA-CB-CG1	5.17	118.65	110.90
1	A	202	GLU	OE1-CD-OE2	-5.02	117.27	123.30
1	A	272	VAL	CG1-CB-CG2	-5.02	102.87	110.90
1	B	202	GLU	OE1-CD-OE2	-5.01	117.28	123.30
1	A	145	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2036	0	2005	17	0
1	B	2010	0	1980	17	0
2	A	6	0	0	0	0
2	B	3	0	0	0	0
3	A	20	0	15	17	0
3	B	12	0	9	1	0
4	A	4	0	6	0	0
5	A	286	0	0	10	1
5	B	254	0	0	5	1
All	All	4631	0	4015	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:LEU:HB3	3:A:412:ACT:H2	1.13	1.08
1:A:89:LEU:CB	3:A:412:ACT:H2	1.93	0.98
1:B:109:GLU:OE1	5:B:501:HOH:O	1.91	0.87
1:A:280:ASN:HA	5:A:505:HOH:O	1.79	0.82
1:A:89:LEU:HB3	3:A:412:ACT:CH3	2.06	0.80
1:B:218:ARG:HD2	1:B:321:LEU:HD21	1.66	0.77
3:A:412:ACT:H1	5:A:693:HOH:O	1.87	0.74
3:A:412:ACT:H3	1:B:294:GLY:HA3	1.72	0.71
1:A:248:HIS:HE1	5:A:738:HOH:O	1.75	0.69
1:B:279:TYR:OH	5:B:502:HOH:O	2.11	0.68
1:A:89:LEU:HD23	3:A:412:ACT:O	1.94	0.67
3:A:412:ACT:H3	1:B:294:GLY:CA	2.26	0.66
3:A:408:ACT:H1	5:A:752:HOH:O	1.95	0.65
1:A:112:LYS:NZ	5:A:503:HOH:O	2.29	0.64
1:B:218:ARG:CD	1:B:321:LEU:HD21	2.31	0.59
1:A:112:LYS:HE2	1:A:119:GLU:CD	2.23	0.59
3:A:412:ACT:H3	1:B:294:GLY:N	2.19	0.57
1:B:112:LYS:NZ	5:B:505:HOH:O	2.37	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:408:ACT:CH3	5:A:752:HOH:O	2.53	0.54
1:A:313:HIS:NE2	5:A:506:HOH:O	2.33	0.54
3:A:412:ACT:H3	1:B:294:GLY:H	1.73	0.53
1:B:152:GLN:HA	5:B:522:HOH:O	2.09	0.52
1:A:145:ARG:HD2	5:A:513:HOH:O	2.09	0.52
1:A:145:ARG:CD	5:A:513:HOH:O	2.59	0.50
1:A:89:LEU:CG	3:A:412:ACT:H2	2.41	0.49
3:A:412:ACT:CH3	1:B:294:GLY:H	2.25	0.49
3:B:405:ACT:C	3:B:406:ACT:H2	2.43	0.49
1:A:294:GLY:H	3:A:406:ACT:H3	1.79	0.47
1:A:306:LYS:HG2	1:A:311:ILE:CD1	2.44	0.47
1:B:304:GLU:HG2	1:B:313:HIS:CE1	2.49	0.47
1:B:218:ARG:HE	1:B:321:LEU:HD11	1.81	0.45
1:A:306:LYS:HG2	1:A:311:ILE:HD13	1.99	0.45
1:B:322:GLU:OE1	1:B:325:HIS:NE2	2.48	0.45
3:A:406:ACT:O	1:B:89:LEU:HD23	2.17	0.45
3:A:412:ACT:C	1:B:294:GLY:H	2.30	0.45
1:A:145:ARG:NE	5:A:513:HOH:O	2.50	0.44
1:B:78:GLY:N	5:B:513:HOH:O	2.52	0.43
1:A:89:LEU:HD23	3:A:412:ACT:C	2.48	0.43

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:710:HOH:O	5:B:621:HOH:O[4_553]	2.13	0.07

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	265/269 (98%)	257 (97%)	7 (3%)	1 (0%)	34 15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	261/269 (97%)	253 (97%)	8 (3%)	0	100	100
All	All	526/538 (98%)	510 (97%)	15 (3%)	1 (0%)	47	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	150	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/212 (99%)	209 (100%)	1 (0%)	88	80
1	B	208/212 (98%)	207 (100%)	1 (0%)	88	80
All	All	418/424 (99%)	416 (100%)	2 (0%)	88	80

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	VAL
1	B	308	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 9 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	ACT	A	412	2	3,3,3	1.96	1 (33%)	3,3,3	1.31	0
3	ACT	A	407	2	3,3,3	0.96	0	3,3,3	0.79	0
3	ACT	A	408	2	3,3,3	2.36	1 (33%)	3,3,3	1.96	2 (66%)
3	ACT	B	404	2	3,3,3	0.66	0	3,3,3	0.80	0
4	EDO	A	410	-	3,3,3	0.52	0	2,2,2	0.39	0
3	ACT	A	409	2	3,3,3	0.73	0	3,3,3	0.45	0
3	ACT	A	406	2	3,3,3	2.13	2 (66%)	3,3,3	1.28	0
3	ACT	B	405	2	3,3,3	0.94	0	3,3,3	0.82	0
3	ACT	B	406	2	3,3,3	1.50	1 (33%)	3,3,3	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	410	-	-	1/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	408	ACT	OXT-C	-4.03	1.11	1.30
3	A	412	ACT	OXT-C	-3.39	1.14	1.30
3	A	406	ACT	OXT-C	-2.62	1.18	1.30
3	B	406	ACT	OXT-C	-2.34	1.19	1.30
3	A	406	ACT	O-C	2.26	1.32	1.22

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	408	ACT	OXT-C-O	-2.55	112.65	122.05
3	A	408	ACT	OXT-C-CH3	2.22	124.36	115.18

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	410	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	412	ACT	13	0
3	A	408	ACT	2	0
3	A	406	ACT	2	0
3	B	405	ACT	1	0
3	B	406	ACT	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	267/269 (99%)	-0.53	5 (1%)	66	67	12, 19, 50, 75	0
1	B	263/269 (97%)	-0.45	10 (3%)	40	38	12, 20, 55, 86	0
All	All	530/538 (98%)	-0.49	15 (2%)	53	51	12, 20, 53, 86	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	327	HIS	3.3
1	B	150	ASP	2.9
1	B	323	HIS	2.8
1	A	77	ILE	2.7
1	A	185	SER	2.5
1	B	185	SER	2.5
1	B	308(F)	THR	2.4
1	A	75	ALA	2.3
1	B	308(E)	ASP	2.3
1	A	150	ASP	2.2
1	B	322	GLU	2.2
1	B	186	GLY	2.2
1	B	152	GLN	2.1
1	B	321	LEU	2.1
1	A	308(E)	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	EDO	A	410	4/4	0.64	0.20	56,60,67,67	0
3	ACT	B	404	4/4	0.91	0.10	28,28,33,33	0
3	ACT	A	409	4/4	0.91	0.14	53,54,59,59	0
3	ACT	B	405	4/4	0.95	0.14	31,39,39,42	0
3	ACT	A	406	4/4	0.95	0.12	16,22,28,44	0
2	ZN	B	403	1/1	0.96	0.04	41,41,41,41	0
3	ACT	A	407	4/4	0.97	0.11	25,33,34,41	0
3	ACT	B	406	4/4	0.98	0.13	25,33,47,49	0
3	ACT	A	412	4/4	0.98	0.22	18,31,44,45	0
3	ACT	A	408	4/4	0.99	0.12	20,26,38,39	0
2	ZN	A	405	1/1	0.99	0.09	49,49,49,49	0
2	ZN	A	411	1/1	0.99	0.04	22,22,22,22	0
2	ZN	B	402	1/1	0.99	0.02	30,30,30,30	0
2	ZN	A	401	1/1	0.99	0.04	22,22,22,22	0
2	ZN	A	403	1/1	0.99	0.02	27,27,27,27	0
2	ZN	A	404	1/1	0.99	0.07	42,42,42,42	0
2	ZN	B	401	1/1	1.00	0.04	22,22,22,22	0
2	ZN	A	402	1/1	1.00	0.03	19,19,19,19	0

6.5 Other polymers (i)

There are no such residues in this entry.