

# Full wwPDB X-ray Structure Validation Report (i)

#### Dec 16, 2023 – 06:53 PM EST

PDB ID	:	3NTB
Title	:	Structure of 6-methylthic naproxen analog bound to mCOX-2.
Authors	:	Duggan, K.C.; Musee, J.; Walters, M.J.; Harp, J.M.; Kiefer, J.R.; Oates, J.A.;
		Marnett, L.J.
Deposited on	:	2010-07-03
Resolution	:	2.27  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	Similar resolution		
	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$		
$R_{free}$	130704	$6980 \ (2.30-2.26)$		
Clashscore	141614	7711 (2.30-2.26)		
Ramachandran outliers	138981	7597 (2.30-2.26)		
Sidechain outliers	138945	7598 (2.30-2.26)		
RSRZ outliers	127900	6849 (2.30-2.26)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			9%		
1	А	587	89%	5%	6%
			11%		
1	В	587	88%	6%	6%
			8%		
1	С	587	88%	6%	6%
			7%		
1	D	587	88%	6%	6%
2	E	3	100%		



Mol	Chain	Length	Quality of chain
2	F	3	100%
2	G	3	100%
2	Н	3	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	Е	3	-	-	-	Х
2	NAG	F	3	-	-	-	Х
2	NAG	G	3	-	-	-	Х
2	NAG	Н	3	-	-	-	Х
4	NAG	А	661	Х	-	-	Х
4	NAG	В	661	Х	-	-	-
4	NAG	С	661	Х	-	-	-
4	NAG	С	681	-	-	-	Х
4	NAG	D	661	Х	-	-	Х



# 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 19636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Δ	559	Total	С	Ν	0	$\mathbf{S}$	0	0	0
1	A	552	4474	2885	750	814	25	0	0	0
1	В	551	Total	С	Ν	0	S	0	0	0
1	I D	551	4465	2880	748	812	25	0	0	0
1	C	559	Total	С	Ν	0	S	0	0	0
	002	4474	2885	750	814	25	0	0	0	
1 D	552	Total	С	Ν	0	S	0	0	0	
		4474	2885	750	814	25	0		0	

• Molecule 1 is a protein called Prostaglandin-endoperoxide synthase 2.

• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace			
2	F	3	Total	С	Ν	0	0	Ο	0	
2		0	42	24	3	15	0	0	U	
2	F	3	Total	С	Ν	0	0	0	0	
2	Ľ		42	24	3	15	0	0	0	
9	С	9	Total	С	Ν	0	0	0	0	
Z G	5	42	24	3	15	0	0	U		
2 H	9	Total	С	Ν	0	0	0	0		
	П	ა	42	24	3	15	0	0	U	

• Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).





Mol	Chain	Residues		Ate	$\mathbf{oms}$		ZeroOcc	AltConf		
3	2 1	1	Total	С	Fe	Ν	0	0	0	
0	A	1	43	34	1	4	4	0		
2	3 B	1	Total	С	Fe	Ν	0	0	0	
0			43	34	1	4	4		0	
2	С	C 1	Total	С	Fe	Ν	0	0	0	
3 U	1	43	34	1	4	4	0	0		
2 D	1	Total	С	Fe	Ν	Ο	0	0		
0	D	1	43	34	1	4	4	0	0	

• Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
1	Λ	1	Total C N O	0	0
4	Л	I	14  8  1  5	0	0
4	Δ	1	Total C N O	0	0
т	11	1	14 8 1 5	0	0
4	В	1	Total C N O	0	0
т	D	1	14 8 1 5	0	0
4	В	1	Total C N O	0	0
	D	I	14 8 1 5	0	0
4	С	1	Total C N O	0	0
	0	I	14 8 1 5	0	0
4	С	1	Total C N O	0	0
	0	1	14 8 1 5	0	0
4	Л	1	Total C N O	0	0
		1	14 8 1 5		0
4	П	1	Total C N O	0	0
<b>T</b>		1	14 8 1 5		0

• Molecule 5 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula:  $C_{14}H_{28}O_6$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total         C         O           20         14         6	0	0
5	А	1	Total         C         O           20         14         6	0	0
5	В	1	Total         C         O           20         14         6	0	0
5	С	1	Total         C         O           20         14         6	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total         C         O           20         14         6	0	0
5	D	1	Total         C         O           20         14         6	0	0

• Molecule 6 is (2S)-2-[6-(methyl sulfanyl)naphthalen-2-yl]propanoic acid (three-letter code: T1N) (formula:  $\rm C_{14}H_{14}O_2S).$ 



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	А	1	Total	С	0	$\mathbf{S}$	0	0
	11	Ť	17	14	2	1	Ŭ	0
6	В	1	Total	С	Ο	$\mathbf{S}$	0	0
0	D	L	17	14	2	1	0	0
6	C	1	Total	С	Ο	S	0	0
0	U	L	17	14	2	1	0	0
6	Л	1	Total	С	Ο	S	0	0
0	D	L	17	14	2	1	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	278	Total O 278 278	0	0
7	В	245	Total O 245 245	0	0
7	С	291	Total         O           291         291	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	295	Total O 295 295	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Prostaglandin-endoperoxide synthase 2





#### LYS ARG ARG SER THR GLU LEU

• Molecule 1: Prostaglandin-endoperoxide synthase 2



# 8

A33	C41	D53	Q54	K56 K56	• [2] [2]		F74	R77	┝	L81	L82	Y91		N101	N1 05		F107	M1 13		Y122	K137		D158	V165	1 1 7 1	1 1 1 1	E186	0192		F1 98	Y234		D.239	L252		8929	D268	1274		H278	P280	E281
-	Q284 F285	A286 V287	G288	4289 E290	V291 F292		P296	L316	K317	Q318	1 378	F329	<mark>0330</mark>	T 225		V344		Y 348	L352		K360 F361	D362	T 377	A378	S379	000	Y385	H388	P389		Q400		1409	F422		Y 466	L472	K4/3	L481		E484 K485	-
D497	V500	A506		0101	D515	I517	F518	E524		A527	P528	Q583	ASP	PRO	PRO	THR	LYS	AL.A	THR	ILE	ASN ALA	SER	ALA SFR	HIS	SER	LEU	ASP	ASP	ASN	PRO THR	VAL	LEU	LYS	ARG	ARG	THR	GLU	LEU				

 $\bullet \ {\rm Molecule \ 2: \ 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose}$ 

Chain E:	100%		-		
NAG1 NAG2 NAG3					
• Molecule 2: 2-acetar opyranose-(1-4)-2-acet	nido-2-deoxy-beta-D-glucopyra amido-2-deoxy-beta-D-glucopyr	nose-(1-4)-2-acetam ranose	ido-2-deox	y-beta-D-glu	ıc
Chain F:	100%		-		
NAG1 NAG2 NAG3					

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:

100%

#### NAG1 NAG2 NAG3 NAG3

 $\bullet$  Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose (1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



# Chain H:

100%

NAG1 NAG2 NAG3



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	181.20Å 134.22Å 121.99Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution(A)	20.48 - 2.27	Depositor
Resolution (A)	19.94 - 2.20	EDS
% Data completeness	98.0 (20.48-2.27)	Depositor
(in resolution range)	96.7 (19.94-2.20)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$< I/\sigma(I) > 1$	$2.23 (at 2.19 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
D D .	0.231 , $0.266$	Depositor
$n, n_{free}$	0.242 , $0.275$	DCC
$R_{free}$ test set	14625  reflections  (10.04%)	wwPDB-VP
Wilson B-factor $(Å^2)$	40.1	Xtriage
Anisotropy	0.349	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, 38.9	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	19636	wwPDB-VP
Average B, all atoms $(Å^2)$	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 50.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.8750e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, T1N, HEM, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles				
MIOI	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5			
1	А	0.34	0/4601	0.47	0/6239			
1	В	0.34	0/4592	0.46	0/6227			
1	С	0.34	0/4601	0.47	0/6239			
1	D	0.34	0/4601	0.46	0/6239			
All	All	0.34	0/18395	0.47	0/24944			

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4474	0	4373	12	0
1	В	4465	0	4365	21	0
1	С	4474	0	4373	20	0
1	D	4474	0	4373	18	0
2	Е	42	0	37	0	0
2	F	42	0	37	0	0
2	G	42	0	37	0	0
2	Н	42	0	37	0	0
3	А	43	0	30	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	43	0	30	0	0
3	С	43	0	30	2	0
3	D	43	0	30	0	0
4	А	28	0	26	0	0
4	В	28	0	26	0	0
4	С	28	0	26	0	0
4	D	28	0	26	0	0
5	А	40	0	56	0	0
5	В	20	0	28	0	0
5	С	20	0	28	0	0
5	D	40	0	56	0	0
6	А	17	0	13	0	0
6	В	17	0	13	0	0
6	С	17	0	13	0	0
6	D	17	0	13	0	0
7	А	278	0	0	0	0
7	В	245	0	0	0	0
7	C	291	0	0	0	0
7	D	295	0	0	0	0
All	All	19636	0	18076	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:472:LEU:HD21	1:A:524:GLU:HG3	1.82	0.60
1:B:352:LEU:HD11	1:B:518:PHE:CE2	2.38	0.58
1:D:198:PHE:CZ	1:D:352:LEU:HD13	2.39	0.58
1:D:198:PHE:HZ	1:D:352:LEU:HD13	1.69	0.57
1:B:352:LEU:HD11	1:B:518:PHE:HE2	1.71	0.55
1:C:447:VAL:HG13	3:C:619:HEM:HBA1	1.88	0.55
1:B:123:LEU:O	1:B:469:ARG:NH2	2.40	0.55
1:B:151:ALA:O	1:B:469:ARG:NH1	2.40	0.54
1:C:352:LEU:HD22	1:C:518:PHE:HE2	1.72	0.54
1:B:472:LEU:HD21	1:B:524:GLU:HG3	1.88	0.54
1:B:380:GLU:HG2	1:B:466:TYR:CE1	2.44	0.53
1:A:289:GLN:HG2	1:A:292:PHE:CE1	2.44	0.52
1:C:382:ASN:O	1:C:386:HIS:HD2	1.93	0.52
1:C:484:GLU:HG2	1:C:485:LYS:H	1.74	0.52



	lo uo puge	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:198:PHE:CZ	1:B:352:LEU:HD13	2.45	0.51
1:D:472:LEU:HD11	1:D:524:GLU:HG3	1.91	0.51
1:C:472:LEU:HD11	1:C:524:GLU:HG3	1.94	0.49
1:B:209:PHE:HB2	1:B:377:ILE:HG13	1.95	0.49
1:C:198:PHE:HZ	1:C:352:LEU:HD21	1.77	0.48
1:C:543:GLN:O	1:D:137:LYS:HE2	2.13	0.48
1:D:281:GLU:HA	1:D:284:GLN:HE21	1.79	0.48
1:A:470:PHE:CD2	1:A:525:LEU:HD22	2.48	0.48
1:C:209:PHE:HB2	1:C:377:ILE:HG13	1.96	0.48
1:C:389:PRO:HB2	1:C:434:VAL:HA	1.95	0.48
1:A:527:ALA:HB3	1:A:528:PRO:HD3	1.95	0.47
1:A:380:GLU:HG2	1:A:466:TYR:CE1	2.51	0.46
1:A:543:GLN:O	1:B:137:LYS:HE2	2.16	0.46
1:D:380:GLU:HG2	1:D:466:TYR:CE1	2.51	0.46
1:A:389:PRO:HB2	1:A:434:VAL:HA	1.99	0.45
1:C:484:GLU:HG2	1:C:485:LYS:N	2.31	0.45
1:B:295:VAL:HG12	1:B:297:GLY:H	1.81	0.45
1:A:344:VAL:HA	1:A:348:TYR:HB3	1.98	0.45
1:C:470:PHE:CD2	1:C:525:LEU:HD22	2.52	0.44
1:B:389:PRO:HB2	1:B:434:VAL:HA	1.99	0.44
1:D:506:ALA:O	1:D:510:GLU:HB2	2.17	0.44
1:A:181:VAL:HG21	1:A:491:LEU:HD21	1.99	0.44
1:D:527:ALA:HB3	1:D:528:PRO:HD3	2.01	0.43
1:B:181:VAL:HG21	1:B:491:LEU:HD21	2.00	0.43
1:C:206:THR:HB	1:C:210:PHE:CD2	2.53	0.43
1:C:138:SER:HB2	1:D:330:GLN:HB3	2.00	0.43
1:C:380:GLU:HG2	1:C:466:TYR:CE1	2.54	0.43
1:C:388:HIS:N	1:C:389:PRO:CD	2.81	0.43
1:B:482:THR:HG22	1:B:509:VAL:HG12	1.99	0.43
1:C:387:TRP:HB2	3:C:619:HEM:HAC	2.01	0.43
1:D:344:VAL:HA	1:D:348:TYR:HB3	2.00	0.43
1:D:497:ASP:HB3	1:D:500:VAL:HG23	2.01	0.43
1:B:388:HIS:N	1:B:389:PRO:CD	2.82	0.43
1:C:274:ILE:HD12	1:C:291:VAL:HG12	2.02	0.42
1:B:484:GLU:HG2	1:B:485:LYS:H	1.84	0.42
1:B:506:ALA:O	1:B:510:GLU:HB2	2.20	0.42
1:D:289:GLN:HG2	1:D:292:PHE:CE1	2.55	0.41
1:D:388:HIS:N	1:D:389:PRO:CD	2.83	0.41
1:C:344:VAL:HA	1:C:348:TYR:HB3	2.01	0.41
1:D:484:GLU:HG2	1:D:485:LYS:H	1.85	0.41
1:A:335:ILE:HA	1:A:559:ILE:HD11	2.02	0.41



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:388:HIS:N	1:A:389:PRO:CD	2.82	0.41
1:B:470:PHE:CD2	1:B:525:LEU:HD22	2.56	0.41
1:D:352:LEU:HD11	1:D:518:PHE:CE2	2.56	0.41
1:B:495:TYR:HE1	1:B:502:GLU:HG3	1.85	0.41
1:A:473:LYS:HA	1:A:474:PRO:HD3	1.96	0.41
1:D:360:LYS:HE3	1:D:362:ASP:HB2	2.03	0.41
1:B:198:PHE:HZ	1:B:352:LEU:HD13	1.86	0.41
1:B:206:THR:HG21	1:B:385:TYR:CE2	2.56	0.40
1:C:287:VAL:HG11	1:C:302:ALA:HB1	2.02	0.40
1:D:274:ILE:HD12	1:D:291:VAL:HG12	2.03	0.40
1:C:518:PHE:HB3	1:C:522:MET:HB3	2.03	0.40
1:B:444:VAL:HG12	1:B:444:VAL:O	2.22	0.40
1:D:192:GLN:HG3	1:D:516:ALA:HA	2.04	0.40

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	550/587~(94%)	536~(98%)	14 (2%)	0	100	100
1	В	549/587~(94%)	534~(97%)	14 (3%)	1 (0%)	47	57
1	С	550/587~(94%)	537~(98%)	13 (2%)	0	100	100
1	D	550/587~(94%)	534 (97%)	16 (3%)	0	100	100
All	All	2199/2348~(94%)	2141 (97%)	57 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	409	TYR



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	493/525~(94%)	488 (99%)	5 (1%)	76	86
1	В	492/525~(94%)	488 (99%)	4 (1%)	81	90
1	С	493/525~(94%)	487 (99%)	6 (1%)	71	82
1	D	493/525~(94%)	488 (99%)	5 (1%)	76	86
All	All	1971/2100~(94%)	1951 (99%)	20~(1%)	76	86

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	176	GLU
1	А	271	VAL
1	А	295	VAL
1	А	385	TYR
1	А	484	GLU
1	В	317	LYS
1	В	385	TYR
1	В	405	LYS
1	В	484	GLU
1	С	107	PHE
1	С	169	LYS
1	С	385	TYR
1	С	409	TYR
1	С	472	LEU
1	С	484	GLU
1	D	56	LYS
1	D	107	PHE
1	D	171	LEU
1	D	385	TYR
1	D	484	GLU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such side chains are listed below:



Mol	Chain	Res	Type
1	А	327	GLN
1	С	369	GLN
1	С	386	HIS
1	С	583	GLN
1	D	284	GLN
1	D	369	GLN
1	D	386	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	I Type Chain Res		Tinle	Bo	Bond lengths			Bond angles		
	туре	Chain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	Е	1	2,1	14,14,15	0.49	0	17,19,21	0.94	1 (5%)
2	NAG	Е	2	2	14,14,15	0.50	0	17,19,21	1.66	6 (35%)
2	NAG	Е	3	2	14,14,15	0.62	0	17,19,21	1.04	1 (5%)
2	NAG	F	1	2,1	14,14,15	0.50	0	17,19,21	0.81	1 (5%)
2	NAG	F	2	2	14,14,15	0.50	0	17,19,21	1.15	2 (11%)
2	NAG	F	3	2	14,14,15	0.52	0	17,19,21	1.69	3 (17%)
2	NAG	G	1	2,1	14,14,15	0.50	0	17,19,21	0.88	1 (5%)
2	NAG	G	2	2	14,14,15	0.53	0	17,19,21	1.21	1 (5%)
2	NAG	G	3	2	14,14,15	0.51	0	17,19,21	1.55	2 (11%)
2	NAG	Н	1	2,1	14,14,15	0.49	0	17,19,21	0.90	1 (5%)



Mal Trme	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Dec	Tinle	Bo	ond leng	$_{\rm ths}$	В	ond ang	les
	туре	Chain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2															
2	NAG	Н	2	2	14,14,15	0.53	0	17,19,21	1.17	2 (11%)															
2	NAG	Н	3	2	14,14,15	0.56	0	17,19,21	1.87	3 (17%)															

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	Е	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	Е	2	2	-	1/6/23/26	0/1/1/1
2	NAG	Е	3	2	-	5/6/23/26	0/1/1/1
2	NAG	F	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	4/6/23/26	0/1/1/1
2	NAG	F	3	2	-	3/6/23/26	0/1/1/1
2	NAG	G	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1
2	NAG	G	3	2	-	4/6/23/26	0/1/1/1
2	NAG	Н	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	Н	2	2	-	5/6/23/26	0/1/1/1
2	NAG	Н	3	2	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All	(24)	bond	angle	outliers	are	listed	below:
-----	------	------	-------	----------	-----	--------	--------

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Η	3	NAG	C1-O5-C5	5.90	120.19	112.19
2	F	3	NAG	C1-O5-C5	5.17	119.20	112.19
2	G	3	NAG	C1-O5-C5	4.73	118.60	112.19
2	G	2	NAG	C2-N2-C7	3.05	127.25	122.90
2	Ε	2	NAG	C2-N2-C7	3.01	127.18	122.90
2	G	3	NAG	O5-C5-C6	2.94	111.82	107.20
2	Н	3	NAG	O5-C5-C6	2.94	111.81	107.20
2	F	2	NAG	C2-N2-C7	2.82	126.91	122.90
2	F	3	NAG	O5-C5-C6	2.81	111.61	107.20
2	Ε	3	NAG	C2-N2-C7	2.70	126.74	122.90
2	Н	1	NAG	C1-O5-C5	2.64	115.78	112.19
2	Н	2	NAG	C2-N2-C7	2.63	126.65	122.90
2	G	1	NAG	C1-O5-C5	2.59	115.71	112.19
2	Н	3	NAG	C2-N2-C7	2.57	126.56	122.90



Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
2	Ε	2	NAG	C1-O5-C5	2.45	115.51	112.19
2	F	3	NAG	C2-N2-C7	2.43	126.37	122.90
2	F	1	NAG	C1-O5-C5	2.43	115.48	112.19
2	Е	1	NAG	C1-O5-C5	2.41	115.46	112.19
2	Ε	2	NAG	C4-C3-C2	2.37	114.49	111.02
2	Ε	2	NAG	O5-C1-C2	-2.35	107.58	111.29
2	Ε	2	NAG	O4-C4-C3	-2.26	105.12	110.35
2	Ε	2	NAG	C3-C4-C5	2.20	114.16	110.24
2	H	2	NAG	O5-C1-C2	-2.17	107.86	111.29
2	F	2	NAG	O5-C1-C2	-2.00	108.12	111.29

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Е	3	NAG	C3-C2-N2-C7
2	Е	3	NAG	C8-C7-N2-C2
2	Е	3	NAG	O7-C7-N2-C2
2	F	3	NAG	O7-C7-N2-C2
2	G	3	NAG	C8-C7-N2-C2
2	G	3	NAG	O7-C7-N2-C2
2	Н	3	NAG	C3-C2-N2-C7
2	F	3	NAG	C8-C7-N2-C2
2	Н	3	NAG	C8-C7-N2-C2
2	Е	3	NAG	O5-C5-C6-O6
2	Н	3	NAG	O7-C7-N2-C2
2	Н	2	NAG	C8-C7-N2-C2
2	F	3	NAG	C1-C2-N2-C7
2	Н	2	NAG	O7-C7-N2-C2
2	Е	3	NAG	C4-C5-C6-O6
2	Н	2	NAG	C1-C2-N2-C7
2	F	2	NAG	C8-C7-N2-C2
2	Н	2	NAG	C4-C5-C6-O6
2	Н	3	NAG	O5-C5-C6-O6
2	G	3	NAG	C3-C2-N2-C7
2	G	3	NAG	O5-C5-C6-O6
2	F	2	NAG	O7-C7-N2-C2
2	Е	2	NAG	C3-C2-N2-C7
2	Н	2	NAG	O5-C5-C6-O6
2	F	2	NAG	C1-C2-N2-C7
2	F	2	NAG	C3-C2-N2-C7
2	G	2	NAG	C3-C2-N2-C7



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	G	2	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.















## 5.6 Ligand geometry (i)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type	Chain	Dec	Link	Bo	ond leng	$_{\rm ths}$	Bond angles		
	туре	Chain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	С	681	1	14,14,15	0.59	0	17,19,21	1.34	3 (17%)
5	BOG	В	6	-	20,20,20	0.48	0	25,25,25	0.60	0
4	NAG	В	681	1	14,14,15	0.54	0	17,19,21	0.79	0
6	T1N	В	1	-	18,18,18	1.03	0	25,25,25	0.83	1 (4%)
4	NAG	В	661	1	14,14,15	0.50	0	17,19,21	1.59	3 (17%)



Mal	Turne	Chain	Dec	Tink	Bo	ond leng	$_{\rm ths}$	Bond angle		les
	туре	Unain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	T1N	C	2	-	18,18,18	1.05	0	25,25,25	0.86	1 (4%)
6	T1N	D	4	-	18,18,18	1.04	0	25,25,25	0.88	0
3	HEM	D	619	1	41,50,50	1.96	7 (17%)	45,82,82	1.70	5 (11%)
3	HEM	А	619	1	41,50,50	1.98	8 (19%)	45,82,82	1.70	5 (11%)
4	NAG	С	661	1	14,14,15	0.46	0	17,19,21	1.19	2 (11%)
5	BOG	С	5	-	20,20,20	0.48	0	25,25,25	0.57	0
4	NAG	А	661	1	14,14,15	0.47	0	17,19,21	1.14	1 (5%)
4	NAG	D	661	1	14,14,15	0.48	0	17,19,21	1.05	1 (5%)
6	T1N	А	3	-	18,18,18	1.04	0	25,25,25	0.93	0
5	BOG	D	8	-	20,20,20	0.49	0	$25,\!25,\!25$	0.57	0
5	BOG	А	7	-	20,20,20	0.51	0	$25,\!25,\!25$	0.52	0
3	HEM	В	619	1,7	41,50,50	1.99	7 (17%)	45,82,82	1.71	6 (13%)
5	BOG	D	703	-	20,20,20	0.50	0	$25,\!25,\!25$	0.58	0
3	HEM	C	619	1	41,50,50	1.99	8 (19%)	45,82,82	1.76	7 (15%)
5	BOG	А	703	-	20,20,20	0.49	0	25,25,25	0.55	0
4	NAG	D	681	1	$14,\!14,\!15$	0.53	0	17,19,21	0.77	0
4	NAG	A	681	1	14,14,15	0.50	0	17,19,21	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	С	681	1	-	2/6/23/26	0/1/1/1
5	BOG	В	6	-	-	4/11/31/31	0/1/1/1
4	NAG	В	681	1	-	2/6/23/26	0/1/1/1
6	T1N	В	1	-	-	0/10/10/10	0/2/2/2
4	NAG	В	661	1	1/1/5/7	4/6/23/26	0/1/1/1
6	T1N	С	2	-	-	0/10/10/10	0/2/2/2
6	T1N	D	4	-	-	0/10/10/10	0/2/2/2
3	HEM	А	619	1	-	5/12/54/54	-
3	HEM	D	619	1	-	5/12/54/54	-
4	NAG	С	661	1	1/1/5/7	2/6/23/26	0/1/1/1
5	BOG	С	5	-	-	3/11/31/31	0/1/1/1
4	NAG	А	661	1	1/1/5/7	3/6/23/26	0/1/1/1
4	NAG	D	661	1	1/1/5/7	4/6/23/26	0/1/1/1
6	T1N	А	3	-	-	0/10/10/10	0/2/2/2



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BOG	D	8	-	-	5/11/31/31	0/1/1/1
5	BOG	А	7	-	-	4/11/31/31	0/1/1/1
3	HEM	В	619	1,7	-	4/12/54/54	-
5	BOG	D	703	-	-	5/11/31/31	0/1/1/1
3	HEM	С	619	1	-	5/12/54/54	-
5	BOG	А	703	-	-	8/11/31/31	0/1/1/1
4	NAG	D	681	1	-	2/6/23/26	0/1/1/1
4	NAG	А	681	1	-	2/6/23/26	0/1/1/1

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	В	619	HEM	C3D-C2D	8.08	1.53	1.36
3	А	619	HEM	C3D-C2D	8.07	1.53	1.36
3	D	619	HEM	C3D-C2D	8.04	1.53	1.36
3	С	619	HEM	C3D-C2D	8.02	1.53	1.36
3	С	619	HEM	C3C-C2C	-4.37	1.34	1.40
3	D	619	HEM	C3C-C2C	-4.33	1.34	1.40
3	А	619	HEM	C3C-C2C	-4.32	1.34	1.40
3	В	619	HEM	C3C-C2C	-4.28	1.34	1.40
3	С	619	HEM	C3C-CAC	3.66	1.55	1.47
3	В	619	HEM	C3C-CAC	3.65	1.55	1.47
3	А	619	HEM	C3C-CAC	3.58	1.55	1.47
3	D	619	HEM	C3C-CAC	3.54	1.55	1.47
3	В	619	HEM	CAB-C3B	2.88	1.55	1.47
3	С	619	HEM	CAB-C3B	2.80	1.55	1.47
3	А	619	HEM	CAB-C3B	2.78	1.55	1.47
3	D	619	HEM	CAB-C3B	2.74	1.54	1.47
3	В	619	HEM	FE-ND	2.70	2.10	1.96
3	С	619	HEM	CAA-C2A	2.70	1.56	1.52
3	А	619	HEM	FE-ND	2.61	2.09	1.96
3	D	619	HEM	FE-ND	2.48	2.09	1.96
3	С	619	HEM	FE-ND	2.44	2.08	1.96
3	В	619	HEM	CAA-C2A	2.42	1.55	1.52
3	С	619	HEM	FE-NB	2.26	2.08	1.96
3	В	619	HEM	FE-NB	2.15	2.07	1.96
3	А	619	HEM	FE-NB	2.14	2.07	1.96
3	А	619	HEM	CAA-C2A	2.14	1.55	1.52
3	С	619	HEM	CMB-C2B	2.03	1.55	1.50
3	D	619	HEM	CAA-C2A	2.02	1.55	1.52



$\alpha$ $\cdot$ $\cdot$ $\cdot$	C	•	
Continued	trom	previous	page
00100000000	J. 00	p. 0000 a0	$p \approx g \circ \cdots$

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	619	HEM	CMD-C2D	2.00	1.55	1.50
3	А	619	HEM	CMB-C2B	2.00	1.55	1.50

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	А	619	HEM	C4D-ND-C1D	6.56	111.85	105.07
3	С	619	HEM	C4D-ND-C1D	6.52	111.80	105.07
3	D	619	HEM	C4D-ND-C1D	6.38	111.67	105.07
3	В	619	HEM	C4D-ND-C1D	6.30	111.58	105.07
4	В	661	NAG	C1-O5-C5	4.46	118.23	112.19
4	С	661	NAG	C1-O5-C5	3.89	117.46	112.19
4	А	661	NAG	C1-O5-C5	3.89	117.46	112.19
3	А	619	HEM	C4B-CHC-C1C	3.43	127.08	122.56
3	С	619	HEM	C4B-CHC-C1C	3.38	127.02	122.56
3	D	619	HEM	C4B-CHC-C1C	3.33	126.96	122.56
3	С	619	HEM	C1B-NB-C4B	3.22	108.40	105.07
4	D	661	NAG	C1-O5-C5	3.21	116.54	112.19
4	В	661	NAG	C2-N2-C7	3.19	127.44	122.90
3	В	619	HEM	C1B-NB-C4B	3.17	108.34	105.07
3	В	619	HEM	C4C-CHD-C1D	3.16	126.72	122.56
4	С	681	NAG	C4-C3-C2	3.14	115.62	111.02
3	D	619	HEM	C4C-CHD-C1D	3.14	126.70	122.56
3	В	619	HEM	C4B-CHC-C1C	3.06	126.60	122.56
3	D	619	HEM	C1B-NB-C4B	3.06	108.23	105.07
3	А	619	HEM	C1B-NB-C4B	3.01	108.18	105.07
3	В	619	HEM	CBA-CAA-C2A	-2.95	107.58	112.62
3	С	619	HEM	C4C-CHD-C1D	2.89	126.38	122.56
4	С	681	NAG	C3-C4-C5	2.83	115.28	110.24
3	А	619	HEM	C4C-CHD-C1D	2.76	126.21	122.56
3	В	619	HEM	C3B-C2B-C1B	2.44	108.30	106.49
3	С	619	HEM	C3B-C2B-C1B	2.35	108.23	106.49
4	В	661	NAG	O5-C5-C6	2.19	110.63	107.20
3	D	619	HEM	C3B-C2B-C1B	2.13	108.07	106.49
4	С	681	NAG	O5-C5-C6	2.12	110.52	107.20
6	В	1	T1N	CAA-SAK-CAM	2.11	108.46	103.38
3	А	619	HEM	C3B-C2B-C1B	2.11	108.05	106.49
6	С	2	T1N	CAA-SAK-CAM	2.09	108.42	103.38
3	С	619	HEM	CMA-C3A-C4A	-2.06	125.30	128.46
4	С	661	NAG	O5-C5-C6	2.06	110.43	107.20
3	С	619	HEM	CBA-CAA-C2A	-2.01	109.19	112.62



All (4) chirality outliers are listed below	<i>'</i> :
---	------------

Mol	Chain	$\mathbf{Res}$	Type	Atom
4	А	661	NAG	C1
4	В	661	NAG	C1
4	С	661	NAG	C1
4	D	661	NAG	C1

All (69) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	661	NAG	C8-C7-N2-C2
4	D	661	NAG	O7-C7-N2-C2
4	А	661	NAG	C8-C7-N2-C2
4	А	661	NAG	O7-C7-N2-C2
4	С	681	NAG	C4-C5-C6-O6
4	С	681	NAG	O5-C5-C6-O6
4	D	661	NAG	C4-C5-C6-O6
5	D	703	BOG	C3'-C4'-C5'-C6'
5	А	7	BOG	C4'-C5'-C6'-C7'
5	В	6	BOG	C3'-C4'-C5'-C6'
4	В	661	NAG	C4-C5-C6-O6
5	А	703	BOG	C4'-C5'-C6'-C7'
5	В	6	BOG	C1'-C2'-C3'-C4'
5	D	8	BOG	C1'-C2'-C3'-C4'
5	А	703	BOG	C2'-C3'-C4'-C5'
5	D	703	BOG	O1-C1'-C2'-C3'
5	А	7	BOG	C1'-C2'-C3'-C4'
5	А	7	BOG	C3'-C4'-C5'-C6'
5	D	8	BOG	C4'-C5'-C6'-C7'
3	А	619	HEM	C2B-C3B-CAB-CBB
3	В	619	HEM	C2B-C3B-CAB-CBB
3	С	619	HEM	C2B-C3B-CAB-CBB
3	D	619	HEM	C2B-C3B-CAB-CBB
5	А	703	BOG	O5-C1-O1-C1'
3	А	619	HEM	C4B-C3B-CAB-CBB
3	В	619	HEM	C4B-C3B-CAB-CBB
3	С	619	HEM	C4B-C3B-CAB-CBB
3	D	619	HEM	C4B-C3B-CAB-CBB
5	D	703	BOG	C4'-C5'-C6'-C7'
5	D	8	BOG	C3'-C4'-C5'-C6'
5	А	703	BOG	C2-C1-O1-C1'
5	В	6	BOG	C4'-C5'-C6'-C7'
4	А	681	NAG	C4-C5-C6-O6
3	D	619	HEM	C2A-CAA-CBA-CGA



Mol	Chain	Res	Type	Atoms
5	D	8	BOG	C5'-C6'-C7'-C8'
4	В	661	NAG	C8-C7-N2-C2
5	С	5	BOG	C5'-C6'-C7'-C8'
5	В	6	BOG	C5'-C6'-C7'-C8'
5	D	8	BOG	C2'-C3'-C4'-C5'
5	D	703	BOG	C5'-C6'-C7'-C8'
5	А	703	BOG	C3'-C4'-C5'-C6'
4	В	661	NAG	C3-C2-N2-C7
5	D	703	BOG	C1'-C2'-C3'-C4'
5	А	703	BOG	C2'-C1'-O1-C1
4	D	681	NAG	C4-C5-C6-O6
4	А	661	NAG	C4-C5-C6-O6
5	А	703	BOG	C1'-C2'-C3'-C4'
4	В	661	NAG	O7-C7-N2-C2
4	D	661	NAG	O5-C5-C6-O6
5	С	5	BOG	C1'-C2'-C3'-C4'
5	А	703	BOG	O1-C1'-C2'-C3'
4	С	661	NAG	C8-C7-N2-C2
3	А	619	HEM	C2A-CAA-CBA-CGA
4	В	681	NAG	C4-C5-C6-O6
4	А	681	NAG	O5-C5-C6-O6
3	С	619	HEM	C2A-CAA-CBA-CGA
3	В	619	HEM	CAA-CBA-CGA-O1A
4	В	681	NAG	O5-C5-C6-O6
4	С	661	NAG	O7-C7-N2-C2
4	D	681	NAG	O5-C5-C6-O6
5	А	7	BOG	C4-C5-C6-O6
3	В	619	HEM	CAA-CBA-CGA-O2A
3	С	619	HEM	CAA-CBA-CGA-O2A
3	С	619	HEM	CAA-CBA-CGA-O1A
3	А	619	HEM	CAA-CBA-CGA-O2A
3	D	619	HEM	CAA-CBA-CGA-O2A
3	D	619	HEM	CAA-CBA-CGA-O1A
3	А	619	HEM	CAA-CBA-CGA-O1A
5	С	5	BOG	C2'-C3'-C4'-C5'

Continued from previous page...

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	С	619	HEM	2	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and similar rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

































# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	2	$\mathbf{OWAB}(\mathbf{A}^2)$	Q<0.9
1	А	552/587~(94%)	0.76	55~(9%) 7	9	35, 49, 69, 84	0
1	В	551/587~(93%)	0.82	64 (11%) 4	6	36, 52, 75, 89	0
1	С	552/587~(94%)	0.67	49 (8%) 9 1	12	34, 47, 63, 77	0
1	D	552/587~(94%)	0.67	44 (7%) 12	15	35, 47, 62, 75	0
All	All	2207/2348~(93%)	0.73	212 (9%) 8	10	34, 48, 68, 89	0

All (212) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	409	TYR	9.9
1	С	583	GLN	7.2
1	С	80	LEU	6.3
1	D	122	TYR	5.3
1	В	33	ALA	5.2
1	В	122	TYR	5.1
1	В	74	PHE	4.8
1	А	278	HIS	4.4
1	С	486	GLU	4.4
1	А	494	LEU	4.2
1	В	481	LEU	4.2
1	А	101	ASN	4.1
1	А	122	TYR	4.1
1	А	81	LEU	4.0
1	D	296	PRO	4.0
1	D	287	VAL	4.0
1	С	122	TYR	3.9
1	В	53	ASP	3.9
1	А	177	VAL	3.8
1	С	422	PHE	3.8
1	В	81	LEU	3.8



Mol

1

1

1

1

1

1

1

1

1

1

1

1

1

1 1

1 1

_			
	399	ASP	3.6
	82	LEU	3.6
	105	ASN	3.6
	399	ASP	3.5
	105	ASN	3.5
	77	ARG	3.5
	158	ASP	3.5
	215	LYS	3.5
	473	LYS	3.5
	158	ASP	3.5
	82	LEU	3.4
	364	GLU	3.4
	53	ASP	3.4
	91	TYR	3.4
	278	HIS	3.3
	399	ASP	3.3
	278	HIS	3.3
	422	PHE	3.3

Continued from previous page... Chain

D

D

В

А

С

В

D

В

D

В

А

А D

В

В В

D

 $\mathbf{Res}$ 

81

422

105(A)

161

181

422

Type

LEU

PHE

ILE

THR

VAL

PHE

RSRZ

3.8

3.7

3.7

3.7

3.7

3.6

1	В	364	GLU	3.4
1	D	53	ASP	3.4
1	А	91	TYR	3.4
1	С	278	HIS	3.3
1	С	399	ASP	3.3
1	D	278	HIS	3.3
1	А	422	PHE	3.3
1	С	493	ALA	3.3
1	В	484	GLU	3.3
1	В	54	GLN	3.3
1	А	80	LEU	3.3
1	С	91	TYR	3.3
1	D	77	ARG	3.3
1	А	583	GLN	3.2
1	А	457	GLU	3.2
1	А	158	ASP	3.2
1	С	75	LEU	3.2
1	С	577	PHE	3.2
1	С	312	VAL	3.2
1	D	101	ASN	3.2
1	В	214	HIS	3.2
1	D	280	PRO	3.2
1	A	239	ASP	3.2
1	В	483	GLY	3.2
1	В	148	TYR	3.1



Continued from previous page...

10101	Unam	nes	Tybe	nsnz
1	A	401	GLU	3.1
1	В	577	PHE	3.1
1	В	287	VAL	3.1
1	В	296	PRO	3.1
1	С	160	PRO	3.1
1	D	318	GLN	3.1
1	В	316	LEU	3.0
1	D	481	LEU	3.0
1	В	278	HIS	3.0
1	В	377	ILE	3.0
1	D	286	ALA	3.0
1	В	105	ASN	3.0
1	С	272	GLU	2.9
1	А	472	LEU	2.9
1	А	33	ALA	2.9
1	В	286	ALA	2.9
1	А	74	PHE	2.9
1	С	148	TYR	2.9
1	В	99	VAL	2.9
1	А	409	TYR	2.9
1	А	82	LEU	2.9
1	С	95	HIS	2.9
1	А	159	CYS	2.8
1	С	582	VAL	2.8
1	В	272	GLU	2.8
1	D	186	GLU	2.8
1	В	160	PRO	2.8
1	С	494	LEU	2.8
1	А	368	ASN	2.8
1	А	170	GLU	2.8
1	А	450	ALA	2.7
1	С	453	ASP	2.7
1	D	252	LEU	2.7
1	В	581	ASN	2.7
1	А	83	LYS	2.7
1	В	416	GLU	2.7
1	С	401	GLU	2.7
1	С	409	TYR	2.7
1	D	234	TYR	2.7
1	В	186	GLU	2.7
1	В	98	GLY	2.6
1	D	583	GLN	2.6



Continued from previous page					
Mol	Chain	Res	Type	RSRZ	
1	С	286	ALA	2.6	
1	С	214	HIS	2.6	
1	С	81	LEU	2.6	
1	В	575	CYS	2.6	
1	А	168	ASN	2.6	
1	А	186	GLU	2.6	
1	С	575	CYS	2.6	
1	С	33	ALA	2.6	
1	D	377	ILE	2.6	
1	А	390	LEU	2.6	
1	А	399	ASP	2.6	
1	С	215	LYS	2.6	
1	А	296	PRO	2.5	
1	А	473	LYS	2.5	
1	А	482	THR	2.5	
1	С	82	LEU	2.5	
1	С	113	MET	2.5	
1	А	486	GLU	2.5	
1	С	239	ASP	2.5	
1	А	491	LEU	2.5	
1	D	59	CYS	2.5	
1	А	272	GLU	2.5	
1	А	222	ARG	2.5	
1	В	352	LEU	2.5	
1	А	441	PRO	2.4	
1	С	59	CYS	2.4	
1	В	493	ALA	2.4	
1	С	428	ARG	2.4	
1	А	116	VAL	2.4	
1	А	515	ASP	2.4	
1	С	77	ARG	2.4	
1	В	491	LEU	2.4	
1	D	54	GLN	2.4	
1	С	295	VAL	2.4	
1	D	74	PHE	2.4	
1	В	290	GLU	2.4	
1	D	515	ASP	2.4	
1	В	198	PHE	2.4	
1	С	36	CYS	2.4	
1	D	500	VAL	2.4	
1	D	506	ALA	2.3	
1	D	473	LYS	2.3	

473LYS2.3Continued on next page...



Continued from previous page					
Mol	Chain	Res	Type	RSRZ	
1	В	101	ASN	2.3	
1	В	426	PHE	2.3	
1	С	53	ASP	2.3	
1	С	158	ASP	2.3	
1	D	91	TYR	2.3	
1	D	328	LEU	2.3	
1	D	268	ASP	2.3	
1	D	379	SER	2.3	
1	В	428	ARG	2.3	
1	С	234	TYR	2.3	
1	С	186	GLU	2.3	
1	В	341	ILE	2.3	
1	А	373	TYR	2.3	
1	В	75	LEU	2.3	
1	В	252	LEU	2.3	
1	С	491	LEU	2.3	
1	С	170	GLU	2.3	
1	А	53	ASP	2.3	
1	В	102	ILE	2.3	
1	А	508	LEU	2.2	
1	В	80	LEU	2.2	
1	D	316	LEU	2.2	
1	D	400	GLN	2.2	
1	В	485	LYS	2.2	
1	D	409	TYR	2.2	
1	В	52	PHE	2.2	
1	В	107	PHE	2.2	
1	D	113	MET	2.2	
1	А	148	TYR	2.2	
1	В	444	VAL	2.2	
1	D	165	VAL	2.2	
1	В	73	GLU	2.2	
1	С	481	LEU	2.2	
1	В	77	ARG	2.2	
1	В	51	GLY	2.2	
1	В	506	ALA	2.2	
1	В	363	PRO	2.2	
1	А	79	LYS	2.2	
1	С	303	THR	2.2	
1	A	93	LEU	2.2	
1	А	562	ALA	2.1	
1	В	221	THR	2.1	

221THR2.1Continued on next page...



Mol	Chain	Res	Type	RSRZ
1	В	159	CYS	2.1
1	D	41	CYS	2.1
1	А	157	ASP	2.1
1	В	234	TYR	2.1
1	С	515	ASP	2.1
1	А	113	MET	2.1
1	А	355	TYR	2.1
1	С	416	GLU	2.1
1	D	352	LEU	2.1
1	С	159	CYS	2.1
1	С	258	GLY	2.1
1	А	98	GLY	2.1
1	В	106	PRO	2.1
1	D	329	PHE	2.1
1	D	335	ILE	2.0
1	А	56	LYS	2.0
1	D	258	GLY	2.0
1	С	395	PHE	2.0
1	D	239	ASP	2.0
1	A	535	MET	2.0
1	A	85	THR	2.0
1	В	515	ASP	2.0
1	В	294	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	NAG	Е	3	14/15	0.48	0.56	$63,\!64,\!65,\!65$	0
2	NAG	G	3	14/15	0.50	0.58	62,63,64,64	0
2	NAG	F	3	14/15	0.57	0.69	$63,\!65,\!66,\!66$	0
2	NAG	G	2	14/15	0.59	0.38	53,56,57,60	0
2	NAG	Н	3	14/15	0.59	0.51	62,64,64,64	0





Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	NAG	Е	2	14/15	0.68	0.26	$53,\!55,\!58,\!60$	0
2	NAG	F	2	14/15	0.77	0.39	54,58,59,61	0
2	NAG	Н	2	14/15	0.79	0.32	53,56,57,60	0
2	NAG	Е	1	14/15	0.86	0.15	40,43,46,50	0
2	NAG	G	1	14/15	0.86	0.18	39,42,45,49	0
2	NAG	F	1	14/15	0.90	0.12	40,42,46,50	0
2	NAG	Н	1	14/15	0.92	0.12	38,41,44,49	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.











# 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
5	BOG	А	7	20/20	0.59	0.35	$64,\!65,\!66,\!66$	0
5	BOG	D	8	20/20	0.63	0.28	$63,\!63,\!63,\!63$	0
5	BOG	А	703	20/20	0.69	0.27	58,58,60,60	0
4	NAG	С	681	14/15	0.70	0.45	$48,\!51,\!53,\!53$	0
4	NAG	В	661	14/15	0.70	0.39	48,51,53,53	0
5	BOG	В	6	20/20	0.76	0.26	72,72,72,72	0
4	NAG	D	661	14/15	0.77	0.43	47,50,52,53	0
4	NAG	А	661	14/15	0.78	0.43	47,50,51,51	0
4	NAG	D	681	14/15	0.78	0.25	46,49,50,50	0
4	NAG	С	661	14/15	0.81	0.38	47,50,51,51	0
4	NAG	А	681	14/15	0.82	0.32	44,47,48,48	0
4	NAG	В	681	14/15	0.82	0.37	48,50,51,51	0



Mol	Type	Chain	Res	Atoms	RSCC	$\mathbf{RSR}$	B-factors(Å <sup>2</sup> )	Q<0.9
5	BOG	D	703	20/20	0.84	0.19	$51,\!51,\!55,\!55$	0
5	BOG	С	5	20/20	0.85	0.18	60,60,62,62	0
3	HEM	А	619	43/43	0.88	0.19	44,45,49,50	0
3	HEM	С	619	43/43	0.88	0.18	41,43,46,48	0
6	T1N	D	4	17/17	0.89	0.16	40,41,41,42	0
6	T1N	С	2	17/17	0.90	0.15	41,41,42,43	0
3	HEM	D	619	43/43	0.90	0.17	42,44,48,50	0
3	HEM	В	619	43/43	0.91	0.14	46,47,49,50	0
6	T1N	А	3	17/17	0.92	0.14	40,41,42,42	0
6	T1N	В	1	17/17	0.92	0.15	43,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.































# 6.5 Other polymers (i)

There are no such residues in this entry.

