



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 06:14 am BST

PDB ID : 5NTW
Title : Structural states of RORgt: X-ray elucidation of molecular mechanisms and binding interactions for natural and synthetic compounds
Authors : Kallen, J.
Deposited on : 2017-04-28
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	
1	B	257	
1	C	257	
1	D	257	
2	P	20	
2	Q	20	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	R	20	 35% 10% 55%
2	S	20	 10% 40% 5% 55%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9687 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear receptor ROR-gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	Total 2046	C 1310	N 359	O 363	S 14	0	10	0
1	B	245	Total 2035	C 1305	N 358	O 359	S 13	0	7	0
1	C	245	Total 2073	C 1330	N 365	O 365	S 13	0	13	0
1	D	245	Total 2066	C 1322	N 364	O 366	S 14	0	12	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	262	GLY	-	expression tag	UNP P51449
A	455	SER	CYS	engineered mutation	UNP P51449
B	262	GLY	-	expression tag	UNP P51449
B	455	SER	CYS	engineered mutation	UNP P51449
C	262	GLY	-	expression tag	UNP P51449
C	455	SER	CYS	engineered mutation	UNP P51449
D	262	GLY	-	expression tag	UNP P51449
D	455	SER	CYS	engineered mutation	UNP P51449

- Molecule 2 is a protein called Nuclear receptor-interacting protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	P	9	Total 70	C 47	N 12	O 11	0	0	0
2	Q	10	Total 79	C 53	N 14	O 12	0	0	0
2	R	9	Total 70	C 47	N 12	O 11	0	0	0
2	S	9	Total 70	C 47	N 12	O 11	0	0	0

- Molecule 3 is (S)-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-7-isopropyl-6-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-6,7-dihydro-5H-pyrrolo[3,4-b]pyridine-3-carboxamide (three-letter code: 98N) (formula: C₂₇H₃₅F₃N₄O₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
3	A	1	Total	C	F	N	O	S	0	0
			38	27	3	4	3	1		
3	B	1	Total	C	F	N	O	S	0	0
			38	27	3	4	3	1		
3	C	1	Total	C	F	N	O	S	0	0
			38	27	3	4	3	1		
3	D	1	Total	C	F	N	O	S	0	0
			38	27	3	4	3	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	229	Total	O	0	0
			229	229		
4	B	235	Total	O	0	0
			235	235		
4	C	270	Total	O	0	0
			270	270		
4	D	268	Total	O	0	0
			268	268		
4	Q	11	Total	O	0	0
			11	11		
4	R	8	Total	O	0	0
			8	8		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	S	5	Total	O	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nuclear receptor ROR-gamma




- Molecule 1: Nuclear receptor ROR-gamma




- Molecule 1: Nuclear receptor ROR-gamma




- Molecule 1: Nuclear receptor ROR-gamma





- Molecule 2: Nuclear receptor-interacting protein 1





- Molecule 2: Nuclear receptor-interacting protein 1



- Molecule 2: Nuclear receptor-interacting protein 1



- Molecule 2: Nuclear receptor-interacting protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	86.19Å 68.47Å 96.72Å 90.00° 110.32° 90.00°	Depositor
Resolution (Å)	19.72 – 1.64 19.54 – 1.64	Depositor EDS
% Data completeness (in resolution range)	98.1 (19.72-1.64) 98.2 (19.54-1.64)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 1.64Å)	Xtrriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.200 , 0.222 0.199 , 0.221	Depositor DCC
R_{free} test set	4950 reflections (3.90%)	wwPDB-VP
Wilson B-factor (Å ²)	17.1	Xtrriage
Anisotropy	0.142	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9687	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 59.95 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6352e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 98N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/2117	0.46	0/2853
1	B	0.38	0/2100	0.48	0/2829
1	C	0.38	0/2157	0.48	0/2905
1	D	0.37	0/2141	0.49	0/2884
2	P	0.31	0/70	0.45	0/94
2	Q	0.35	0/79	0.52	0/105
2	R	0.33	0/70	0.49	0/94
2	S	0.36	0/70	0.47	0/94
All	All	0.37	0/8804	0.48	0/11858

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2046	0	2074	6	0
1	B	2035	0	2067	11	0
1	C	2073	0	2113	9	0
1	D	2066	0	2091	7	0
2	P	70	0	79	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Q	79	0	92	0	0
2	R	70	0	79	2	0
2	S	70	0	79	1	0
3	A	38	0	0	0	0
3	B	38	0	0	2	0
3	C	38	0	0	2	0
3	D	38	0	0	0	0
4	A	229	0	0	4	0
4	B	235	0	0	1	0
4	C	270	0	0	2	0
4	D	268	0	0	3	0
4	Q	11	0	0	0	0
4	R	8	0	0	0	0
4	S	5	0	0	1	0
All	All	9687	0	8674	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:479:HIS:CD2	3:B:601:98N:F38	2.20	0.85
1:D:494:VAL:O	1:D:503[A]:LYS:HE3	1.80	0.81
1:C:322[B]:HIS:CD2	4:C:729:HOH:O	2.43	0.71
1:A:322:HIS:CD2	1:C:322[A]:HIS:CD2	2.80	0.70
1:C:437:ARG:HD2	1:C:440:GLU:OE1	1.95	0.67
1:B:479:HIS:HD2	3:B:601:98N:F38	1.69	0.62
1:C:479:HIS:HD2	3:C:601:98N:F37	1.81	0.54
1:C:479:HIS:CD2	3:C:601:98N:F37	2.52	0.53
1:C:492:ILE:HD12	2:P:503:LEU:HB3	1.92	0.51
1:B:324[A]:LEU:HD21	1:B:358:MET:SD	2.51	0.50
1:B:495:GLN:HB2	4:B:789:HOH:O	2.13	0.48
1:D:494:VAL:O	1:D:503[A]:LYS:CE	2.59	0.48
1:B:482:ARG:HA	1:B:485:ILE:HD12	1.96	0.46
1:C:350[B]:ILE:HD11	2:R:502:GLN:HG2	1.97	0.45
1:B:324[A]:LEU:HD11	1:B:358:MET:SD	2.57	0.45
1:D:358:MET:HB3	4:D:702:HOH:O	2.16	0.45
1:B:487:GLN:HE22	1:B:494:VAL:HB	1.81	0.45
1:D:412:PHE:HB3	1:D:417:ILE:HG13	1.99	0.44
1:A:324[A]:LEU:HD23	1:A:502:TYR:HE1	1.82	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:282:ARG:NH2	4:D:704:HOH:O	2.49	0.44
2:S:499:THR:N	4:S:601:HOH:O	2.51	0.44
1:A:412:PHE:HB3	1:A:417:ILE:HG13	2.00	0.43
1:A:451:HIS:HD2	4:A:913:HOH:O	2.01	0.43
1:D:451:HIS:HE1	4:D:926:HOH:O	2.00	0.43
1:C:298[A]:ASN:ND2	4:C:704:HOH:O	2.51	0.43
1:B:412:PHE:HB3	1:B:417:ILE:HG13	2.01	0.43
2:R:499:THR:N	2:R:502:GLN:HE21	2.19	0.41
1:B:392:GLY:O	1:B:482:ARG:NH2	2.53	0.41
1:D:324[A]:LEU:HD21	1:D:358:MET:SD	2.61	0.41
4:A:807:HOH:O	1:B:485:ILE:HG12	2.19	0.41
1:A:499:PRO:HD3	4:A:790:HOH:O	2.21	0.41
1:A:395:GLU:HG2	4:A:923:HOH:O	2.20	0.40
1:B:478:GLN:O	1:B:482:ARG:HG2	2.22	0.40
1:C:270[A]:ILE:HD13	1:C:270[A]:ILE:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/257 (98%)	250 (99%)	3 (1%)	0	100	100
1	B	250/257 (97%)	246 (98%)	4 (2%)	0	100	100
1	C	256/257 (100%)	252 (98%)	4 (2%)	0	100	100
1	D	255/257 (99%)	251 (98%)	4 (2%)	0	100	100
2	P	7/20 (35%)	7 (100%)	0	0	100	100
2	Q	8/20 (40%)	8 (100%)	0	0	100	100
2	R	7/20 (35%)	7 (100%)	0	0	100	100
2	S	7/20 (35%)	7 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1043/1108 (94%)	1028 (99%)	15 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/229 (100%)	229 (100%)	0	100	100
1	B	226/229 (99%)	226 (100%)	0	100	100
1	C	232/229 (101%)	232 (100%)	0	100	100
1	D	231/229 (101%)	231 (100%)	0	100	100
2	P	8/19 (42%)	8 (100%)	0	100	100
2	Q	9/19 (47%)	9 (100%)	0	100	100
2	R	8/19 (42%)	8 (100%)	0	100	100
2	S	8/19 (42%)	8 (100%)	0	100	100
All	All	951/992 (96%)	951 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	329	GLN
1	A	451	HIS
1	B	429	HIS
1	B	460	GLN
1	B	487	GLN
1	B	488	HIS
1	D	322	HIS
1	D	451	HIS
2	Q	502	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	S	507	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	98N	D	601	-	38,41,41	1.12	3 (7%)	52,61,61	1.69	12 (23%)
3	98N	B	601	-	38,41,41	1.11	2 (5%)	52,61,61	1.65	16 (30%)
3	98N	C	601	-	38,41,41	1.03	2 (5%)	52,61,61	1.75	16 (30%)
3	98N	A	601	-	38,41,41	1.15	2 (5%)	52,61,61	1.67	13 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	98N	D	601	-	-	6/32/54/54	0/4/4/4
3	98N	B	601	-	-	0/32/54/54	0/4/4/4
3	98N	C	601	-	-	9/32/54/54	0/4/4/4
3	98N	A	601	-	-	9/32/54/54	0/4/4/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	98N	C24-S23	3.24	1.81	1.77
3	D	601	98N	C24-S23	2.70	1.81	1.77
3	A	601	98N	C35-C32	2.67	1.54	1.50
3	B	601	98N	C24-S23	2.64	1.80	1.77
3	D	601	98N	C35-C32	2.58	1.54	1.50
3	B	601	98N	C35-C32	2.50	1.54	1.50
3	C	601	98N	C35-C32	2.37	1.54	1.50
3	C	601	98N	C24-S23	2.32	1.80	1.77
3	D	601	98N	C19-C20	2.01	1.40	1.38

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	98N	C1-C2-N3	-4.11	122.03	125.27
3	D	601	98N	C5-C6-C1	-3.75	115.76	121.47
3	A	601	98N	C1-C2-N3	-3.73	122.33	125.27
3	D	601	98N	C1-C2-N3	-3.66	122.39	125.27
3	C	601	98N	O27-S23-C24	3.48	111.37	108.29
3	A	601	98N	C5-C6-C1	-3.46	116.20	121.47
3	C	601	98N	C5-C6-C1	-3.44	116.24	121.47
3	C	601	98N	C8-N9-C10	3.38	112.36	108.11
3	C	601	98N	C6-C5-C4	3.34	121.39	117.95
3	D	601	98N	C8-N9-C10	3.28	112.24	108.11
3	B	601	98N	C5-C6-C1	-3.13	116.70	121.47
3	A	601	98N	C8-N9-C10	3.13	112.05	108.11
3	A	601	98N	O27-S23-C24	3.11	111.04	108.29
3	D	601	98N	C19-N18-C17	3.09	121.78	117.82
3	C	601	98N	F38-C35-C32	-3.01	108.24	112.53
3	D	601	98N	C29-C28-N9	-3.00	108.08	113.39
3	C	601	98N	C1-C2-N3	-2.94	122.95	125.27
3	B	601	98N	F38-C35-C32	-2.91	108.38	112.53
3	D	601	98N	C6-C5-C4	2.80	120.83	117.95
3	C	601	98N	C30-C31-C32	-2.75	106.33	111.84
3	A	601	98N	F37-C35-C32	-2.69	108.69	112.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	98N	C19-N18-C17	2.65	121.22	117.82
3	A	601	98N	C33-C32-C35	-2.64	109.04	111.76
3	A	601	98N	F38-C35-C32	-2.62	108.79	112.53
3	A	601	98N	C6-C5-C4	2.62	120.65	117.95
3	C	601	98N	C17-C16-N15	-2.61	107.05	112.71
3	D	601	98N	C17-C16-N15	-2.59	107.09	112.71
3	B	601	98N	C29-C28-N9	-2.59	108.81	113.39
3	C	601	98N	C29-C28-N9	-2.58	108.84	113.39
3	A	601	98N	C19-N18-C17	2.55	121.09	117.82
3	D	601	98N	F38-C35-C32	-2.54	108.91	112.53
3	C	601	98N	F37-C35-C32	-2.50	108.97	112.53
3	B	601	98N	C17-C16-N15	-2.50	107.30	112.71
3	D	601	98N	O27-S23-C24	2.49	110.49	108.29
3	C	601	98N	F38-C35-F37	2.47	112.39	106.87
3	D	601	98N	F36-C35-C32	-2.46	109.02	112.53
3	C	601	98N	C19-N18-C17	2.42	120.93	117.82
3	D	601	98N	C6-C1-C2	2.39	121.11	118.86
3	B	601	98N	F36-C35-C32	-2.35	109.18	112.53
3	C	601	98N	C12-C11-C10	2.33	113.87	110.72
3	C	601	98N	C5-C4-N3	-2.24	120.54	123.67
3	B	601	98N	C8-N9-C10	2.24	110.92	108.11
3	B	601	98N	C34-C29-C30	2.22	114.76	109.33
3	B	601	98N	C31-C32-C35	-2.21	109.49	111.76
3	A	601	98N	C6-C1-C2	2.21	120.94	118.86
3	A	601	98N	C30-C31-C32	-2.19	107.44	111.84
3	A	601	98N	C17-C16-N15	-2.17	108.01	112.71
3	B	601	98N	F37-C35-F36	2.16	111.70	106.87
3	B	601	98N	C30-C31-C32	-2.14	107.55	111.84
3	B	601	98N	F37-C35-C32	-2.13	109.49	112.53
3	A	601	98N	C34-C29-C30	2.11	114.50	109.33
3	B	601	98N	C6-C1-C2	2.11	120.85	118.86
3	C	601	98N	F36-C35-C32	-2.09	109.55	112.53
3	B	601	98N	C12-C11-C10	2.06	113.49	110.72
3	B	601	98N	C6-C5-C4	2.05	120.06	117.95
3	C	601	98N	C34-C33-C32	-2.05	107.74	111.84
3	D	601	98N	F37-C35-C32	-2.04	109.62	112.53

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	601	98N	C31-C32-C35-F36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	601	98N	C31-C32-C35-F37
3	D	601	98N	C31-C32-C35-F38
3	D	601	98N	C33-C32-C35-F36
3	D	601	98N	C33-C32-C35-F37
3	D	601	98N	C33-C32-C35-F38
3	C	601	98N	N9-C10-C11-C12
3	C	601	98N	C31-C32-C35-F36
3	C	601	98N	C31-C32-C35-F37
3	C	601	98N	C31-C32-C35-F38
3	C	601	98N	C33-C32-C35-F36
3	C	601	98N	C33-C32-C35-F37
3	C	601	98N	C33-C32-C35-F38
3	A	601	98N	C2-C10-C11-C12
3	A	601	98N	N9-C10-C11-C12
3	A	601	98N	N9-C10-C11-C13
3	A	601	98N	C31-C32-C35-F36
3	A	601	98N	C31-C32-C35-F37
3	A	601	98N	C31-C32-C35-F38
3	A	601	98N	C33-C32-C35-F37
3	C	601	98N	N9-C10-C11-C13
3	A	601	98N	C33-C32-C35-F38
3	A	601	98N	C33-C32-C35-F36
3	C	601	98N	C2-C10-C11-C12


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	601	98N	2	0
3	C	601	98N	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/257 (95%)	0.27	10 (4%) 37 35	10, 18, 29, 35	0
1	B	245/257 (95%)	0.10	6 (2%) 59 58	9, 16, 26, 32	0
1	C	245/257 (95%)	0.03	4 (1%) 72 73	8, 14, 23, 33	0
1	D	245/257 (95%)	0.11	6 (2%) 59 58	9, 16, 25, 31	0
2	P	9/20 (45%)	1.80	2 (22%) 0 0	27, 31, 43, 52	0
2	Q	10/20 (50%)	0.73	1 (10%) 7 5	17, 19, 25, 36	0
2	R	9/20 (45%)	0.26	0 100 100	16, 18, 22, 24	0
2	S	9/20 (45%)	1.21	2 (22%) 0 0	19, 22, 34, 43	0
All	All	1017/1108 (91%)	0.16	31 (3%) 50 48	8, 16, 27, 52	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	P	507	HIS	5.8
2	S	507	HIS	4.9
1	C	339	SER	4.6
1	B	508	THR	4.4
2	P	506	GLY	4.3
1	A	508	THR	4.1
1	B	469	LYS	3.8
1	A	488	HIS	3.5
1	C	302	ARG	3.4
1	B	302	ARG	3.3
2	Q	508	LYS	3.0
2	S	506	GLY	2.9
1	D	294	ARG	2.9
1	A	294	ARG	2.8
1	D	272	HIS	2.7
1	A	458	HIS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	288	ARG	2.5
1	A	290	GLU	2.5
1	B	272	HIS	2.5
1	A	288	ARG	2.4
1	B	294	ARG	2.4
1	D	310	ARG	2.3
1	A	469	LYS	2.3
1	A	264	TYR	2.3
1	A	272	HIS	2.3
1	C	303[A]	GLU	2.2
1	D	458[A]	HIS	2.1
1	C	294	ARG	2.1
1	D	469	LYS	2.1
1	A	270[A]	ILE	2.1
1	B	288	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	98N	B	601	38/38	0.94	0.11	12,17,33,36	0
3	98N	A	601	38/38	0.94	0.12	14,21,35,36	0
3	98N	C	601	38/38	0.95	0.11	11,17,28,31	0
3	98N	D	601	38/38	0.95	0.11	12,18,29,30	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 98N B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 98N A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 98N C 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 98N D 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.