Full wwPDB X-ray Structure Validation Report (i)

Aug 7, 2023 – 07:22 PM EDT

PDB ID	:	1NUH
Title	:	The crystal structure of human phosphoglucose isomerase complexed with 5-phosphoarabinonate
Authors	:	Davies, C.
Deposited on	:	2003-01-31
Resolution	:	2.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	558	7% 75% 23%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4578 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glucose phosphate isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	555	4424	2819	778	809	18	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

- Molecule 3 is 5-PHOSPHOARABINONIC ACID (three-letter code: PA5) (formula: $C_5H_{11}O_9P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
3	A	1	15	5	9	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	109	109	109	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: glucose phosphate isomerase

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α, β, γ	94.40Å 94.40Å 137.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.51 47.82 – 2.51	Depositor EDS
% Data completeness (in resolution range)	98.4 (50.00-2.51) 97.7 (47.82-2.51)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$< I/\sigma(I) >$ [1]	2.72 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.214 , 0.268 0.243 , 0.281	Depositor DCC
R_{free} test set	1101 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å^2)	27.3	Xtriage
Anisotropy	0.212	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 38.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.52$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.89	EDS
Total number of atoms	4578	wwPDB-VP
Average B, all atoms (Å^2)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PA5, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.55	0/4532	0.82	11/6137 (0.2%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	505	ASP	CB-CG-OD2	7.10	124.69	118.30
1	A	150	ASP	CB-CG-OD2	6.76	124.38	118.30
1	A	371	ASP	CB-CG-OD2	6.72	124.35	118.30
1	A	404	ASP	CB-CG-OD2	6.14	123.83	118.30
1	A	290	ASP	CB-CG-OD2	5.84	123.55	118.30
1	A	139	ASP	CB-CG-OD2	5.68	123.41	118.30
1	A	53	ASP	CB-CG-OD2	5.59	123.34	118.30
1	A	126	ASP	CB-CG-OD2	5.58	123.33	118.30
1	A	457	ASP	CB-CG-OD2	5.47	123.22	118.30
1	A	355	ASP	CB-CG-OD2	5.31	123.08	118.30
1	A	510	ASP	CB-CG-OD2	5.22	123.00	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4424	0	4375	90	0
2	A	30	0	0	2	0
3	A	15	0	8	0	0
4	A	109	0	0	7	1
All	All	4578	0	4383	90	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:366:SER:OG	1:A:368:THR:HG23	1.68	0.94
1:A:14:GLN:HE22	1:A:17:ARG:NH1	1.78	0.81
1:A:6:ASP:OD1	4:A:609:HOH:O	1.99	0.80
1:A:523:LYS:O	1:A:526:PRO:HD2	1.84	0.77
1:A:530:GLY:O	1:A:552:ARG:NH2	2.18	0.77
1:A:77:GLU:HG3	4:A:615:HOH:O	1.83	0.77
1:A:249:THR:HG22	1:A:253:LYS:HE3	1.68	0.75
1:A:136:ARG:NE	2:A:560:SO4:O1	2.18	0.69
1:A:324:ILE:HG23	1:A:328:ASN:HD22	1.56	0.69
1:A:328:ASN:HD21	1:A:504:TRP:HE1	1.37	0.69
1:A:35:ARG:HD2	4:A:637:HOH:O	1.91	0.68
1:A:330:PHE:HB3	1:A:400:MET:CE	2.24	0.67
1:A:14:GLN:NE2	1:A:17:ARG:HH11	1.94	0.66
1:A:330:PHE:HB3	1:A:400:MET:HE3	1.78	0.66
1:A:391:TYR:HA	1:A:394:ILE:HD12	1.78	0.66
1:A:315:ALA:HB3	1:A:316:PRO:HD3	1.78	0.65
1:A:366:SER:OG	1:A:368:THR:CG2	2.45	0.64
1:A:283:ILE:O	1:A:287:VAL:HG22	1.98	0.62
1:A:29:PHE:CE2	1:A:35:ARG:HG2	2.36	0.61
1:A:387:GLN:HE22	1:A:427:ASN:HB3	1.66	0.60
1:A:14:GLN:CD	1:A:17:ARG:HH11	2.04	0.60
1:A:332:CYS:SG	1:A:400:MET:HE2	2.41	0.60
1:A:332:CYS:SG	1:A:400:MET:CE	2.90	0.60
1:A:33:LYS:HD3	1:A:33:LYS:H	1.67	0.59
1:A:314:ASN:OD1	1:A:316:PRO:HD2	2.03	0.58
1:A:74:ARG:HD2	4:A:609:HOH:O	2.02	0.58
1:A:14:GLN:NE2	1:A:17:ARG:NH1	2.47	0.58
1:A:45:THR:O	1:A:46:ASN:HB2	2.03	0.58
1:A:518:LYS:HG2	4:A:648:HOH:O	2.04	0.57
1:A:225:LYS:NZ	1:A:255:PHE:O	2.38	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:541:THR:O	1:A:545:ILE:HG13	2.06	0.56
1:A:482:THR:HB	1:A:483:PRO:HD2	1.90	0.54
1:A:19:HIS:O	1:A:20:ARG:C	2.47	0.53
1:A:183:VAL:HG12	1:A:220:ASN:ND2	2.24	0.53
1:A:324:ILE:HG23	1:A:328:ASN:ND2	2.24	0.52
1:A:330:PHE:CD1	1:A:400:MET:HE1	2.45	0.52
1:A:312:GLU:H	1:A:312:GLU:CD	2.13	0.51
1:A:353:GLN:O	1:A:357:GLU:HB3	2.11	0.50
1:A:462:LEU:HB3	1:A:463:PRO:HD3	1.93	0.50
1:A:245:LEU:HD13	1:A:279:ILE:HA	1.94	0.49
1:A:325:TRP:O	1:A:329:CYS:HB2	2.13	0.48
1:A:171:LYS:N	1:A:172:PRO:CD	2.76	0.48
1:A:186:ILE:HB	1:A:216:GLU:CG	2.44	0.47
1:A:537:HIS:HB2	1:A:542:ASN:OD1	2.15	0.46
1:A:102:LEU:HD11	1:A:269:VAL:HG22	1.98	0.46
1:A:191:ILE:HG23	1:A:192:ALA:N	2.30	0.46
1:A:206:ILE:HD13	1:A:281:LEU:HD23	1.98	0.46
1:A:171:LYS:HE3	1:A:286:HIS:HE1	1.81	0.46
1:A:332:CYS:SG	1:A:400:MET:HE3	2.56	0.46
1:A:490:VAL:O	1:A:494:GLU:HG3	2.16	0.46
1:A:400:MET:HE2	1:A:400:MET:HB3	1.79	0.45
1:A:103:ARG:CB	1:A:296:LEU:HB3	2.47	0.45
1:A:169:ALA:HA	1:A:343:TYR:HB3	1.99	0.45
1:A:394:ILE:O	1:A:471:ARG:NE	2.49	0.45
1:A:435:LEU:HD23	1:A:435:LEU:HA	1.80	0.44
1:A:103:ARG:HB2	1:A:296:LEU:HB3	1.99	0.44
1:A:152:ILE:HD12	1:A:152:ILE:N	2.32	0.44
1:A:141:LYS:O	2:A:561:SO4:O2	2.35	0.44
1:A:133:GLN:HE21	1:A:133:GLN:HB3	1.63	0.44
1:A:345:HIS:HA	1:A:382:PRO:HG3	2.00	0.44
1:A:98:LEU:HB2	1:A:268:TRP:CE3	2.53	0.43
1:A:164:LEU:C	1:A:164:LEU:HD23	2.38	0.43
1:A:229:LEU:HD23	1:A:229:LEU:HA	1.85	0.43
1:A:178:PRO:O	1:A:286:HIS:HD2	2.01	0.43
1:A:74:ARG:NH1	1:A:505:ASP:HB2	2.34	0.43
1:A:330:PHE:CB	1:A:400:MET:HE3	2.46	0.42
1:A:127:LYS:HE2	1:A:262:MET:O	2.19	0.42
1:A:255:PHE:HD1	1:A:257:ILE:HG13	1.84	0.42
1:A:446:LYS:HE2	1:A:446:LYS:HB3	1.87	0.42
1:A:138:GLY:O	1:A:141:LYS:HE2	2.19	0.42
1:A:155:GLY:O	1:A:160:ASP:HB2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:339:PRO:O	1:A:382:PRO:HA	2.19	0.42
1:A:84:PHE:HB3	1:A:307:ARG:CZ	2.49	0.42
1:A:405:PHE:HB3	1:A:428:PHE:CE1	2.55	0.41
1:A:33:LYS:H	1:A:33:LYS:CD	2.27	0.41
1:A:250:THR:O	1:A:254:GLU:HG3	2.21	0.41
1:A:407:ILE:HG13	1:A:408:PRO:HD2	2.02	0.41
1:A:239:ALA:HA	4:A:634:HOH:O	2.20	0.41
1:A:335:HIS:O	1:A:378:VAL:HA	2.20	0.41
1:A:125:LEU:HD23	1:A:125:LEU:HA	1.90	0.41
1:A:191:ILE:CG2	1:A:192:ALA:N	2.82	0.41
1:A:356:MET:HA	4:A:588:HOH:O	2.20	0.41
1:A:80:ARG:HG3	1:A:306:PHE:CE2	2.56	0.41
1:A:19:HIS:O	1:A:21:SER:N	2.54	0.41
1:A:116:ASP:O	1:A:119:PRO:HD2	2.21	0.41
1:A:212:PHE:HB3	1:A:251:LYS:HB3	2.03	0.40
1:A:387:GLN:HG2	1:A:391:TYR:CE2	2.56	0.40
1:A:195:LEU:HD13	1:A:227:TRP:CG	2.57	0.40
1:A:482:THR:HB	1:A:483:PRO:CD	2.51	0.40
1:A:524:ILE:H	1:A:524:ILE:HG12	1.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:567:HOH:O	4:A:567:HOH:O[7_555]	1.39	0.81

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	553/558 (99%)	530 (96%)	23 (4%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	473/477 (99%)	451 (95%)	22 (5%)	26 49

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	THR
1	A	6	ASP
1	A	30	ASP
1	A	33	LYS
1	A	34	ASP
1	A	35	ARG
1	A	41	LEU
1	A	43	LEU
1	A	65	ARG
1	A	103	ARG
1	A	133	GLN
1	A	180	VAL
1	A	184	SER
1	A	191	ILE
1	A	193	LYS
1	A	212	PHE
1	A	282	SER
1	A	368	THR
1	A	507	ASN
1	A	515	GLU
1	A	518	LYS
1	A	523	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	HIS
1	A	57	ASN
1	A	107	ASN
1	A	122	ASN
1	A	133	GLN
1	A	286	HIS
1	A	291	ASN
1	A	294	GLN
1	A	328	ASN
1	A	335	HIS
1	A	353	GLN
1	A	359	ASN
1	A	387	GLN
1	A	427	ASN
1	A	474	ASN
1	A	500	GLN
1	A	507	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	SO4	A	558	-	4,4,4	0.13	0	6,6,6	0.32	0
2	SO4	A	562	-	4,4,4	0.22	0	6,6,6	0.35	0
2	SO4	A	561	1	4,4,4	0.10	0	6,6,6	0.15	0
2	SO4	A	559	-	4,4,4	0.20	0	6,6,6	0.15	0
2	SO4	A	560	-	4,4,4	0.22	0	6,6,6	0.34	0
2	SO4	A	563	-	4,4,4	0.13	0	6,6,6	0.22	0
3	PA5	A	564	-	14,14,14	1.04	2 (14%)	19,20,20	1.26	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PA5	A	564	-	-	1/18/18/18	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	564	PA5	C2-C1	-2.31	1.49	1.52
3	A	564	PA5	O1-C1	-2.08	1.23	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	564	PA5	C3-C2-C1	-2.75	105.87	109.32
3	A	564	PA5	O1-C1-C2	2.36	119.66	113.27

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	564	PA5	C5-O5-P-O1P

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	561	SO4	1	0
2	A	560	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	555/558 (99%)	0.75	39 (7%)	16	16	15, 23, 39, 47	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	32	ASN	5.6
1	A	30	ASP	4.1
1	A	19	HIS	3.8
1	A	10	GLN	3.5
1	A	553	GLU	3.5
1	A	531	SER	3.3
1	A	451	ALA	3.2
1	A	18	GLU	3.2
1	A	519	GLN	3.0
1	A	554	ALA	3.0
1	A	379	TRP	2.9
1	A	456	GLU	2.8
1	A	336	ALA	2.8
1	A	269	VAL	2.8
1	A	460	ARG	2.8
1	A	133	GLN	2.7
1	A	138	GLY	2.7
1	A	534	VAL	2.6
1	A	33	LYS	2.6
1	A	23	LEU	2.6
1	A	378	VAL	2.4
1	A	347	PHE	2.4
1	A	8	GLN	2.4
1	A	438	GLY	2.4
1	A	490	VAL	2.3
1	A	257	ILE	2.3
1	A	231	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	12	LEU	2.2
1	A	28	LEU	2.2
1	A	34	ASP	2.2
1	A	29	PHE	2.2
1	A	14	GLN	2.2
1	A	542	ASN	2.2
1	A	390	PHE	2.1
1	A	5	ARG	2.1
1	A	437	ARG	2.1
1	A	258	ASP	2.0
1	A	290	ASP	2.0
1	A	139	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	SO4	A	560	5/5	0.82	0.53	53,54,55,55	0
2	SO4	A	562	5/5	0.83	0.34	53,53,54,54	0
2	SO4	A	559	5/5	0.85	0.37	55,56,57,57	0
2	SO4	A	558	5/5	0.89	0.30	48,48,50,51	0
2	SO4	A	561	5/5	0.91	0.40	60,60,60,61	0
2	SO4	A	563	5/5	0.93	0.37	58,58,58,59	0
3	PA5	A	564	15/15	0.93	0.19	21,27,30,31	0

6.5 Other polymers (i)

There are no such residues in this entry.