

Full wwPDB X-ray Structure Validation Report (i)

Mar 23, 2024 – 03:20 PM EDT

PDB ID : 1NWK

Title : CRYSTAL STRUCTURE OF MONOMERIC ACTIN IN THE ATP STATE

Authors : Graceffa, P.; Dominguez, R.

Deposited on : 2003-02-06

Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(\AA))$
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain				
1	A	375	76%	18%	• 6%		



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 3179 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Actin, alpha skeletal muscle.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	٨	354	Total	С	N	О	S	0	19	0
1	A	304	2853	1806	476	550	21	U	12	


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	73	HIC	HIS	modified residue	UNP P68135

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	4	Total Ca 4 4	0	0


• Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).





\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	Р	0	0
3	A	1	31	10	6	12	3	U	

• Molecule 4 is TETRAMETHYLRHODAMINE-5-MALEIMIDE (three-letter code: RHO) (formula: $C_{28}H_{25}N_3O_5$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf		
4	A	1	Total 36	C 28		O 5	0	0

• Molecule 5 is water.

\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
5	A	255	Total O 255 255	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Actin, alpha skeletal muscle





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	C 1 2 1	Depositor	
Cell constants	112.22Å 37.37Å 85.03Å	Depositor	
a, b, c, α , β , γ	90.00° 108.13° 90.00°	Depositor	
Resolution (Å)	24.35 - 1.85	Depositor	
% Data completeness	98.2 (24.35-1.85)	Depositor	
(in resolution range)	30.2 (24.00-1.00)		
R_{merge}	0.09	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	CNS 1.0	Depositor	
R, R_{free}	0.187 , 0.226	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	3179	wwPDB-VP	
Average B, all atoms (Å ²)	34.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, HIC, CA, RHO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain	Bond	$\mathbf{lengths}$	Bond angles		
MIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.30	0/2900	0.56	0/3929	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2853	0	2812	55	0
2	A	4	0	0	0	0
3	A	31	0	13	0	0
4	A	36	0	24	1	0
5	A	255	0	0	7	0
All	All	3179	0	2849	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:A:291[A]:LYS:HD3	1:A:325:MET:SD	2.21	0.80
1:A:299:MET:HE3	1:A:329:ILE:HG21	1.64	0.78
1:A:369:ILE:HD13	1:A:369:ILE:O	1.86	0.75
1:A:170:ALA:O	1:A:172:PRO:HD3	1.91	0.70
1:A:335:ARG:HA	1:A:338[B]:SER:OG	1.92	0.70
1:A:73:HIC:HZ3	1:A:179:ASP:HB3	1.78	0.64
1:A:334:GLU:H	1:A:334:GLU:CD	2.01	0.64
1:A:28:ARG:HG3	5:A:1432:HOH:O	1.99	0.61
1:A:113:LYS:HB3	1:A:371:HIS:CE1	2.38	0.58
1:A:37:ARG:HG2	1:A:37:ARG:HH11	1.70	0.57
1:A:275:HIS:CD2	1:A:275:HIS:H	2.23	0.57
1:A:314:GLN:HE22	1:A:327:ILE:HB	1.68	0.57
1:A:162:ASN:HD22	1:A:176:MET:HB2	1.70	0.56
1:A:334:GLU:HG2	5:A:1620:HOH:O	2.06	0.56
1:A:299:MET:HE3	1:A:329:ILE:CG2	2.35	0.56
1:A:190:MET:HG2	1:A:209:VAL:HG21	1.86	0.55
1:A:194[B]:THR:HA	1:A:198:TYR:O	2.07	0.55
1:A:194[A]:THR:HA	1:A:198:TYR:O	2.08	0.54
1:A:196:ARG:HD3	5:A:1577:HOH:O	2.07	0.54
1:A:250:ILE:HG13	1:A:253:GLU:HB2	1.90	0.54
1:A:299:MET:CE	1:A:313:MET:HG3	2.37	0.53
1:A:299:MET:HE1	1:A:313:MET:HG3	1.89	0.53
1:A:109:PRO:HG3	5:A:1550:HOH:O	2.09	0.52
1:A:297:ASN:HB2	1:A:329:ILE:HD13	1.92	0.52
1:A:303:THR:HG22	1:A:303:THR:O	2.10	0.52
1:A:80:ASP:O	1:A:84:LYS:HG2	2.11	0.49
1:A:162:ASN:ND2	1:A:277:THR:HB	2.28	0.49
1:A:210:ARG:O	1:A:214:GLU:HG3	2.12	0.49
1:A:37:ARG:HG2	1:A:37:ARG:NH1	2.28	0.47
1:A:257:CYS:HB3	1:A:258:PRO:CD	2.44	0.47
1:A:357:ILE:CD1	1:A:370:VAL:HA	2.47	0.45
1:A:252:ASN:HA	1:A:255:PHE:CE2	2.51	0.45
1:A:332:PRO:HA	1:A:333:PRO:HD3	1.92	0.45
1:A:356:TRP:CE2	4:A:1381:RHO:H161	2.52	0.45
1:A:10[B]:CYS:HB2	1:A:105:LEU:HD23	1.98	0.44
1:A:159:VAL:HG22	1:A:160:THR:N	2.32	0.44
1:A:115:ASN:O	1:A:119[B]:MET:HG3	2.18	0.44
1:A:180:LEU:HD11	1:A:260:THR:HG22	1.99	0.44
1:A:367:PRO:O	1:A:370:VAL:HG12	2.17	0.43
1:A:211:ASP:O	1:A:215:LYS:HG2	2.18	0.43
1:A:222[A]:ASP:OD2	1:A:225:ASN:OD1	2.36	0.42
1:A:303:THR:O	1:A:303:THR:CG2	2.67	0.42

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:121:GLN:O	1:A:125:GLU:HG3	2.19	0.42
1:A:334:GLU:CD	1:A:334:GLU:N	2.72	0.42
1:A:53:TYR:N	1:A:53:TYR:CD1	2.88	0.42
1:A:62:ARG:NH1	5:A:1594:HOH:O	2.51	0.42
1:A:113:LYS:HG3	5:A:1510:HOH:O	2.20	0.42
1:A:162:ASN:ND2	1:A:277:THR:CB	2.83	0.42
1:A:335:ARG:HA	1:A:338[A]:SER:HB3	2.00	0.41
1:A:330:ILE:N	1:A:330:ILE:HD12	2.35	0.41
1:A:12:ASN:HB3	1:A:71:ILE:CD1	2.51	0.41
1:A:207:GLU:OE1	1:A:210:ARG:NH2	2.54	0.41
1:A:162:ASN:HD21	1:A:277:THR:CB	2.34	0.40
1:A:37:ARG:HA	1:A:38:PRO:HD3	1.89	0.40
1:A:291[B]:LYS:HE3	5:A:1470:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	361/375~(96%)	353 (98%)	8 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/317 (98%)	309 (99%)	2 (1%)	86 83	

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	334	GLU
1	A	369	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	162	ASN
1	A	275	HIS
1	A	314	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Ros	$\operatorname{Res} \left \operatorname{Link} \right $	Bond lengths			Bond angles		
IVIOI	Туре	Chain	rtes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	HIC	A	73	1	8,11,12	0.85	0	6,14,16	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HIC	A	73	1	-	1/5/6/8	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	73	HIC	CA-CB-CG-ND1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	73	HIC	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain		n Res	Dag	Dag	Dag	Link	В	ond leng	gths	В	ond ang	gles
MOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2			
4	RHO	A	1381	-	39,40,40	3.13	24 (61%)	52,60,60	1.94	11 (21%)			
3	ANP	A	1380	2	29,33,33	1.82	8 (27%)	31,52,52	1.25	4 (12%)			

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	RHO	A	1381	-	-	6/18/58/58	0/5/5/5
3	ANP	A	1380	2	-	3/14/38/38	0/3/3/3

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
4	A	1381	RHO	C8-C14	11.02	1.45	1.35
4	A	1381	RHO	C-N	6.12	1.52	1.37
4	A	1381	RHO	C17-C8	4.42	1.55	1.49
4	A	1381	RHO	C25-C24	-4.06	1.41	1.48
3	A	1380	ANP	C2-N1	4.00	1.41	1.33
4	A	1381	RHO	C13-C12	-3.91	1.42	1.49
4	A	1381	RHO	C26-C27	-3.81	1.41	1.48
3	A	1380	ANP	PB-O2B	-3.64	1.47	1.56
4	A	1381	RHO	C24-N2	-3.55	1.34	1.40
4	A	1381	RHO	C27-N2	-3.41	1.34	1.40
3	A	1380	ANP	PG-O2G	-3.38	1.47	1.56
4	A	1381	RHO	C13-C14	-3.36	1.43	1.50
3	A	1380	ANP	PG-O3G	-3.29	1.47	1.56
3	A	1380	ANP	C2-N3	3.26	1.37	1.32
4	A	1381	RHO	C6-N	-2.85	1.39	1.45
4	A	1381	RHO	C18-C17	2.83	1.45	1.40
4	A	1381	RHO	C4-C3	2.80	1.44	1.39
4	A	1381	RHO	C7-N	-2.77	1.39	1.45
4	A	1381	RHO	C12-C11	2.70	1.44	1.35
4	A	1381	RHO	C19-C20	2.62	1.44	1.39
3	A	1380	ANP	O4'-C1'	2.53	1.44	1.41
4	A	1381	RHO	C21-C20	2.52	1.44	1.39
4	A	1381	RHO	C22-C17	2.51	1.43	1.39
4	A	1381	RHO	С5-С	2.30	1.43	1.39
4	A	1381	RHO	C3-C2	2.21	1.44	1.40
3	A	1380	ANP	PA-O2A	-2.13	1.45	1.55
4	A	1381	RHO	C19-C18	2.12	1.43	1.39
4	A	1381	RHO	C5-C4	2.11	1.42	1.38
4	A	1381	RHO	C16-N1	-2.07	1.39	1.46
4	A	1381	RHO	C15-N1	-2.05	1.39	1.46
3	A	1380	ANP	C4-N3	2.04	1.38	1.35
4	A	1381	RHO	C10-C9	2.01	1.42	1.36

All (15) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
4	A	1381	RHO	C18-C17-C8	-6.23	114.67	123.33
4	A	1381	RHO	C22-C17-C8	4.56	126.73	118.30
4	A	1381	RHO	C12-C13-C14	4.15	122.72	112.29
4	A	1381	RHO	C19-C20-N2	4.01	124.08	119.12
4	A	1381	RHO	C17-C8-C3	-3.85	115.45	119.94
3	A	1380	ANP	O4'-C1'-C2'	-3.18	102.28	106.93
3	A	1380	ANP	C5-C6-N6	3.18	125.18	120.35
4	A	1381	RHO	OH4-C24-N2	2.58	130.29	124.66
4	A	1381	RHO	C4-C3-C8	2.54	128.36	123.28
4	A	1381	RHO	OH5-C27-N2	2.45	130.01	124.66
4	A	1381	RHO	C4-C3-C2	-2.43	115.02	118.21
4	A	1381	RHO	C18-C19-C20	2.27	124.66	120.28
4	A	1381	RHO	OH3-C23-OH2	-2.10	118.69	123.35
3	A	1380	ANP	O2'-C2'-C3'	2.09	118.58	111.82
3	A	1380	ANP	O4'-C4'-C5'	-2.07	102.56	109.37

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1380	ANP	PB-N3B-PG-O1G
3	A	1380	ANP	PG-N3B-PB-O1B
4	A	1381	RHO	C21-C20-N2-C24
4	A	1381	RHO	C19-C20-N2-C24
4	A	1381	RHO	C21-C20-N2-C27
3	A	1380	ANP	PG-N3B-PB-O3A
4	A	1381	RHO	C19-C20-N2-C27
4	A	1381	RHO	С19-С18-С23-ОН3
4	A	1381	RHO	C19-C18-C23-OH2

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1381	RHO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

