



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 11, 2021 – 01:00 PM EDT

PDB ID : 2O1X
Title : 1-deoxy-D-xylulose 5-phosphate synthase (DXS) from *Deinococcus radiodurans*
Authors : Xiang, S.; Usunow, G.; Lange, G.; Busch, M.; Tong, L.
Deposited on : 2006-11-29
Resolution : 2.90 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	629	
1	B	629	
1	C	629	
1	D	629	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 16910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-deoxy-D-xylulose-5-phosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	578	4362	2753	771	821	17	0	0	0
1	B	579	4376	2761	775	823	17	0	0	0
1	C	538	4063	2558	724	766	15	0	0	0
1	D	530	4001	2521	715	750	15	0	0	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	130	THR	ALA	engineered mutation	UNP Q9RUB5
B	130	THR	ALA	engineered mutation	UNP Q9RUB5
C	130	THR	ALA	engineered mutation	UNP Q9RUB5
D	130	THR	ALA	engineered mutation	UNP Q9RUB5


- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		

- Molecule 3 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: C₁₂H₁₉N₄O₇P₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
3	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
3	B	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
3	C	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
3	D	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.26Å 154.06Å 124.88Å 90.00° 98.91° 90.00°	Depositor
Resolution (Å)	30.00 – 2.90 29.25 – 2.90	Depositor EDS
% Data completeness (in resolution range)	97.4 (30.00-2.90) 97.4 (29.25-2.90)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 2.90Å)	Xtrriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.209 , 0.272 0.207 , 0.266	Depositor DCC
R_{free} test set	3206 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	40.7	Xtrriage
Anisotropy	0.152	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	16910	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, TPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.53	0/4452	0.85	19/6049 (0.3%)
1	B	0.53	0/4465	0.87	16/6066 (0.3%)
1	C	0.43	0/4139	0.80	18/5621 (0.3%)
1	D	0.83	6/4076 (0.1%)	0.80	15/5534 (0.3%)
All	All	0.60	6/17132 (0.0%)	0.83	68/23270 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	36	GLU	CD-OE2	28.08	1.56	1.25
1	D	628	GLU	CD-OE1	23.04	1.50	1.25
1	D	628	GLU	CD-OE2	18.05	1.45	1.25
1	D	36	GLU	CD-OE1	16.34	1.43	1.25
1	D	28	GLU	CD-OE2	9.40	1.35	1.25

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	182	ASP	CB-CG-OD2	7.46	125.02	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	561	ASP	CB-CG-OD2	7.41	124.97	118.30
1	B	310	ASP	CB-CG-OD2	7.27	124.84	118.30
1	A	154	ASP	CB-CG-OD2	7.21	124.79	118.30
1	C	422	ASP	CB-CG-OD2	7.20	124.78	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	191	VAL	Peptide
1	B	539	LYS	Peptide
1	C	438	PHE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4362	0	4337	101	0
1	B	4376	0	4352	140	0
1	C	4063	0	4054	115	0
1	D	4001	0	3997	106	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	26	0	16	2	0
3	B	26	0	16	4	0
3	C	26	0	16	0	0
3	D	26	0	16	1	0
All	All	16910	0	16804	449	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 449 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:439:ASP:OD2	1:D:477:ARG:HD2	1.47	1.11
1:C:75:ARG:HG3	1:C:75:ARG:HH11	0.93	1.09
1:B:440:LEU:HD21	1:B:477:ARG:HD3	1.34	1.08
1:B:365:ARG:HG3	1:B:365:ARG:HH11	1.19	1.02
1:C:423:ARG:HD2	1:C:480:ARG:HG3	1.44	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	574/629 (91%)	536 (93%)	34 (6%)	4 (1%)	22	54
1	B	575/629 (91%)	531 (92%)	32 (6%)	12 (2%)	7	26
1	C	532/629 (85%)	473 (89%)	48 (9%)	11 (2%)	7	26
1	D	524/629 (83%)	470 (90%)	44 (8%)	10 (2%)	8	28
All	All	2205/2516 (88%)	2010 (91%)	158 (7%)	37 (2%)	9	31

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	507	ASP
1	B	194	MET
1	B	195	ASN
1	B	244	SER
1	B	439	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	452/493 (92%)	404 (89%)	48 (11%)	6	20
1	B	453/493 (92%)	404 (89%)	49 (11%)	6	20
1	C	422/493 (86%)	380 (90%)	42 (10%)	7	23
1	D	414/493 (84%)	384 (93%)	30 (7%)	14	39
All	All	1741/1972 (88%)	1572 (90%)	169 (10%)	8	25

5 of 169 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	351	GLU
1	D	71	SER
1	C	405	GLN
1	C	543	GLU
1	D	185	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	470	HIS
1	D	595	GLN
1	C	181	ASN
1	D	580	ASN
1	D	405	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TPP	D	1004	2	22,27,27	2.24	5 (22%)	29,40,40	1.92	11 (37%)
3	TPP	A	1001	2	22,27,27	1.82	5 (22%)	29,40,40	1.91	10 (34%)
3	TPP	B	1002	2	22,27,27	1.46	3 (13%)	29,40,40	1.81	8 (27%)
3	TPP	C	1003	2	22,27,27	2.95	7 (31%)	29,40,40	2.03	8 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TPP	D	1004	2	-	3/16/17/17	0/2/2/2
3	TPP	A	1001	2	-	3/16/17/17	0/2/2/2
3	TPP	B	1002	2	-	5/16/17/17	0/2/2/2
3	TPP	C	1003	2	-	4/16/17/17	0/2/2/2

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1003	TPP	C6-C5	11.51	1.56	1.50
3	D	1004	TPP	C6-C5	7.33	1.54	1.50
3	A	1001	TPP	PB-O1B	4.51	1.65	1.50
3	D	1004	TPP	PB-O1B	4.20	1.64	1.50
3	B	1002	TPP	C4-N3	-4.18	1.36	1.39

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1003	TPP	C6-C5-C4	4.80	131.28	127.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1003	TPP	C5-C4-N3	3.96	115.50	107.57
3	B	1002	TPP	CM2-C2'-N1'	3.86	121.38	117.14
3	B	1002	TPP	C5-C4-N3	3.62	114.83	107.57
3	C	1003	TPP	C7'-N3-C2	-3.61	118.83	125.35

There are no chirality outliers.

5 of 15 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	1001	TPP	C4'-C5'-C7'-N3
3	B	1002	TPP	C4'-C5'-C7'-N3
3	C	1003	TPP	C4'-C5'-C7'-N3
3	D	1004	TPP	C4'-C5'-C7'-N3
3	D	1004	TPP	C5-C6-C7-O7


There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1004	TPP	1	0
3	A	1001	TPP	2	0
3	B	1002	TPP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	578/629 (91%)	-0.48	6 (1%) 82 82	-5, 10, 34, 53	0
1	B	579/629 (92%)	-0.49	7 (1%) 79 79	-3, 11, 39, 75	0
1	C	538/629 (85%)	0.37	48 (8%) 9 7	6, 65, 139, 191	0
1	D	530/629 (84%)	0.35	52 (9%) 7 5	7, 60, 133, 175	0
All	All	2225/2516 (88%)	-0.08	113 (5%) 28 24	-5, 24, 121, 191	0

The worst 5 of 113 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	15	GLN	6.1
1	D	46	SER	6.1
1	D	49	GLY	5.6
1	C	321	ALA	5.5
1	C	8	SER	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	TPP	D	1004	26/26	0.92	0.16	40,44,48,48	0
2	MG	C	2003	1/1	0.93	0.12	33,33,33,33	0
2	MG	D	2004	1/1	0.93	0.12	35,35,35,35	0
3	TPP	C	1003	26/26	0.93	0.16	50,50,52,53	0
2	MG	B	2002	1/1	0.93	0.21	8,8,8,8	0
2	MG	A	2001	1/1	0.97	0.21	9,9,9,9	0
3	TPP	A	1001	26/26	0.98	0.11	2,2,4,5	0
3	TPP	B	1002	26/26	0.98	0.13	2,3,4,6	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around TPP C 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around TPP A 1001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.