



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2021 – 10:17 am BST

PDB ID : 7O26
Title : Complex-B bound [FeFe]-hydrogenase maturase HydE from *T. Maritima* (5'dA + Methionine)
Authors : Rohac, R.; Martin, L.; Liu, L.; Basu, D.; Tao, L.; Britt, R.D.; Rauchfuss, T.; Nicolet, Y.
Deposited on : 2021-03-30
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	

2 Entry composition [i](#)


There are 15 unique types of molecules in this entry. The entry contains 3492 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called [FeFe] hydrogenase maturase subunit HydE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	347	2891	1856	489	519	27	0	27	0

- Molecule 2 is CYSTEINE (three-letter code: CYS) (formula: C₃H₇NO₂S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	7	3	1	2	1	0	1

- Molecule 3 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		

- Molecule 4 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	1
			2	1	1		
4	A	1	Total	C	O	0	1
			2	1	1		

- Molecule 5 is CYANIDE ION (three-letter code: CYN) (formula: CN) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	1
			2	1	1		

- Molecule 6 is METHIONINE (three-letter code: MET) (formula: $C_5H_{11}NO_2S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			9	5	1	2	1		

- Molecule 7 is HYDROSULFURIC ACID (three-letter code: H2S) (formula: H_2S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total S 1 1	0	1
7	A	1	Total S 1 1	0	1

- Molecule 8 is 5'-DEOXYADENOSINE (three-letter code: 5AD) (formula: C₁₀H₁₃N₅O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C N O 18 10 5 3	0	0

- Molecule 9 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	Fe	S	0	1
			8	4	4		

- Molecule 10 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	1
			6	3	3		

- Molecule 11 is 3-[(3-CHOLAMIDOPROPYL)DIMETHYLAMMONIO]-1-PROPANESULFONATE (three-letter code: CPS) (formula: C₃₂H₅₈N₂O₇S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
11	A	1	Total	C	N	O	S	0	0
			42	32	2	7	1		
11	A	1	Total	C	N	O	S	0	0
			42	32	2	7	1		
11	A	1	Total	C	N	O		0	0
			29	24	1	4			
11	A	1	Total	C	N	O		0	0
			29	24	1	4			
11	A	1	Total	C	O			0	1
			50	44	6				

- Molecule 12 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	A	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 13 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	1	Total	I	0	0
			1	1		

- Molecule 14 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	A	1	Total	Cl	0	1
			2	2		


- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	339	Total	O	0	3
			341	341		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: [FeFe] hydrogenase maturase subunit HydE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.88Å 79.84Å 86.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.81 – 1.50 43.81 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.2 (43.81-1.50) 99.5 (43.81-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 1.50Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.178 , 0.216 0.178 , 0.216	Depositor DCC
R_{free} test set	2854 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtrriage
Anisotropy	0.337	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 38.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3492	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, CPS, FES, FE2, CYN, CMO, H2S, IOD, CL, 5AD, GOL, CSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/3021	0.66	0/4079

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2891	0	2948	22	0
2	A	7	0	3	2	0
3	A	1	0	0	0	0
4	A	4	0	0	0	0
5	A	2	0	0	0	0
6	A	9	0	8	1	0
7	A	2	0	0	1	0
8	A	18	0	13	3	0
9	A	8	0	0	0	0
10	A	6	0	3	1	0
11	A	192	0	264	6	0
12	A	8	0	0	0	0
13	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
14	A	2	0	0	1	0
15	A	341	0	0	7	1
All	All	3492	0	3239	31	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:322[C]:CYS:SG	15:A:820:HOH:O	2.30	0.89
1:A:321:PRO:HB3	11:A:416[A]:CPS:H11A	1.69	0.73
11:A:415:CPS:O3	15:A:501:HOH:O	2.12	0.67
1:A:334[B]:PRO:HB3	15:A:797:HOH:O	1.94	0.67
1:A:246:PHE:HD1	1:A:274[A]:ILE:HD11	1.62	0.64
1:A:41[A]:GLU:OE1	1:A:44:ARG:NH2	2.31	0.63
1:A:142[A]:ARG:HH12	1:A:189:GLU:CD	2.03	0.61
6:A:406:MET:SD	8:A:409:5AD:H5'2	2.46	0.56
1:A:94:ARG:HG3	1:A:129[B]:MET:SD	2.46	0.55
1:A:306:TYR:CE1	1:A:309[B]:LYS:HG2	2.46	0.51
10:A:411[A]:GOL:H12	15:A:714:HOH:O	2.10	0.51
1:A:217:LEU:HD11	15:A:509:HOH:O	2.11	0.51
1:A:306:TYR:CE1	1:A:309[A]:LYS:HG2	2.48	0.48
1:A:246:PHE:CD1	1:A:274[A]:ILE:HD11	2.46	0.48
1:A:142[B]:ARG:NH1	1:A:189:GLU:OE2	2.43	0.48
1:A:12:ARG:NH1	1:A:14:GLU:OE2	2.47	0.47
1:A:26:ILE:O	1:A:251:LYS:NZ	2.48	0.46
11:A:413:CPS:H31	11:A:413:CPS:H28B	1.60	0.46
2:A:401[A]:CYS:SG	8:A:409:5AD:H5'1	2.55	0.46
11:A:415:CPS:H3	11:A:415:CPS:H1	1.75	0.46
1:A:311[A]:CYS:HB3	7:A:407[A]:H2S:S	2.58	0.44
2:A:401[A]:CYS:HA	15:A:724:HOH:O	2.18	0.44
1:A:54:ARG:HB2	1:A:291[A]:MET:HG2	2.00	0.44
11:A:415:CPS:O3	11:A:415:CPS:H14	2.18	0.43
1:A:7:LEU:O	1:A:11:GLU:HG3	2.18	0.43
1:A:306:TYR:OH	14:A:419[C]:CL:CL	2.63	0.43
1:A:309[A]:LYS:HE3	1:A:309[A]:LYS:HB3	1.75	0.43
1:A:229:PRO:HG3	1:A:274[A]:ILE:HD12	2.01	0.43
11:A:416[A]:CPS:H8	15:A:715:HOH:O	2.20	0.42
1:A:229:PRO:O	8:A:409:5AD:H2	2.21	0.41
1:A:292:PRO:HB2	1:A:319[A]:CYS:HB3	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
15:A:786:HOH:O	15:A:793:HOH:O[3_745]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	372/348 (107%)	362 (97%)	10 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	317/303 (105%)	305 (96%)	12 (4%)	33 7

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ARG
1	A	22	GLU
1	A	274[A]	ILE
1	A	274[B]	ILE
1	A	290	ILE
1	A	309[A]	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	309[B]	LYS
1	A	319[A]	CYS
1	A	319[B]	CYS
1	A	322[A]	CYS
1	A	322[B]	CYS
1	A	322[C]	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	A	183	1	3,6,7	0.51	0	0,6,8	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	183	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 4 are monoatomic and 2 are modelled with single atom - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CMO	A	404[C]	-	0,1,1	0.00	-	-		
11	CPS	A	416[A]	-	28,28,45	0.82	0	46,46,70	1.39	8 (17%)
4	CMO	A	403[C]	-	0,1,1	0.00	-	-		
11	CPS	A	415	-	32,32,45	0.71	1 (3%)	51,51,70	1.46	7 (13%)
11	CPS	A	413	-	45,45,45	1.29	1 (2%)	69,70,70	1.07	5 (7%)
12	SF4	A	417	1,6	0,12,12	0.00	-	-		
11	CPS	A	414	-	32,32,45	0.58	0	51,51,70	0.99	3 (5%)
2	CYS	A	401[A]	3	3,6,6	0.76	0	1,7,7	1.65	0
9	FES	A	410[B]	1	0,4,4	0.00	-	-		
11	CPS	A	412	-	45,45,45	1.45	3 (6%)	69,70,70	1.03	3 (4%)
6	MET	A	406	12	4,8,8	0.40	0	2,9,9	0.71	0
9	FES	A	410[A]	1,7	0,4,4	0.00	-	-		
5	CYN	A	405[A]	-	0,1,1	0.00	-	-		
11	CPS	A	416[B]	-	28,28,45	0.70	0	46,46,70	1.18	2 (4%)
8	5AD	A	409	-	17,20,20	3.86	8 (47%)	15,30,30	1.99	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	CPS	A	416[A]	-	-	2/4/69/90	0/4/4/4
11	CPS	A	415	-	-	1/9/74/90	0/4/4/4
11	CPS	A	413	-	-	13/25/90/90	0/4/4/4
12	SF4	A	417	1,6	-	-	0/6/5/5
11	CPS	A	414	-	-	0/9/74/90	0/4/4/4
2	CYS	A	401[A]	3	-	0/2/6/6	-
11	CPS	A	412	-	-	1/25/90/90	0/4/4/4
9	FES	A	410[B]	1	-	-	0/1/1/1
6	MET	A	406	12	-	1/4/8/8	-
9	FES	A	410[A]	1,7	-	-	0/1/1/1
11	CPS	A	416[B]	-	-	4/4/69/90	0/4/4/4
8	5AD	A	409	-	-	0/0/20/20	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	409	5AD	C3'-C2'	-9.57	1.27	1.53
11	A	412	CPS	C32-S	-7.90	1.66	1.77
8	A	409	5AD	C3'-C4'	7.43	1.63	1.52
11	A	413	CPS	C32-S	-7.21	1.67	1.77
8	A	409	5AD	O4'-C4'	-5.45	1.28	1.44
8	A	409	5AD	O4'-C1'	4.93	1.48	1.41
8	A	409	5AD	O2'-C2'	4.02	1.52	1.43
8	A	409	5AD	C2'-C1'	3.31	1.58	1.53
8	A	409	5AD	C6-N6	3.09	1.45	1.34
8	A	409	5AD	O3'-C3'	3.07	1.50	1.43
11	A	412	CPS	C5-C6	-2.36	1.51	1.55
11	A	415	CPS	C2-C19	-2.24	1.52	1.56
11	A	412	CPS	C28-N2	-2.03	1.45	1.50

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	A	416[B]	CPS	C5-C9-C20	-4.85	113.70	119.49
8	A	409	5AD	N3-C2-N1	-4.50	121.65	128.68
11	A	415	CPS	C3-C19-C2	-4.14	109.46	113.73
8	A	409	5AD	C5'-C4'-C3'	-4.04	111.46	115.70
11	A	412	CPS	O1S-S-C32	3.77	111.46	106.92
11	A	416[A]	CPS	C3-C19-C2	-3.41	110.21	113.73
11	A	416[A]	CPS	C5-C9-C20	-3.12	115.76	119.49
11	A	415	CPS	C6-C5-C4	2.96	110.16	107.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	A	415	CPS	C1-C2-C15	2.93	112.11	107.77
11	A	413	CPS	O2S-S-C32	2.69	110.12	105.77
11	A	416[A]	CPS	C2-C19-C18	-2.69	108.93	111.82
8	A	409	5AD	C3'-C2'-C1'	2.62	104.92	100.98
11	A	413	CPS	O3S-S-C32	2.61	110.06	106.92
11	A	415	CPS	C19-C3-C4	-2.60	110.86	114.30
11	A	416[A]	CPS	C10-C5-C4	2.53	111.64	109.07
11	A	412	CPS	C14-C13-C12	-2.52	107.55	110.55
11	A	415	CPS	C19-C18-C6	-2.47	106.32	109.71
11	A	415	CPS	C1-C2-C19	-2.46	107.48	111.35
11	A	416[A]	CPS	C11-C2-C19	-2.44	107.81	111.18
11	A	414	CPS	C10-C5-C4	2.34	111.45	109.07
11	A	414	CPS	C10-C5-C6	-2.32	107.59	111.21
11	A	416[A]	CPS	C14-C13-C12	-2.30	107.81	110.55
11	A	416[A]	CPS	C5-C6-C18	-2.28	111.83	114.74
11	A	414	CPS	C23-C22-C20	-2.24	110.42	114.52
11	A	415	CPS	C11-C2-C19	-2.24	108.10	111.18
11	A	413	CPS	O1S-S-C32	2.22	109.59	106.92
11	A	413	CPS	O1S-S-O3S	-2.20	106.34	113.95
11	A	416[A]	CPS	C1-C2-C15	2.18	110.98	107.77
11	A	412	CPS	C23-C22-C20	2.08	118.32	114.52
11	A	416[B]	CPS	C10-C5-C9	-2.06	107.99	111.21
11	A	413	CPS	C26-C27-N2	-2.01	111.12	115.38

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	406	MET	C-CA-CB-CG
11	A	413	CPS	C31-C32-S-O3S
11	A	413	CPS	C31-C32-S-O1S
11	A	413	CPS	O1-C24-N1-C25
11	A	413	CPS	C31-C32-S-O2S
11	A	413	CPS	C23-C24-N1-C25
11	A	416[B]	CPS	C21-C20-C9-C5
11	A	416[B]	CPS	C22-C20-C9-C5
11	A	413	CPS	C9-C20-C22-C23
11	A	416[B]	CPS	C21-C20-C9-C8
11	A	416[B]	CPS	C22-C20-C9-C8
11	A	413	CPS	C26-C27-N2-C29
11	A	413	CPS	C26-C27-N2-C28
11	A	413	CPS	C31-C30-N2-C27

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
11	A	413	CPS	C26-C27-N2-C30
11	A	415	CPS	C9-C20-C22-C23
11	A	413	CPS	C31-C30-N2-C29
11	A	413	CPS	C20-C22-C23-C24
11	A	413	CPS	C31-C30-N2-C28
11	A	412	CPS	C9-C20-C22-C23
11	A	416[A]	CPS	C21-C20-C9-C5
11	A	416[A]	CPS	C22-C20-C9-C5


There are no ring outliers.




6 monomers are involved in 10 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	A	416[A]	CPS	2	0
11	A	415	CPS	3	0
11	A	413	CPS	1	0
2	A	401[A]	CYS	2	0
6	A	406	MET	1	0
8	A	409	5AD	3	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	346/348 (99%)	-0.40	2 (0%) 89 91	13, 21, 41, 71	4 (1%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	345	PHE	4.2
1	A	347	THR	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	A	183	7/8	0.98	0.05	16,17,25,25	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
11	CPS	A	415	29/42	0.55	0.27	40,45,46,50	0
11	CPS	A	416[A]	25/42	0.78	0.20	23,25,27,31	25
11	CPS	A	416[B]	25/42	0.78	0.20	31,35,42,43	25
11	CPS	A	413	42/42	0.86	0.15	18,23,55,56	3
5	CYN	A	405[A]	2/2	0.86	0.16	22,22,22,22	2
10	GOL	A	411[A]	6/6	0.87	0.10	30,34,38,45	1
7	H2S	A	408[A]	1/1	0.89	0.23	20,20,20,20	1
6	MET	A	406	9/9	0.89	0.12	31,33,36,36	0
8	5AD	A	409	18/18	0.92	0.07	17,20,28,31	0
11	CPS	A	414	29/42	0.94	0.07	18,21,37,39	0
7	H2S	A	407[A]	1/1	0.94	0.05	18,18,18,18	1
4	CMO	A	403[C]	2/2	0.95	0.13	19,19,19,21	2
9	FES	A	410[A]	4/4	0.95	0.14	21,23,25,26	4
9	FES	A	410[B]	4/4	0.95	0.14	23,28,28,29	4
4	CMO	A	404[C]	2/2	0.95	0.10	19,19,19,21	0
11	CPS	A	412	42/42	0.95	0.08	14,17,28,30	5
2	CYS	A	401[A]	7/7	0.96	0.08	20,26,32,32	7
12	SF4	A	417	8/8	0.98	0.05	19,20,21,23	0
14	CL	A	419[B]	1/1	0.98	0.05	24,24,24,24	1
14	CL	A	419[C]	1/1	0.98	0.05	31,31,31,31	1
3	FE2	A	402	1/1	1.00	0.04	21,21,21,21	1
13	IOD	A	418	1/1	1.00	0.05	22,22,22,22	1

6.5 Other polymers [i](#)

There are no such residues in this entry.