

Full wwPDB X-ray Structure Validation Report (i)

Aug 17, 2022 – 04:27 PM EDT

PDB ID : 4O5T

Title: Crystal structure of Diels-Alderase CE20 in complex with a product analog

Authors: Beck, T.; Preiswerk, N.; Mayer, C.; Hilvert, D.

Deposited on : 2013-12-20

Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.29

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.29

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\#\text{Entries, resolution range}(\mathring{\mathbf{A}}))$
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	337	88%	6% • 5%
1	В	337	87%	9% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	XDA	В	400	-	-	_	X



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9279 atoms, of which 4451 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Diisopropyl-fluorophosphatase.

Mol	Chain	Residues			Atom	S			ZeroOcc	AltConf	Trace
1	A	319	Total 4581	C 1510	H 2211	N 402	O 441	S 17	0	4	0
1	В	326	Total 4557	C 1516		N 400	O 436	S 17	0	0	0

There are 116 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	MET	VAL	engineered mutation	UNP Q7SIG4
A	21	THR	GLU	engineered mutation	UNP Q7SIG4
A	33	VAL	ILE	engineered mutation	UNP Q7SIG4
A	36	SER	PRO	SEE REMARK 999	UNP Q7SIG4
A	37	PRO	GLU	SEE REMARK 999	UNP Q7SIG4
A	38	LEU	VAL	SEE REMARK 999	UNP Q7SIG4
A	39	SER	GLU	SEE REMARK 999	UNP Q7SIG4
A	40	GLU	VAL	SEE REMARK 999	UNP Q7SIG4
A	41	ALA	ASN	SEE REMARK 999	UNP Q7SIG4
A	42	LEU	GLY	SEE REMARK 999	UNP Q7SIG4
A	43	ILE	LYS	SEE REMARK 999	UNP Q7SIG4
A	44	ASN	PRO	SEE REMARK 999	UNP Q7SIG4
A	45	ALA	ALA	SEE REMARK 999	UNP Q7SIG4
A	46	ASN	-	SEE REMARK 999	UNP Q7SIG4
A	47	SER	-	SEE REMARK 999	UNP Q7SIG4
A	48	LEU	-	SEE REMARK 999	UNP Q7SIG4
A	49	ALA	-	SEE REMARK 999	UNP Q7SIG4
A	50	GLU	-	SEE REMARK 999	UNP Q7SIG4
A	51	ALA	-	SEE REMARK 999	UNP Q7SIG4
A	52	TYR	-	SEE REMARK 999	UNP Q7SIG4
A	53	GLU	-	SEE REMARK 999	UNP Q7SIG4
A	54	ALA	-	SEE REMARK 999	UNP Q7SIG4
A	55	ARG	-	SEE REMARK 999	UNP Q7SIG4
A	56	SER	-	SEE REMARK 999	UNP Q7SIG4
A	57	ASP	-	SEE REMARK 999	UNP Q7SIG4



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	58	ALA	-	SEE REMARK 999	UNP Q7SIG4
A	63	HIS	ARG	engineered mutation	UNP Q7SIG4
A	85	SER	ILE	engineered mutation	UNP Q7SIG4
A	87	ILE	ALA	engineered mutation	UNP Q7SIG4
A	121	ASN	LYS	engineered mutation	UNP Q7SIG4
A	128	CYS	ARG	engineered mutation	UNP Q7SIG4
A	133	ALA	ASN	engineered mutation	UNP Q7SIG4
A	134	TYR	ASP	engineered mutation	UNP Q7SIG4
A	151	GLY	GLU	engineered mutation	UNP Q7SIG4
A	157	PHE	TYR	engineered mutation	UNP Q7SIG4
A	159	ILE	ARG	engineered mutation	UNP Q7SIG4
A	161	LEU	MET	engineered mutation	UNP Q7SIG4
A	162	ARG	GLN	engineered mutation	UNP Q7SIG4
A	186	CYS	PHE	engineered mutation	UNP Q7SIG4
A	188	ALA	ASN	engineered mutation	UNP Q7SIG4
A	208	GLN	THR	engineered mutation	UNP Q7SIG4
A	223	ASN	LYS	engineered mutation	UNP Q7SIG4
A	238	LYS	GLU	engineered mutation	UNP Q7SIG4
A	242	ALA	ASP	engineered mutation	UNP Q7SIG4
A	245	VAL	ASP	engineered mutation	UNP Q7SIG4
A	284	ALA	SER	engineered mutation	UNP Q7SIG4
A	301	ASP	GLU	engineered mutation	UNP Q7SIG4
A	322	SER	LEU	engineered mutation	UNP Q7SIG4
A	328	GLY	-	expression tag	UNP Q7SIG4
A	329	SER	-	expression tag	UNP Q7SIG4
A	330	LEU	_	expression tag	UNP Q7SIG4
A	331	GLU	-	expression tag	UNP Q7SIG4
A	332	HIS	-	expression tag	UNP Q7SIG4
A	333	HIS	_	expression tag	UNP Q7SIG4
A	334	HIS	-	expression tag	UNP Q7SIG4
A	335	HIS	-	expression tag	UNP Q7SIG4
A	336	HIS	_	expression tag	UNP Q7SIG4
A	337	HIS	-	expression tag	UNP Q7SIG4
В	13	MET	VAL	engineered mutation	UNP Q7SIG4
В	21	THR	GLU	engineered mutation	UNP Q7SIG4
В	33	VAL	ILE	engineered mutation	UNP Q7SIG4
В	36	SER	PRO	SEE REMARK 999	UNP Q7SIG4
В	37	PRO	GLU	SEE REMARK 999	UNP Q7SIG4
В	38	LEU	VAL	SEE REMARK 999	UNP Q7SIG4
В	39	SER	GLU	SEE REMARK 999	UNP Q7SIG4
В	40	GLU	VAL	SEE REMARK 999	UNP Q7SIG4
В	41	ALA	ASN	SEE REMARK 999	UNP Q7SIG4



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	42	LEU	GLY	SEE REMARK 999	UNP Q7SIG4
В	43	ILE	LYS	SEE REMARK 999	UNP Q7SIG4
В	44	ASN	PRO	SEE REMARK 999	UNP Q7SIG4
В	45	ALA	ALA	SEE REMARK 999	UNP Q7SIG4
В	46	ASN	-	SEE REMARK 999	UNP Q7SIG4
В	47	SER	-	SEE REMARK 999	UNP Q7SIG4
В	48	LEU	-	SEE REMARK 999	UNP Q7SIG4
В	49	ALA	-	SEE REMARK 999	UNP Q7SIG4
В	50	GLU	-	SEE REMARK 999	UNP Q7SIG4
В	51	ALA	-	SEE REMARK 999	UNP Q7SIG4
В	52	TYR	-	SEE REMARK 999	UNP Q7SIG4
В	53	GLU	-	SEE REMARK 999	UNP Q7SIG4
В	54	ALA	-	SEE REMARK 999	UNP Q7SIG4
В	55	ARG	-	SEE REMARK 999	UNP Q7SIG4
В	56	SER	-	SEE REMARK 999	UNP Q7SIG4
В	57	ASP	-	SEE REMARK 999	UNP Q7SIG4
В	58	ALA	-	SEE REMARK 999	UNP Q7SIG4
В	63	HIS	ARG	engineered mutation	UNP Q7SIG4
В	85	SER	ILE	engineered mutation	UNP Q7SIG4
В	87	ILE	ALA	engineered mutation	UNP Q7SIG4
В	121	ASN	LYS	engineered mutation	UNP Q7SIG4
В	128	CYS	ARG	engineered mutation	UNP Q7SIG4
В	133	ALA	ASN	engineered mutation	UNP Q7SIG4
В	134	TYR	ASP	engineered mutation	UNP Q7SIG4
В	151	GLY	GLU	engineered mutation	UNP Q7SIG4
В	157	PHE	TYR	engineered mutation	UNP Q7SIG4
В	159	ILE	ARG	engineered mutation	UNP Q7SIG4
В	161	LEU	MET	engineered mutation	UNP Q7SIG4
В	162	ARG	GLN	engineered mutation	UNP Q7SIG4
В	186	CYS	PHE	engineered mutation	UNP Q7SIG4
В	188	ALA	ASN	engineered mutation	UNP Q7SIG4
В	208	GLN	THR	engineered mutation	UNP Q7SIG4
В	223	ASN	LYS	engineered mutation	UNP Q7SIG4
В	238	LYS	GLU	engineered mutation	UNP Q7SIG4
В	242	ALA	ASP	engineered mutation	UNP Q7SIG4
В	245	VAL	ASP	engineered mutation	UNP Q7SIG4
В	284	ALA	SER	engineered mutation	UNP Q7SIG4
В	301	ASP	GLU	engineered mutation	UNP Q7SIG4
В	322	SER	LEU	engineered mutation	UNP Q7SIG4
В	328	GLY	-	expression tag	UNP Q7SIG4
В	329	SER	-	expression tag	UNP Q7SIG4
В	330	LEU	-	expression tag	UNP Q7SIG4



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	331	GLU	-	expression tag	UNP Q7SIG4
В	332	HIS	-	expression tag	UNP Q7SIG4
В	333	HIS	-	expression tag	UNP Q7SIG4
В	334	HIS	-	expression tag	UNP Q7SIG4
В	335	HIS	-	expression tag	UNP Q7SIG4
В	336	HIS	-	expression tag	UNP Q7SIG4
В	337	HIS	-	expression tag	UNP Q7SIG4

 $\bullet \ \, Molecule \ 2 \ is \ 4-\{[2-(phosphonooxy)ethyl]carbamoyl\} benzyl \ [(1R,6S)-6-(dimethylcarbamoyl) \ \, cyclohex-2-en-1-yl]carbamate \ (three-letter \ code: \ XDA) \ (formula: \ C_{20}H_{28}N_3O_8P).$



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf				
2	٨	1	Total	С	Н	N	О	Р	0	0		
	A	1	58	20	26	3	8	1	0			
9	D	1	Total	С	Н	N	О	Р	0	0		
2	D	В	1	58	20	26	3	8	1	U	U	

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	15	Total O 15 15	0	0
3	В	10	Total O 10 10	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Diisopropyl-fluorophosphatase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	88.65Å 128.94Å 106.53Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.08 - 2.90	Depositor
Resolution (A)	43.04 - 2.90	EDS
% Data completeness	97.0 (43.08-2.90)	Depositor
(in resolution range)	97.0 (43.04-2.90)	EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.50 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
D D.	0.228 , 0.288	Depositor
R, R_{free}	0.227 , 0.277	DCC
R_{free} test set	655 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	78.2	Xtriage
Anisotropy	0.521	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37, 60.7	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9279	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.84% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: XDA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.49	0/2439	0.65	0/3324	
1	В	0.50	0/2431	0.66	0/3316	
All	All	0.50	0/4870	0.66	0/6640	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2370	2211	2202	16	0
1	В	2369	2188	2188	21	1
2	A	32	26	28	4	0
2	В	32	26	28	15	0
3	A	15	0	0	0	0
3	В	10	0	0	0	0
All	All	4828	4451	4446	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

A + a 1	A 4 a 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	$overlap (\AA)$
2:B:400:XDA:C13	2:B:400:XDA:H10B	1.54	1.36
2:B:400:XDA:C13	2:B:400:XDA:C10	2.27	1.13
2:B:400:XDA:H10B	2:B:400:XDA:O14	1.48	1.11
2:B:400:XDA:H10B	2:B:400:XDA:N12	1.67	1.09
2:B:400:XDA:H10B	2:B:400:XDA:C3	1.83	1.04
2:B:400:XDA:H10B	2:B:400:XDA:H3	1.51	0.93
2:A:400:XDA:C13	2:A:400:XDA:H10B	1.98	0.93
2:B:400:XDA:C10	2:B:400:XDA:C3	2.48	0.91
2:B:400:XDA:C10	2:B:400:XDA:N12	2.37	0.87
1:B:299:GLU:OE2	1:B:301:ASP:HB2	1.77	0.85
2:B:400:XDA:C10	2:B:400:XDA:O14	2.22	0.80
1:A:94[A]:ASP:O	1:A:95[A]:ALA:CB	2.35	0.75
1:A:94[A]:ASP:O	1:A:95[A]:ALA:HB3	1.88	0.74
2:A:400:XDA:H10B	2:A:400:XDA:O14	1.89	0.71
1:A:280:PHE:CG	1:A:299:GLU:HG3	2.28	0.69
2:B:400:XDA:C10	2:B:400:XDA:H3	2.16	0.69
1:B:86:PRO:O	1:B:87:ILE:HD12	1.94	0.67
2:B:400:XDA:C19	2:B:400:XDA:H26B	2.22	0.67
2:B:400:XDA:N12	2:B:400:XDA:N9	2.42	0.65
1:B:193:ARG:NH1	1:B:267:PRO:O	2.32	0.63
1:A:299:GLU:OE2	1:A:301:ASP:HB2	1.99	0.62
1:A:46:ASN:O	1:A:47:SER:CB	2.48	0.62
1:A:280:PHE:CD1	1:A:299:GLU:HG3	2.35	0.61
1:B:48:LEU:O	1:B:51:ALA:N	2.34	0.60
1:B:280:PHE:CG	1:B:299:GLU:HG3	2.37	0.59
1:B:299:GLU:OE2	1:B:301:ASP:CB	2.49	0.58
1:B:86:PRO:C	1:B:87:ILE:HD12	2.25	0.57
2:B:400:XDA:C19	2:B:400:XDA:C26	2.86	0.53
1:B:143:LEU:HD23	1:B:145:ILE:HD11	1.91	0.51
2:B:400:XDA:C26	2:B:400:XDA:H19	2.41	0.50
1:A:94[B]:ASP:OD1	1:A:94[B]:ASP:N	2.39	0.50
1:A:299:GLU:OE2	1:A:301:ASP:CB	2.59	0.50
1:A:108:LEU:HD23	1:A:118:GLU:HA	1.93	0.49
1:B:161:LEU:O	1:B:185:GLN:NE2	2.44	0.49
1:A:257:TRP:CD1	2:A:400:XDA:H11B	2.48	0.49
1:B:182:THR:OG1	1:B:183:ALA:N	2.45	0.47
1:B:206:ALA:HA	1:B:213:LEU:HD23	1.97	0.47
2:B:400:XDA:H26B	2:B:400:XDA:H19	1.94	0.46
1:A:215:SER:HB3	1:A:230:TRP:HB2	1.97	0.46
1:A:86:PRO:HB3	1:A:100:VAL:CG1	2.46	0.46
1:B:193:ARG:HB3	1:B:202:GLN:HB3	1.98	0.46



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
2:A:400:XDA:C13	2:A:400:XDA:C10	2.85	0.44
1:B:289:LYS:O	1:B:292:THR:OG1	2.32	0.44
1:B:280:PHE:CD2	1:B:299:GLU:HG3	2.53	0.43
1:A:247:ASP:OD1	1:A:251:ASN:N	2.50	0.43
1:B:193:ARG:HH21	1:B:246:PHE:HE1	1.65	0.43
1:B:276:ILE:HD11	1:B:310:TRP:CZ3	2.53	0.43
1:B:195:MET:HB2	1:B:197:ASP:OD1	2.18	0.43
1:A:181:ASP:OD2	1:A:216:TYR:OH	2.29	0.42
1:A:21:THR:OG1	1:A:22:GLY:N	2.51	0.42
1:B:17:ILE:O	1:B:303:ASN:HB3	2.19	0.41
1:B:299:GLU:OE2	1:B:301:ASP:N	2.50	0.41
1:A:278:CYS:HB3	1:A:280:PHE:CE1	2.56	0.40
1:B:159:ILE:HG23	1:B:161:LEU:HB2	2.02	0.40
1:B:102:ASP:OD1	1:B:105:LEU:N	2.53	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:B:197:ASP:OD2	1:B:197:ASP:OD2[3_554]	2.13	0.07

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percent	tiles
1	A	319/337~(95%)	302 (95%)	15 (5%)	2 (1%)	25	58
1	В	322/337~(96%)	306 (95%)	16 (5%)	0	100	100
All	All	641/674 (95%)	608 (95%)	31 (5%)	2 (0%)	41	71

All (2) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	47	SER
1	A	49	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	234/277 (84%)	232 (99%)	2 (1%)	78 93		
1	В	228/277 (82%)	222 (97%)	6 (3%)	46 77		
All	All	462/554 (83%)	454 (98%)	8 (2%)	60 86		

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	161	LEU
1	A	301	ASP
1	В	75	CYS
1	В	78	GLU
1	В	158	THR
1	В	159	ILE
1	В	161	LEU
1	В	327	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trme	oe Chain	Dog	Timle	Bond lengths			Bond angles		
MIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XDA	A	400	-	33,33,33	1.16	1 (3%)	39,45,45	1.57	8 (20%)
2	XDA	В	400	-	33,33,33	1.14	1 (3%)	39,45,45	1.32	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XDA	A	400	-	-	7/29/40/40	0/2/2/2
2	XDA	В	400	-	-	10/29/40/40	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
2	A	400	XDA	O15-C13	5.96	1.46	1.35
2	В	400	XDA	O15-C13	5.67	1.46	1.35

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
2	В	400	XDA	O15-C13-N12	4.46	119.56	110.50
2	A	400	XDA	C20-C23-N25	3.75	125.14	117.09
2	A	400	XDA	O15-C13-N12	3.32	117.24	110.50
2	В	400	XDA	O15-C13-O14	-3.08	118.34	124.25



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
2	A	400	XDA	C16-O15-C13	3.05	122.73	115.93
2	A	400	XDA	O24-C23-C20	-2.92	115.72	120.94
2	A	400	XDA	C6-C1-C2	-2.77	118.68	123.61
2	A	400	XDA	O15-C13-O14	-2.44	119.57	124.25
2	В	400	XDA	C20-C23-N25	2.23	121.87	117.09
2	A	400	XDA	C6-C5-C4	2.11	114.05	110.98
2	В	400	XDA	C16-O15-C13	2.03	120.45	115.93
2	A	400	XDA	C3-N12-C13	2.02	124.10	121.61

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	400	XDA	C3-C4-C7-O8
2	В	400	XDA	C3-C4-C7-N9
2	В	400	XDA	C27-O28-P29-O31
2	В	400	XDA	C27-O28-P29-O30
2	В	400	XDA	O24-C23-N25-C26
2	В	400	XDA	C20-C23-N25-C26
2	A	400	XDA	O24-C23-N25-C26
2	A	400	XDA	C20-C23-N25-C26
2	В	400	XDA	C27-O28-P29-O32
2	A	400	XDA	C19-C20-C23-N25
2	В	400	XDA	N12-C13-O15-C16
2	A	400	XDA	C21-C20-C23-N25
2	A	400	XDA	C19-C20-C23-O24
2	A	400	XDA	C21-C20-C23-O24
2	В	400	XDA	O14-C13-O15-C16
2	A	400	XDA	C5-C4-C7-N9
2	В	400	XDA	N25-C26-C27-O28

There are no ring outliers.


2 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	400	XDA	4	0
2	В	400	XDA	15	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will



also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(Å^2)$	Q < 0.9
1	A	319/337 (94%)	0.20	10 (3%) 49 44	51, 76, 107, 131	0
1	В	326/337~(96%)	0.20	5 (1%) 73 73	49, 76, 101, 123	0
All	All	645/674 (95%)	0.20	15 (2%) 60 58	49, 76, 104, 131	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	2	GLU	4.3
1	A	80	ASN	4.1
1	A	66	LEU	2.7
1	В	43	ILE	2.5
1	A	81	GLY	2.4
1	A	28	ASN	2.4
1	В	45	ALA	2.3
1	В	276	ILE	2.2
1	A	112	THR	2.1
1	A	107	LEU	2.1
1	A	79	VAL	2.1
1	A	54	ALA	2.1
1	A	94[A]	ASP	2.0
1	В	329	SER	2.0
1	A	30	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	XDA	В	400	32/32	0.78	0.40	48,54,65,65	58
2	XDA	A	400	32/32	0.88	0.32	57,61,64,65	58

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

