



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2022 – 02:01 pm BST

PDB ID : 7O5R
Title : Crystal structure of holo-SwHPA-Mn (hydroxyketoacid aldolase) from *Sphingomonas wittichii* RW1
Authors : Laustsen, J.; Justo, I.; Marsden, S.R.; Hanefeld, U.; Bento, I.
Deposited on : 2021-04-09
Resolution : 1.65 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.31.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	
1	B	252	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8144 atoms, of which 3839 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hydroxyketoacid aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	251	3782	1190	1894	336	354	8	103	3	0
1	B	252	3877	1219	1945	343	362	8	109	9	0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	2
			2	2		
2	B	1	Total	Mn	0	0
			1	1		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		
3	B	2	Total	K	0	0
			2	2		

- Molecule 4 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Br	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	230	Total	O	0	9
			239	239		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	234	Total 239	O 239	0	5



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: hydroxyketoacid aldolase



- Molecule 1: hydroxyketoacid aldolase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	71.54Å 71.54Å 225.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	75.13 – 1.65 75.13 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.9 (75.13-1.65) 100.0 (75.13-1.65)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.160 , 0.193 0.170 , 0.202	Depositor DCC
R_{free} test set	2506 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	13.5	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.008 for -h-k,k,-l	Xtriage
Reported twinning fraction	0.756 for H, K, L 0.244 for K, H, -L	Depositor
Outliers	0 of 51788 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8144	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, MN, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.63	1/1927 (0.1%)	0.95	0/2621
1	B	0.59	0/1987	0.98	2/2702 (0.1%)
All	All	0.61	1/3914 (0.0%)	0.97	2/5323 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	145	GLU	CD-OE1	7.48	1.33	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	97	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	B	69	ARG	NE-CZ-NH1	5.40	123.00	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	147	ARG	Sidechain
1	A	199	ARG	Sidechain
1	B	112	ARG	Sidechain
1	B	125	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1888	1894	1888	20	0
1	B	1932	1945	1941	23	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
4	A	1	0	0	1	0
5	A	239	0	0	5	1
5	B	239	0	0	5	2
All	All	4305	3839	3829	43	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:97:ARG:NH1	1:A:101:GLN:NE2	2.28	0.82
1:B:15:LEU:HD12	1:B:228[B]:VAL:HG23	1.67	0.74
1:A:102:THR:HG21	5:A:564:HOH:O	1.91	0.69
1:A:93:MET:HE3	1:A:95:ASN:HD21	1.58	0.68
1:A:97:ARG:NH1	1:A:101:GLN:HE21	1.93	0.67
1:B:101:GLN:NE2	5:B:405:HOH:O	2.29	0.65
1:A:97:ARG:HH11	1:A:101:GLN:NE2	1.95	0.64
1:B:198:GLU:OE2	1:B:199:ARG:HD2	1.98	0.64
1:A:21:LEU:HB3	5:A:588:HOH:O	2.01	0.61
1:A:97:ARG:HH12	1:A:101:GLN:HE21	1.47	0.61
1:B:197[B]:ARG:HG3	1:B:207:VAL:HG11	1.82	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:THR:HG21	1:A:182[A]:VAL:HG12	1.84	0.59
1:B:75:PRO:O	1:B:79:MET:HG3	2.04	0.58
1:B:170[A]:THR:HG22	1:B:181:LEU:O	2.03	0.57
1:A:208:GLY:HA2	1:A:227:PHE:O	2.06	0.54
1:A:101:GLN:HG2	5:A:410[B]:HOH:O	2.06	0.54
1:B:3:LYS:NZ	1:B:63:GLY:O	2.27	0.53
1:B:209:ILE:HG21	1:B:220:LYS:HG3	1.92	0.52
1:A:170:THR:HG21	1:A:182[B]:VAL:HG22	1.93	0.51
1:B:197[B]:ARG:CG	1:B:207:VAL:HG11	2.40	0.50
1:B:5:ARG:HD3	1:B:136:GLU:OE1	2.12	0.50
1:B:211:CYS:HB2	1:B:229:THR:O	2.12	0.50
1:B:132:ALA:O	5:B:403:HOH:O	2.20	0.49
1:A:93:MET:CE	1:A:95:ASN:HD21	2.24	0.48
1:B:197[A]:ARG:CZ	5:B:471:HOH:O	2.62	0.48
1:A:170:THR:CG2	1:A:182[A]:VAL:HG12	2.43	0.48
1:B:208:GLY:HA2	1:B:227:PHE:O	2.14	0.47
1:B:32:ARG:HA	5:B:417:HOH:O	2.15	0.47
1:A:207:VAL:N	5:A:412:HOH:O	2.48	0.46
1:A:48:ASP:OD1	1:A:51:GLN:HG3	2.16	0.46
1:A:182[B]:VAL:HG21	4:A:304:BR:BR	2.71	0.45
1:B:16:ALA:HB3	1:B:229:THR:HG22	1.99	0.45
1:B:119:PRO:HG2	1:B:123[B]:SER:OG	2.18	0.44
1:B:53:ALA:N	1:B:54:PRO:HD2	2.33	0.44
1:A:63:GLY:HA2	5:A:507[A]:HOH:O	2.18	0.44
1:A:51:GLN:O	1:A:55:MET:HG3	2.18	0.43
1:B:201:HIS:HE1	1:B:226:ASP:OD2	2.01	0.42
1:B:150[B]:LEU:HD11	1:B:192:ALA:HB1	2.02	0.42
1:A:150:LEU:HD12	1:A:150:LEU:HA	1.90	0.41
1:B:32:ARG:C	5:B:417:HOH:O	2.58	0.41
1:A:40:ILE:HG21	1:A:52:VAL:HG23	2.01	0.41
1:B:49:PHE:O	1:B:52:VAL:HG12	2.20	0.41
1:B:48:ASP:OD1	1:B:51:GLN:HG3	2.22	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:551:HOH:O	5:B:621:HOH:O[3_665]	2.03	0.17
5:B:404:HOH:O	5:B:449:HOH:O[2_555]	2.12	0.08
5:A:587:HOH:O	5:A:603:HOH:O[2_655]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/252 (100%)	244 (97%)	8 (3%)	0	100	100
1	B	259/252 (103%)	250 (96%)	8 (3%)	1 (0%)	34	16
All	All	511/504 (101%)	494 (97%)	16 (3%)	1 (0%)	47	28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	118	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/189 (101%)	190 (100%)	1 (0%)	88	81
1	B	198/189 (105%)	197 (100%)	1 (0%)	88	81
All	All	389/378 (103%)	387 (100%)	2 (0%)	88	81

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	LEU
1	B	114	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	GLN
1	B	201	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/252 (99%)	0.62	5 (1%) 65 67	7, 12, 21, 47	0
1	B	252/252 (100%)	0.69	9 (3%) 42 43	7, 12, 22, 39	0
All	All	503/504 (99%)	0.65	14 (2%) 53 53	7, 12, 22, 47	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	251	LEU	6.4
1	B	1	HIS	4.4
1	B	251	LEU	3.6
1	B	132	ALA	3.4
1	B	11	GLY	3.1
1	A	182[A]	VAL	2.7
1	B	111	PRO	2.7
1	B	227	PHE	2.4
1	B	128	SER	2.3
1	B	131	LEU	2.2
1	B	8	TRP	2.1
1	A	84	ALA	2.1
1	A	138	VAL	2.1
1	A	181	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MN	A	302[B]	1/1	0.80	0.12	13,13,13,13	1
2	MN	A	301[A]	1/1	0.98	0.08	11,11,11,11	1
2	MN	B	301	1/1	0.99	0.09	11,11,11,11	1
4	BR	A	304	1/1	0.99	0.05	18,18,18,18	1
3	K	B	302	1/1	1.00	0.11	12,12,12,12	0
3	K	B	303	1/1	1.00	0.10	19,19,19,19	1
3	K	A	303	1/1	1.00	0.09	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.