



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 3, 2023 – 03:59 pm GMT

PDB ID : 1OC2
Title : The structure of NADH in the dTDP-D-glucose dehydratase (RmlB) enzyme
Authors : Beis, K.; Naismith, J.H.
Deposited on : 2003-02-04
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	
1	B	348	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	1352	-	-	X	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6465 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DTDP-GLUCOSE 4,6-DEHYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	346	Total	C	N	O	S	0	13	0
			2787	1760	474	550	3			
1	B	347	Total	C	N	O	S	0	4	1
			2750	1737	471	539	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	313	GLU	GLN	conflict	UNP Q8GIP9
B	313	GLU	GLN	conflict	UNP Q8GIP9

- Molecule 2 is THYMIDINE-5'-DIPHOSPHO-BETA-D-XYLOSE (three-letter code: TDX) (formula: C₁₅H₂₄N₂O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			34	15	2	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	B	1	34	15	2	15	2	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	44	21	7	14	2	0	0
3	B	1	44	21	7	14	2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	404	Total O 404 404	0	0
5	B	353	Total O 353 353	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DTDP-GLUCOSE 4,6-DEHYDRATASE



- Molecule 1: DTDP-GLUCOSE 4,6-DEHYDRATASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.41Å 94.83Å 183.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	91.29 – 1.50 34.74 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.3 (91.29-1.50) 98.3 (34.74-1.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.170 , 0.200 0.183 , 0.206	Depositor DCC
R_{free} test set	16689 reflections (9.89%)	wwPDB-VP
Wilson B-factor (Å ²)	15.1	Xtrriage
Anisotropy	0.075	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 54.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6465	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, SO4, TDX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.06	9/2924 (0.3%)	1.01	17/3973 (0.4%)
1	B	0.90	2/2838 (0.1%)	0.90	9/3860 (0.2%)
All	All	0.98	11/5762 (0.2%)	0.96	26/7833 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	49[A]	GLU	CD-OE2	16.95	1.44	1.25
1	A	49[B]	GLU	CD-OE2	16.95	1.44	1.25
1	B	131[A]	ASP	CB-CG	-10.24	1.30	1.51
1	B	131[B]	ASP	CB-CG	-10.24	1.30	1.51
1	A	163[A]	SER	CB-OG	-9.14	1.30	1.42
1	A	163[B]	SER	CB-OG	-9.14	1.30	1.42
1	A	131[A]	ASP	CB-CG	-7.90	1.35	1.51
1	A	131[B]	ASP	CB-CG	-7.90	1.35	1.51
1	A	131[A]	ASP	CA-CB	6.10	1.67	1.53
1	A	131[B]	ASP	CA-CB	6.10	1.67	1.53
1	A	220	GLU	CD-OE2	5.04	1.31	1.25

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	49[A]	GLU	OE1-CD-OE2	-10.90	110.22	123.30
1	A	49[B]	GLU	OE1-CD-OE2	-10.90	110.22	123.30
1	A	276	ASP	CB-CG-OD2	9.26	126.64	118.30
1	B	276	ASP	CB-CG-OD2	8.50	125.95	118.30
1	A	54	ASP	CB-CG-OD2	8.04	125.54	118.30
1	A	300	ASP	CB-CG-OD2	7.97	125.48	118.30
1	A	126	ASP	CB-CG-OD1	7.32	124.89	118.30
1	A	3	GLN	CA-CB-CG	7.24	129.33	113.40
1	B	310	ASP	CB-CG-OD2	7.18	124.76	118.30
1	A	310	ASP	CB-CG-OD2	6.96	124.56	118.30
1	B	245	ARG	NE-CZ-NH1	6.91	123.75	120.30
1	B	300	ASP	CB-CG-OD2	6.83	124.45	118.30
1	B	327	ASP	CB-CG-OD2	6.38	124.04	118.30
1	A	131[A]	ASP	CB-CA-C	-6.03	98.34	110.40
1	A	131[B]	ASP	CB-CA-C	-6.03	98.34	110.40
1	A	131[A]	ASP	CB-CG-OD2	5.87	123.58	118.30
1	A	131[B]	ASP	CB-CG-OD2	5.87	123.58	118.30
1	B	30	ASP	CB-CG-OD1	5.81	123.53	118.30
1	B	291	TYR	CB-CG-CD2	-5.62	117.63	121.00
1	A	77	ASP	CB-CG-OD2	5.59	123.33	118.30
1	A	70	ASP	CB-CG-OD2	5.53	123.28	118.30
1	A	135	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	327	ASP	CB-CG-OD2	5.38	123.15	118.30
1	B	126	ASP	CB-CG-OD1	5.21	122.99	118.30
1	B	245	ARG	NE-CZ-NH2	-5.16	117.72	120.30
1	A	37	ASP	CB-CG-OD2	5.14	122.93	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	49[A]	GLU	Sidechain
1	A	49[B]	GLU	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2787	0	2655	32	0
1	B	2750	0	2627	44	0
2	A	34	0	22	3	0
2	B	34	0	22	2	0
3	A	44	0	26	4	0
3	B	44	0	26	4	0
4	A	15	0	0	4	0
5	A	404	0	0	16	0
5	B	353	0	0	19	0
All	All	6465	0	5378	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:19[B]:ASN:ND2	1:B:230:THR:HG23	1.58	1.18
1:A:265[A]:GLU:OE2	5:A:2278:HOH:O	1.60	1.18
1:A:265[A]:GLU:CD	5:A:2278:HOH:O	1.82	1.16
4:A:1352:SO4:O1	5:A:2398:HOH:O	1.71	1.07
1:A:131[B]:ASP:OD1	5:A:2169:HOH:O	1.71	1.07
4:A:1352:SO4:S	5:A:2398:HOH:O	2.17	1.02
1:B:154[B]:ASN:OD1	5:B:2196:HOH:O	1.80	0.97
1:B:19[B]:ASN:HD22	1:B:230:THR:HG23	1.14	0.97
4:A:1352:SO4:O1	5:A:2401:HOH:O	1.82	0.96
1:A:131[B]:ASP:OD2	5:A:2170:HOH:O	1.84	0.95
1:B:154[B]:ASN:OD1	5:B:2194:HOH:O	1.89	0.90
4:A:1352:SO4:S	5:A:2401:HOH:O	2.32	0.88
1:B:19[B]:ASN:HD22	1:B:230:THR:CG2	1.87	0.85
1:B:2:SER:OG	5:B:2001:HOH:O	1.95	0.84
1:B:19[B]:ASN:ND2	1:B:230:THR:CG2	2.42	0.82
1:B:19[B]:ASN:HD21	1:B:231:ASN:HD22	1.28	0.81
1:B:19[B]:ASN:CG	5:B:2015:HOH:O	2.20	0.81
1:B:19[A]:ASN:HB3	5:B:2015:HOH:O	1.82	0.79
1:B:229:HIS:HD2	1:B:231:ASN:H	1.31	0.78
1:B:22:HIS:HE1	1:B:47:ASN:O	1.68	0.77
1:B:2:SER:HA	5:B:2002:HOH:O	1.85	0.76
1:B:19[B]:ASN:ND2	5:B:2015:HOH:O	2.21	0.73
1:B:131[B]:ASP:OD1	5:B:2165:HOH:O	2.07	0.73
1:B:19[B]:ASN:HD21	1:B:231:ASN:ND2	1.88	0.72
1:A:131[B]:ASP:CG	5:A:2169:HOH:O	2.24	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:327:ASP:HB2	5:A:2352:HOH:O	1.98	0.64
1:B:3:GLN:NE2	1:B:242:THR:O	2.31	0.63
2:A:1349:TDX:H5D	5:A:2211:HOH:O	1.98	0.61
1:B:2:SER:CA	5:B:2002:HOH:O	2.45	0.60
3:B:1348:NAD:H4N	5:B:2218:HOH:O	2.02	0.59
1:A:81[A]:HIS:CE1	1:A:109:LEU:HD12	2.38	0.58
1:A:154:ASN:HA	1:B:154[B]:ASN:OD1	2.03	0.58
1:A:188:CYS:HB3	3:A:1350:NAD:H5N	1.84	0.57
1:B:283:ASP:OD1	1:B:287:HIS:HD2	1.87	0.57
1:B:131[B]:ASP:CG	5:B:2165:HOH:O	2.42	0.57
1:A:197:HIS:HD2	1:A:199:GLU:H	1.54	0.56
1:B:188:CYS:HB3	3:B:1348:NAD:H5N	1.88	0.56
1:A:16:ILE:HD11	3:A:1350:NAD:C7N	2.37	0.55
1:B:313:GLU:HG3	5:B:2312:HOH:O	2.06	0.54
1:A:173:LYS:HE2	1:A:185:ILE:HD12	1.89	0.54
1:B:131[A]:ASP:CG	1:B:289:LEU:H	2.12	0.53
1:B:173:LYS:HE2	1:B:185:ILE:HD12	1.92	0.51
1:A:81[A]:HIS:ND1	1:A:109:LEU:CD1	2.74	0.51
1:A:305:THR:HG23	5:A:2329:HOH:O	2.10	0.51
1:A:81[A]:HIS:CE1	1:A:109:LEU:CD1	2.94	0.50
3:B:1348:NAD:C3N	2:B:1349:TDX:H4'	2.42	0.50
1:B:155:TYR:OH	5:B:2199:HOH:O	2.18	0.50
1:B:122:HIS:HE1	1:B:169:ASP:OD1	1.95	0.50
1:A:127:GLU:OE1	2:A:1349:TDX:H5B	2.11	0.49
1:B:62:ASP:OD2	1:B:100:HIS:HE1	1.96	0.49
1:A:23:TYR:OH	1:A:28:HIS:HE1	1.95	0.49
1:B:131[A]:ASP:OD1	1:B:289:LEU:N	2.47	0.48
1:A:265[B]:GLU:HG2	5:A:2278:HOH:O	2.14	0.47
1:A:259:ASN:O	1:A:262[A]:GLU:HG2	2.15	0.47
1:B:122:HIS:CE1	1:B:169:ASP:OD1	2.67	0.47
1:A:275:LYS:O	5:A:2278:HOH:O	2.21	0.47
1:A:62:ASP:OD2	1:A:100:HIS:HE1	1.98	0.46
1:A:347:ILE:O	5:A:2385:HOH:O	2.21	0.46
1:B:173:LYS:CE	1:B:185:ILE:HD12	2.46	0.46
1:B:120:PHE:O	1:B:183:ALA:HA	2.17	0.45
1:A:32:HIS:HE1	5:A:2068:HOH:O	1.99	0.45
1:A:197:HIS:CD2	1:A:199:GLU:H	2.35	0.45
2:B:1349:TDX:H5D	5:B:2218:HOH:O	2.18	0.44
1:A:159:SER:O	1:A:163[A]:SER:HB2	2.18	0.44
1:B:19[B]:ASN:ND2	1:B:231:ASN:HD22	2.06	0.44
1:A:120:PHE:O	1:A:183:ALA:HA	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16:ILE:HD11	3:B:1348:NAD:C7N	2.48	0.43
2:A:1349:TDX:H4'	3:A:1350:NAD:C3N	2.48	0.43
1:A:16:ILE:HD11	3:A:1350:NAD:N7N	2.34	0.43
1:A:173:LYS:CE	1:A:185:ILE:HD12	2.49	0.43
1:A:96[B]:SER:OG	1:A:97:PRO:HD3	2.19	0.42
1:B:32:HIS:HE1	5:B:2073:HOH:O	2.02	0.42
1:A:201:PHE:O	1:A:205:GLN:HG2	2.20	0.41
1:B:81:HIS:HD2	1:B:122:HIS:HD2	1.68	0.41
1:B:258:LYS:HD3	5:B:2269:HOH:O	2.20	0.41
1:B:28:HIS:CE1	1:B:238:TRP:CE2	3.08	0.41
1:B:28:HIS:HD2	5:B:2001:HOH:O	2.03	0.41
1:B:186:SER:HA	1:B:250:TYR:O	2.21	0.41
1:B:16:ILE:HG22	1:B:82:TYR:CE2	2.56	0.40
1:B:263:VAL:HA	5:B:2269:HOH:O	2.21	0.40
1:A:163[A]:SER:HB3	1:B:171:ILE:HG22	2.02	0.40
1:A:191:ASN:HA	1:A:228:ILE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/348 (103%)	348 (98%)	9 (2%)	0	100	100
1	B	349/348 (100%)	341 (98%)	8 (2%)	0	100	100
All	All	706/696 (101%)	689 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	302/291 (104%)	290 (96%)	12 (4%)	31 6
1	B	293/291 (101%)	286 (98%)	7 (2%)	49 19
All	All	595/582 (102%)	576 (97%)	19 (3%)	49 10

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49[A]	GLU
1	A	49[B]	GLU
1	A	131[A]	ASP
1	A	131[B]	ASP
1	A	139	PRO
1	A	143	GLU
1	A	152[A]	GLU
1	A	152[B]	GLU
1	A	163[A]	SER
1	A	163[B]	SER
1	A	191	ASN
1	A	245	ARG
1	B	3	GLN
1	B	131[A]	ASP
1	B	131[B]	ASP
1	B	191	ASN
1	B	246	MET
1	B	287	HIS
1	B	332	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	A	32	HIS
1	A	100	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	122	HIS
1	A	141	HIS
1	A	196	GLN
1	A	197	HIS
1	A	231	ASN
1	A	260	ASN
1	A	280	HIS
1	A	307	GLN
1	A	339	ASN
1	B	22	HIS
1	B	28	HIS
1	B	32	HIS
1	B	87	HIS
1	B	100	HIS
1	B	122	HIS
1	B	191	ASN
1	B	196	GLN
1	B	197	HIS
1	B	229	HIS
1	B	233	HIS
1	B	260	ASN
1	B	273	GLN
1	B	280	HIS
1	B	287	HIS
1	B	339	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	1351	-	4,4,4	0.42	0	6,6,6	0.72	0
4	SO4	A	1353	-	4,4,4	0.48	0	6,6,6	0.71	0
3	NAD	B	1348	-	42,48,48	2.40	11 (26%)	50,73,73	3.30	10 (20%)
3	NAD	A	1350	-	42,48,48	2.11	7 (16%)	50,73,73	2.87	13 (26%)
2	TDX	A	1349	-	33,36,36	1.35	5 (15%)	50,55,55	1.45	8 (16%)
4	SO4	A	1352	-	4,4,4	0.19	0	6,6,6	0.36	0
2	TDX	B	1349	-	33,36,36	1.07	3 (9%)	50,55,55	1.82	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TDX	A	1349	-	-	1/21/50/50	0/3/3/3
3	NAD	B	1348	-	-	6/26/62/62	0/5/5/5
2	TDX	B	1349	-	-	1/21/50/50	0/3/3/3
3	NAD	A	1350	-	-	4/26/62/62	0/5/5/5

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1348	NAD	C4N-C3N	9.59	1.55	1.39
3	A	1350	NAD	C4N-C3N	9.02	1.54	1.39
3	B	1348	NAD	C5N-C4N	5.37	1.50	1.38
3	B	1348	NAD	C2N-N1N	5.11	1.41	1.35
3	A	1350	NAD	C2N-N1N	4.78	1.40	1.35
3	A	1350	NAD	C5N-C4N	4.49	1.48	1.38
3	B	1348	NAD	C6N-N1N	4.28	1.45	1.35
2	A	1349	TDX	C4-N3	-3.78	1.31	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1350	NAD	C6N-N1N	3.75	1.44	1.35
3	A	1350	NAD	C4A-N3A	3.51	1.40	1.35
2	B	1349	TDX	C4-N3	-2.99	1.33	1.38
3	B	1348	NAD	C2B-C1B	-2.91	1.49	1.53
2	A	1349	TDX	C2-N1	-2.90	1.33	1.38
2	A	1349	TDX	O2-C2	2.85	1.28	1.23
2	A	1349	TDX	C2-N3	-2.81	1.33	1.38
3	B	1348	NAD	C4A-N3A	2.81	1.39	1.35
3	B	1348	NAD	C7N-N7N	2.77	1.38	1.33
3	B	1348	NAD	O4D-C4D	-2.70	1.39	1.45
2	B	1349	TDX	O2-C2	2.65	1.27	1.23
3	B	1348	NAD	C6N-C5N	-2.59	1.32	1.38
3	B	1348	NAD	O3D-C3D	2.43	1.48	1.43
2	B	1349	TDX	C6-N1	-2.43	1.33	1.38
3	A	1350	NAD	C6N-C5N	-2.20	1.33	1.38
2	A	1349	TDX	C4'-C3'	-2.19	1.49	1.52
3	B	1348	NAD	O4B-C1B	2.15	1.44	1.41
3	A	1350	NAD	C2A-N3A	2.13	1.35	1.32

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1348	NAD	C5N-C4N-C3N	-13.81	104.00	120.34
3	B	1348	NAD	C6N-N1N-C2N	-12.86	110.25	121.97
3	A	1350	NAD	C5N-C4N-C3N	-11.28	107.00	120.34
2	B	1349	TDX	O4-C4-C5	-7.80	115.86	124.90
3	A	1350	NAD	O7N-C7N-C3N	-7.69	110.42	119.63
3	B	1348	NAD	C3N-C7N-N7N	7.67	126.96	117.75
3	B	1348	NAD	O7N-C7N-C3N	-7.39	110.78	119.63
3	A	1350	NAD	C6N-N1N-C2N	-7.00	115.59	121.97
3	A	1350	NAD	C2N-C3N-C4N	6.50	125.62	118.26
2	B	1349	TDX	C5-C4-N3	5.16	119.71	115.31
3	A	1350	NAD	C3N-C7N-N7N	4.22	122.82	117.75
3	A	1350	NAD	C3N-C2N-N1N	-4.08	116.44	120.43
2	B	1349	TDX	C4-N3-C2	-3.92	122.27	127.35
3	B	1348	NAD	N3A-C2A-N1A	-3.85	122.67	128.68
3	A	1350	NAD	N3A-C2A-N1A	-3.78	122.78	128.68
3	A	1350	NAD	C2A-N1A-C6A	3.68	125.05	118.75
2	A	1349	TDX	N3-C2-N1	3.56	119.62	114.89
2	B	1349	TDX	C6-C5-C4	-3.10	115.44	118.03
3	A	1350	NAD	N6A-C6A-N1A	3.05	124.91	118.57
3	A	1350	NAD	C3D-C2D-C1D	-3.05	96.39	100.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1349	TDX	C4'-C3'-C2'	-2.92	105.84	110.89
2	A	1349	TDX	C4-N3-C2	-2.90	123.60	127.35
2	A	1349	TDX	C5-C4-N3	2.86	117.76	115.31
3	B	1348	NAD	C5A-C6A-N6A	-2.70	116.25	120.35
2	B	1349	TDX	N3-C2-N1	2.64	118.39	114.89
3	A	1350	NAD	O7N-C7N-N7N	2.63	126.31	122.58
2	A	1349	TDX	C5-C6-N1	-2.60	120.67	123.34
2	A	1349	TDX	C2B-C1B-N1	2.59	119.74	113.77
2	A	1349	TDX	O4'-C4'-C5'	-2.54	103.95	109.15
3	A	1350	NAD	C6N-C5N-C4N	2.37	122.88	119.44
2	B	1349	TDX	O4B-C1B-N1	-2.32	103.72	107.86
3	B	1348	NAD	O4B-C1B-C2B	-2.26	103.62	106.93
2	A	1349	TDX	O4-C4-C5	-2.26	122.28	124.90
3	B	1348	NAD	C6N-C5N-C4N	2.21	122.66	119.44
3	B	1348	NAD	C2N-C3N-C4N	2.15	120.70	118.26
3	B	1348	NAD	C4A-C5A-N7A	-2.15	107.16	109.40
2	B	1349	TDX	O4-C4-N3	2.14	124.23	120.12
3	A	1350	NAD	O2N-PN-O1N	2.02	122.23	112.24

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1350	NAD	O4D-C1D-N1N-C2N
3	A	1350	NAD	O4D-C1D-N1N-C6N
3	A	1350	NAD	C2D-C1D-N1N-C6N
3	B	1348	NAD	C5D-O5D-PN-O1N
3	B	1348	NAD	O4D-C1D-N1N-C6N
3	B	1348	NAD	C2D-C1D-N1N-C6N
2	B	1349	TDX	C1'-O1'-PB-O3A
3	B	1348	NAD	C5D-O5D-PN-O3
3	B	1348	NAD	C5D-O5D-PN-O2N
2	A	1349	TDX	C1'-O1'-PB-O3A
3	B	1348	NAD	O4B-C4B-C5B-O5B
3	A	1350	NAD	O4B-C4B-C5B-O5B

There are no ring outliers.

5 monomers are involved in 15 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1348	NAD	4	0



Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1350	NAD	4	0
2	A	1349	TDX	3	0
4	A	1352	SO4	4	0
2	B	1349	TDX	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	346/348 (99%)	0.42	16 (4%) 32 35	10, 14, 23, 35	5 (1%)
1	B	347/348 (99%)	0.83	42 (12%) 4 4	12, 18, 28, 39	9 (2%)
All	All	693/696 (99%)	0.63	58 (8%) 11 11	10, 16, 27, 39	14 (2%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	121	HIS	7.7
1	B	272	GLY	6.9
1	B	85	GLU	5.4
1	A	145	PRO	5.3
1	B	141	HIS	5.2
1	B	275	LYS	5.1
1	A	141	HIS	4.3
1	B	334	GLU	4.2
1	A	139	PRO	3.9
1	B	2	SER	3.9
1	B	347	ILE	3.7
1	B	278	TYR	3.6
1	B	276	ASP	3.5
1	A	131[A]	ASP	3.4
1	B	123	VAL	3.3
1	A	275	LYS	3.2
1	B	332	GLU	3.1
1	A	332	GLU	3.0
1	B	262	GLU	3.0
1	B	331	ALA	3.0
1	B	327	ASP	2.9
1	B	16	ILE	2.9
1	B	274	PRO	2.8
1	B	265	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	80	VAL	2.7
1	B	313	GLU	2.6
1	B	145	PRO	2.6
1	B	82	TYR	2.6
1	B	341	ALA	2.6
1	A	15	PHE	2.6
1	A	152[A]	GLU	2.6
1	B	273	GLN	2.5
1	A	276	ASP	2.5
1	B	348	LYS	2.5
1	B	269	GLU	2.4
1	B	308	PHE	2.4
1	A	123	VAL	2.4
1	B	222	LYS	2.4
1	A	143	GLU	2.4
1	B	248	GLU	2.3
1	B	209	ILE	2.3
1	A	151	ALA	2.3
1	B	210	LEU	2.3
1	A	144	GLY	2.2
1	B	220	GLU	2.2
1	B	152	GLU	2.2
1	B	345	GLU	2.2
1	B	172	VAL	2.2
1	A	54	ASP	2.1
1	B	320	GLN	2.1
1	B	309	THR	2.1
1	A	49[A]	GLU	2.1
1	B	211	ALA	2.1
1	B	300	ASP	2.1
1	A	269	GLU	2.1
1	B	346	VAL	2.0
1	B	324	ASP	2.0
1	B	131[A]	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	A	1352	5/5	0.74	0.80	90,90,90,90	0
2	TDX	B	1349	34/34	0.95	0.12	18,20,22,25	0
3	NAD	A	1350	44/44	0.96	0.15	12,14,25,26	0
3	NAD	B	1348	44/44	0.96	0.13	16,18,31,32	0
2	TDX	A	1349	34/34	0.96	0.15	12,14,17,19	0
4	SO4	A	1351	5/5	0.97	0.15	26,26,30,30	0
4	SO4	A	1353	5/5	0.98	0.17	29,31,33,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around TDX B 1349:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD A 1350:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD B 1348:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around TDX A 1349:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.