Full wwPDB X-ray Structure Validation Report (i)

Oct 28, 2021 – 02:03 pm BST

PDB ID	:	7OJT
Title	:	Crystal structure of unliganded PatA, a membrane associated acyltransferase from Mycobacterium smegmatis
Authors	:	Anso, I.; Wang, L.; Marina, A.; Paez-Perez, E.D.; Perrone, S.; Lowary, T.L.; Trastoy, B.; Guerin, M.E.
Deposited on	:	2021-05-17
Resolution	:	3.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1013 (3.84-3.52)
Clashscore	141614	1070 (3.84-3.52)
Ramachandran outliers	138981	1036 (3.84-3.52)
Sidechain outliers	138945	1033 (3.84-3.52)
RSRZ outliers	127900	1471 (3.86-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	70% 10% 20%
1	B	313	67% 12% 20%
1	C	313	.% 73% 7% 20%
1	D	313	.% 72% 8% 20%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 7134 atoms, of which 8 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol mannoside acyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	249	1787	1140	311	330	6	0	0	0
1	B	249	1793	1141	308	336	8	0	0	0
1	C	249	1724	1086	302	330	6	0	0	0
1	D	249	1816	1159	314	335	8	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP A0QWG5
A	1	VAL	-	expression tag	UNP A0QWG5
A	305	SER	-	expression tag	UNP A0QWG5
A	306	ARG	-	expression tag	UNP A0QWG5
A	307	HIS	-	expression tag	UNP A0QWG5
A	308	HIS	-	expression tag	UNP A0QWG5
A	309	HIS	-	expression tag	UNP A0QWG5
A	310	HIS	-	expression tag	UNP A0QWG5
A	311	HIS	-	expression tag	UNP A0QWG5
A	312	HIS	-	expression tag	UNP A0QWG5
B	0	MET	-	initiating methionine	UNP A0QWG5
B	1	VAL	-	expression tag	UNP A0QWG5
B	305	SER	-	expression tag	UNP A0QWG5
B	306	ARG	-	expression tag	UNP A0QWG5
B	307	HIS	-	expression tag	UNP A0QWG5
B	308	HIS	-	expression tag	UNP A0QWG5
B	309	HIS	-	expression tag	UNP A0QWG5
B	310	HIS	-	expression tag	UNP A0QWG5
B	311	HIS	-	expression tag	UNP A0QWG5
B	312	HIS	-	expression tag	UNP A0QWG5
C	0	MET	-	initiating methionine	UNP A0QWG5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	VAL	-	expression tag	UNP A0QWG5
C	305	SER	-	expression tag	UNP A0QWG5
C	306	ARG	-	expression tag	UNP A0QWG5
C	307	HIS	-	expression tag	UNP A0QWG5
C	308	HIS	-	expression tag	UNP A0QWG5
C	309	HIS	-	expression tag	UNP A0QWG5
C	310	HIS	-	expression tag	UNP A0QWG5
C	311	HIS	-	expression tag	UNP A0QWG5
C	312	HIS	-	expression tag	UNP A0QWG5
D	0	MET	-	initiating methionine	UNP A0QWG5
D	1	VAL	-	expression tag	UNP A0QWG5
D	305	SER	-	expression tag	UNP A0QWG5
D	306	ARG	-	expression tag	UNP A0QWG5
D	307	HIS	-	expression tag	UNP A0QWG5
D	308	HIS	-	expression tag	UNP A0QWG5
D	309	HIS	-	expression tag	UNP A0QWG5
D	310	HIS	-	expression tag	UNP A0QWG5
D	311	HIS	-	expression tag	UNP A0QWG5
D	312	HIS	-	expression tag	UNP A0QWG5

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	14	3	8	3	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphatidylinositol mannoside acyltransferase

Chain A:



• Molecule 1: Phosphatidylinositol mannoside acyltransferase

Chain B:



• Molecule 1: Phosphatidylinositol mannoside acyltransferase

Chain C:



• Molecule 1: Phosphatidylinositol mannoside acyltransferase

Chain D:

%
72%
8%
20%


MET VAL THR LEU SER GLY ARG ILE PRO LEU GLY GLY GLN VAL THR ASP LEU GLY TYR ALA ALA GLY TRP ARG LEU VAL ARG ALA MET PRO GLU ALA MET ALA GLN GLY VAL PHE GLY ALA GLY ALA ARG TYR ALA ALA ARG ASN GLY GLY
P50
N56 L57
V60 V61
A116

W130
G134
E149
F169
L174
P180 P181
F182 E183
A199
V208
F212
E215
A221
A233 L234 F235
W240
A273
D282 W283 H284 M285 L286 Q287 P288 Q289 W290
R298
ARG ALA ARG LEU GLY THR SER ARG HIS HIS HIS HIS HIS HIS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	81.31Å 92.97Å 81.09Å 90.00° 90.33° 90.00°	Depositor
Resolution (Å)	46.48 – 3.67 48.75 – 3.67	Depositor EDS
% Data completeness (in resolution range)	99.0 (46.48-3.67) 99.5 (48.75-3.67)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.79 (at 3.67Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.268 , 0.290 0.268 , 0.294	Depositor DCC
R_{free} test set	664 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å^2)	126.7	Xtriage
Anisotropy	0.593	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	(Not available) , (Not available)	EDS
L-test for twinning[2]	$< \|L\| > = 0.42$, $< L^2 > = 0.25$	Xtriage
Estimated twinning fraction	0.054 for -l,k,h 0.177 for -h,-k,l 0.057 for l,-k,h	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	7134	wwPDB-VP
Average B, all atoms (Å^2)	124.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.14% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.29	0/1840	0.53	0/2533
1	B	0.28	0/1846	0.51	0/2540
1	C	0.28	0/1770	0.52	0/2440
1	D	0.28	0/1869	0.52	0/2567
All	All	0.28	0/7325	0.52	0/10080

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1787	0	1586	23	0
1	B	1793	0	1586	23	0
1	C	1724	0	1472	16	0
1	D	1816	0	1638	19	0
2	A	6	8	8	0	0
All	All	7126	8	6290	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:285:MET:HE2	1:C:287:GLN:H	1.33	0.94
1:B:285:MET:HE2	1:B:287:GLN:H	1.38	0.89
1:D:285:MET:HE2	1:D:287:GLN:H	1.43	0.84
1:A:285:MET:HE2	1:A:287:GLN:H	1.45	0.82
1:C:285:MET:CE	1:C:287:GLN:H	1.96	0.78
1:C:285:MET:HE2	1:C:287:GLN:N	2.01	0.73
1:B:285:MET:CE	1:B:287:GLN:H	2.02	0.72
1:A:285:MET:CE	1:A:287:GLN:H	2.04	0.71
1:D:285:MET:CE	1:D:287:GLN:H	2.05	0.70
1:B:285:MET:HE2	1:B:287:GLN:N	2.10	0.66
1:B:107:ILE:HG13	1:B:111:TRP:NE1	2.11	0.66
1:A:81:ALA:HB2	1:B:81:ALA:HB2	1.79	0.64
1:A:285:MET:HE2	1:A:287:GLN:N	2.14	0.63
1:B:134:GLY:HA3	1:B:169:PHE:CZ	2.34	0.62
1:C:108:ASP:HA	1:C:111:TRP:HD1	1.65	0.61
1:A:134:GLY:HA3	1:A:169:PHE:CZ	2.36	0.60
1:A:57:LEU:O	1:A:61:VAL:HG22	2.02	0.59
1:C:134:GLY:HA3	1:C:169:PHE:CZ	2.38	0.59
1:D:285:MET:HE2	1:D:287:GLN:N	2.15	0.58
1:A:108:ASP:HA	1:A:111:TRP:HD1	1.68	0.58
1:B:160:PHE:O	1:B:164:ARG:HD2	2.05	0.56
1:A:107:ILE:HG13	1:A:111:TRP:NE1	2.21	0.55
1:A:212:PHE:CE1	1:A:273:ALA:HB2	2.43	0.53
1:C:240:TRP:HB2	1:C:282:ASP:OD2	2.08	0.53
1:C:57:LEU:O	1:C:61:VAL:HG22	2.08	0.53
1:B:68:VAL:HG11	1:B:73:ILE:HD11	1.91	0.53
1:B:57:LEU:O	1:B:61:VAL:HG22	2.10	0.52
1:A:56:ASN:O	1:A:60:VAL:HG23	2.11	0.51
1:A:144:PHE:HB2	1:A:194:PRO:O	2.10	0.50
1:A:215:GLU:HB2	1:A:290:TRP:CD1	2.47	0.50
1:A:210:VAL:HB	1:A:266:GLN:OE1	2.13	0.49
1:B:108:ASP:HA	1:B:111:TRP:HD1	1.78	0.49
1:C:157:TYR:O	1:C:161:VAL:HG23	2.14	0.47
1:B:240:TRP:HB2	1:B:282:ASP:OD2	2.14	0.47
1:A:117:GLY:HA2	1:A:192:ASN:CG	2.35	0.47
1:B:297:GLU:O	1:B:298:ARG:CB	2.63	0.46
1:A:136:TRP:CE2	1:A:140:ASN:ND2	2.83	0.46
1:B:137:LEU:HD22	1:B:144:PHE:CD2	2.51	0.46
1:D:208:VAL:HG23	1:D:221:ALA:HB2	1.97	0.46
1:A:235:PHE:CE1	1:A:253:GLU:HB3	2.50	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:212:PHE:CE1	1:D:273:ALA:HB2	2.51	0.45
1:D:285:MET:HE3	1:D:285:MET:HA	1.99	0.45
1:C:108:ASP:HA	1:C:111:TRP:CD1	2.49	0.45
1:D:285:MET:CE	1:D:285:MET:HA	2.45	0.45
1:C:136:TRP:CE2	1:C:140:ASN:ND2	2.85	0.45
1:B:136:TRP:CE2	1:B:140:ASN:ND2	2.85	0.45
1:B:212:PHE:CE1	1:B:273:ALA:HB2	2.53	0.44
1:C:107:ILE:O	1:C:111:TRP:CD1	2.71	0.44
1:A:134:GLY:HA3	1:A:169:PHE:HZ	1.80	0.43
1:D:180:PRO:HD2	1:D:183:GLU:OE1	2.19	0.43
1:B:56:ASN:O	1:B:60:VAL:HG23	2.18	0.43
1:D:169:PHE:N	1:D:169:PHE:CD1	2.86	0.43
1:A:285:MET:HA	1:A:285:MET:HE3	2.00	0.43
1:D:149:GLU:HA	1:D:174:LEU:HD12	2.00	0.43
1:C:266:GLN:NE2	1:C:270:ASP:OD1	2.50	0.43
1:A:169:PHE:CD1	1:A:169:PHE:N	2.85	0.42
1:B:184:VAL:O	1:B:188:ARG:HG2	2.19	0.42
1:A:109:HIS:HB3	1:A:235:PHE:CE1	2.54	0.42
1:B:163:TYR:CZ	1:B:167:LEU:HD11	2.54	0.42
1:C:285:MET:CE	1:C:285:MET:HA	2.49	0.42
1:D:56:ASN:O	1:D:60:VAL:HG23	2.19	0.42
1:A:55:ARG:O	1:A:58:ALA:HB3	2.19	0.42
1:A:208:VAL:HG11	1:A:262:THR:HA	2.00	0.42
1:B:130:TRP:CH2	1:B:284:HIS:CE1	3.08	0.42
1:D:57:LEU:O	1:D:61:VAL:HG22	2.19	0.42
1:D:130:TRP:CH2	1:D:284:HIS:CE1	3.07	0.42
1:D:134:GLY:HA3	1:D:169:PHE:CZ	2.55	0.42
1:D:215:GLU:HB2	1:D:290:TRP:CD1	2.54	0.42
1:A:108:ASP:HA	1:A:111:TRP:CD1	2.51	0.42
1:B:157:TYR:O	1:B:161:VAL:HG23	2.19	0.42
1:B:208:VAL:HG11	1:B:262:THR:HA	2.02	0.41
1:C:195:ILE:CD1	1:C:226:LEU:HD13	2.50	0.41
1:C:208:VAL:HG23	1:C:221:ALA:HB2	2.02	0.41
1:B:189:LEU:CD1	1:B:226:LEU:HB3	2.51	0.41
1:B:164:ARG:HB3	1:B:169:PHE:CD2	2.56	0.41
1:D:180:PRO:HA	1:D:181:PRO:HD3	1.94	0.41
1:D:208:VAL:CG2	1:D:221:ALA:HB2	2.51	0.41
1:D:233:ALA:HB3	1:D:235:PHE:CE2	2.55	0.40
1:C:285:MET:HA	1:C:285:MET:HE3	2.03	0.40
1:D:240:TRP:HB2	1:D:282:ASP:OD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/313 (79%)	241 (98%)	6 (2%)	0	100	100
1	B	247/313 (79%)	243 (98%)	4 (2%)	0	100	100
1	C	247/313 (79%)	242 (98%)	5 (2%)	0	100	100
1	D	247/313 (79%)	243 (98%)	4 (2%)	0	100	100
All	All	988/1252 (79%)	969 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/244 (65%)	158 (100%)	0	100	100
1	B	161/244 (66%)	161 (100%)	0	100	100
1	C	143/244 (59%)	143 (100%)	0	100	100
1	D	165/244 (68%)	165 (100%)	0	100	100
All	All	627/976 (64%)	627 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	109	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	GOL	A	401	-	5,5,5	0.75	0	5,5,5	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	401	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	249/313 (79%)	0.01	0	100	100	93, 122, 151, 163	0
1	B	249/313 (79%)	0.02	1 (0%)	92	88	89, 114, 143, 154	0
1	C	249/313 (79%)	0.09	3 (1%)	79	68	103, 132, 163, 176	0
1	D	249/313 (79%)	0.16	3 (1%)	79	68	97, 126, 154, 169	0
All	All	996/1252 (79%)	0.07	7 (0%)	87	80	89, 124, 154, 176	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	199	ALA	3.2
1	D	199	ALA	2.7
1	C	150	ARG	2.7
1	D	116	ALA	2.5
1	D	289	GLN	2.5
1	B	199	ALA	2.2
1	C	57	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	GOL	A	401	6/6	0.84	0.31	111,137,154,154	0

6.5 Other polymers (i)

There are no such residues in this entry.