



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2024 – 06:34 PM EDT

PDB ID : 2OKG
Title : Structure of effector binding domain of central glycolytic gene regulator (CggR) from *B. subtilis*
Authors : Rezacova, P.; Moy, S.F.; Joachimiak, A.; Otwinowski, Z.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2007-01-16
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	255	85% 11% ..
1	B	255	84% 13% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	G3H	B	501	X	-	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4367 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Central glycolytic gene regulator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	249	1926	1210	332	370	4	10	0	9	0
1	B	251	1956	1228	339	377	3	9	0	11	0


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	86	SER	-	cloning artifact	UNP O32253
A	87	ASN	-	cloning artifact	UNP O32253
A	88	ALA	-	cloning artifact	UNP O32253
A	127	MSE	MET	modified residue	UNP O32253
A	135	MSE	MET	modified residue	UNP O32253
A	159	MSE	MET	modified residue	UNP O32253
A	160	MSE	MET	modified residue	UNP O32253
A	193	MSE	MET	modified residue	UNP O32253
A	236	MSE	MET	modified residue	UNP O32253
A	247	MSE	MET	modified residue	UNP O32253
A	290	MSE	MET	modified residue	UNP O32253
B	86	SER	-	cloning artifact	UNP O32253
B	87	ASN	-	cloning artifact	UNP O32253
B	88	ALA	-	cloning artifact	UNP O32253
B	127	MSE	MET	modified residue	UNP O32253
B	135	MSE	MET	modified residue	UNP O32253
B	159	MSE	MET	modified residue	UNP O32253
B	160	MSE	MET	modified residue	UNP O32253
B	193	MSE	MET	modified residue	UNP O32253
B	236	MSE	MET	modified residue	UNP O32253
B	247	MSE	MET	modified residue	UNP O32253
B	290	MSE	MET	modified residue	UNP O32253

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Cl 3 3	0	0

- Molecule 3 is GLYCERALDEHYDE-3-PHOSPHATE (three-letter code: G3H) (formula: C₃H₇O₆P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O P 10 3 6 1	0	0


- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	214	Total O 214 214	0	0
4	B	258	Total O 258 258	0	3

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Central glycolytic gene regulator

Chain A:  85% 11% ..



- Molecule 1: Central glycolytic gene regulator

Chain B:  84% 13% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.60Å 83.80Å 116.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.04 – 1.65 34.04 – 1.65	Depositor EDS
% Data completeness (in resolution range)	96.0 (34.04-1.65) 96.0 (34.04-1.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.61 (at 1.65Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.196 , 0.239 0.245 , 0.277	Depositor DCC
R_{free} test set	3089 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	21.1	Xtrriage
Anisotropy	0.153	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4367	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G3H, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	4/1963 (0.2%)	0.76	1/2625 (0.0%)
1	B	0.80	5/1997 (0.3%)	0.73	4/2672 (0.1%)
All	All	0.78	9/3960 (0.2%)	0.75	5/5297 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	330	GLU	CD-OE2	-9.79	1.14	1.25
1	B	120	SER	CB-OG	9.11	1.54	1.42
1	B	330	GLU	CG-CD	8.80	1.65	1.51
1	B	330	GLU	CD-OE1	8.76	1.35	1.25
1	A	179	GLY	C-O	8.51	1.37	1.23
1	B	330	GLU	CB-CG	7.61	1.66	1.52
1	A	249	GLN	CD-OE1	6.50	1.38	1.24
1	A	184	ASN	C-N	5.13	1.45	1.34
1	A	338	ARG	CZ-NH1	5.06	1.39	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	338	ARG	NE-CZ-NH2	-7.89	116.36	120.30
1	B	330	GLU	OE1-CD-OE2	6.94	131.63	123.30
1	B	255	LEU	CB-CG-CD1	5.80	120.87	111.00
1	B	178	LEU	CB-CG-CD1	5.62	120.56	111.00
1	B	330	GLU	CB-CG-CD	-5.49	99.38	114.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1926	0	1993	27	0
1	B	1956	0	2019	34	0
2	A	3	0	0	0	0
3	B	10	0	5	1	0
4	A	214	0	0	0	0
4	B	258	0	0	7	0
All	All	4367	0	4017	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:ILE:CD1	1:A:134[B]:CYS:SG	2.53	0.96
1:B:158:GLU:HG3	1:B:159[A]:MSE:HE2	1.51	0.91
1:A:111:ILE:HD13	1:A:134[B]:CYS:SG	2.14	0.87
1:B:158:GLU:HG3	1:B:159[A]:MSE:CE	2.12	0.78
1:B:134[B]:CYS:SG	1:B:138[B]:ARG:NH1	2.62	0.72
1:B:129:ARG:HG3	1:B:159[B]:MSE:HE2	1.72	0.71
1:A:111:ILE:HD12	1:A:134[B]:CYS:SG	2.30	0.71
1:A:127[B]:MSE:HE2	1:A:304:VAL:HG12	1.75	0.68
1:B:99:LYS:O	1:B:103:GLU:HG3	1.94	0.68
1:B:228:LEU:HD21	1:B:288:VAL:HG22	1.77	0.66
1:A:219[A]:ILE:HG23	1:A:228:LEU:HD11	1.78	0.65
1:A:219[A]:ILE:HG23	1:A:228:LEU:CD1	2.28	0.64
1:A:334:LYS:O	1:A:338:ARG:HG3	1.99	0.63
1:A:99:LYS:NZ	1:A:103:GLU:OE2	2.36	0.59
1:B:247:MSE:SE	3:B:501:G3H:H32	2.55	0.57
1:A:89:LYS:HD2	1:B:89:LYS:H	1.69	0.57
1:A:191:ALA:O	1:A:195:GLU:HG3	2.06	0.55
1:B:94:LEU:HD21	1:B:330:GLU:HG3	1.89	0.55
1:B:159[A]:MSE:HE3	4:B:610:HOH:O	2.06	0.55
1:B:228:LEU:CD2	1:B:288:VAL:HG22	2.37	0.54
1:A:127[B]:MSE:HE1	1:A:305:ALA:N	2.23	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:ARG:NE	1:B:114:SER:HB3	2.24	0.53
1:B:124:LYS:HA	1:B:127:MSE:HE3	1.91	0.53
1:B:106[B]:ASN:ND2	4:B:692:HOH:O	2.41	0.52
1:B:268:THR:CG2	1:B:275:PHE:HB2	2.39	0.52
1:A:154:GLU:HA	1:A:189:ILE:HD12	1.92	0.52
1:A:87:ASN:HD22	1:A:88:ALA:H	1.58	0.51
1:A:129:ARG:N	1:A:159[A]:MSE:HG2	2.26	0.51
1:B:175[B]:ARG:HA	4:B:709[B]:HOH:O	2.10	0.51
1:B:191:ALA:O	1:B:195[B]:GLU:HG3	2.13	0.49
1:B:90:ASP:OD1	1:B:95:THR:HG21	2.13	0.48
1:B:176:GLY:O	1:B:178:LEU:HD13	2.12	0.48
1:A:87:ASN:HD22	1:A:88:ALA:N	2.12	0.48
1:A:215:TYR:O	1:A:219[A]:ILE:HG13	2.13	0.48
1:B:247:MSE:HE2	1:B:251:ARG:CZ	2.43	0.47
1:A:127[B]:MSE:HE2	1:A:304:VAL:CG1	2.44	0.47
1:B:88:ALA:N	4:B:720:HOH:O	2.48	0.46
1:B:92:LEU:HD22	4:B:664:HOH:O	2.15	0.46
1:B:247:MSE:HE1	1:B:250:ARG:NH2	2.31	0.46
1:A:149:GLY:O	1:A:185:GLN:HG3	2.16	0.46
1:B:134[B]:CYS:SG	1:B:138[B]:ARG:CZ	3.04	0.46
1:B:247:MSE:HE1	1:B:250:ARG:HH21	1.81	0.46
1:B:106[A]:ASN:ND2	4:B:637:HOH:O	2.49	0.45
1:A:219[A]:ILE:CG2	1:A:228:LEU:CD1	2.94	0.44
1:B:191:ALA:O	1:B:195[A]:GLU:HG3	2.18	0.44
1:B:228:LEU:HD21	1:B:288:VAL:CG2	2.45	0.44
1:B:180:GLU:OE2	1:B:247:MSE:HE1	2.17	0.43
1:A:334:LYS:O	1:A:338:ARG:CG	2.66	0.43
1:B:268:THR:HG23	1:B:275:PHE:HB2	2.01	0.43
1:A:149:GLY:O	1:A:185:GLN:CG	2.67	0.43
1:A:129:ARG:HA	1:A:159[A]:MSE:CG	2.49	0.42
1:B:154:GLU:OE1	4:B:754:HOH:O	2.21	0.42
1:B:92:LEU:HD13	1:B:114:SER:OG	2.19	0.42
1:A:129:ARG:HA	1:A:159[A]:MSE:HG3	2.01	0.42
1:B:247:MSE:HE2	1:B:251:ARG:NE	2.35	0.42
1:A:190[A]:CYS:SG	1:A:203:LEU:HD23	2.61	0.41
1:A:221[B]:GLU:HA	1:A:222:PRO:HD2	1.97	0.40
1:A:106:ASN:HB3	1:A:321:PRO:HA	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/255 (100%)	252 (99%)	2 (1%)	0	100	100
1	B	260/255 (102%)	259 (100%)	1 (0%)	0	100	100
All	All	514/510 (101%)	511 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/199 (106%)	206 (98%)	4 (2%)	57	34
1	B	212/199 (106%)	209 (99%)	3 (1%)	67	46
All	All	422/398 (106%)	415 (98%)	7 (2%)	65	39

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	A	190[A]	CYS
1	A	190[B]	CYS
1	A	222	PRO
1	B	178	LEU
1	B	262	ASP
1	B	287	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	A	106	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	G3H	B	501	-	8,9,9	2.36	1 (12%)	10,12,12	1.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	G3H	B	501	-	1/1/2/3	2/7/8/8	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	501	G3H	O1-C1	6.31	1.45	1.19

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	B	501	G3H	C2

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	501	G3H	O1-C1-C2-C3
3	B	501	G3H	O2-C2-C3-O1P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	501	G3H	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.