



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 04:19 am BST

PDB ID : 1OLC  
Title : OLIGO-PEPTIDE BINDING PROTEIN (OPPA) COMPLEXED WITH  
LYS-LYS-LYS-ALA  
Authors : Tame, J.; Wilkinson, A.J.  
Deposited on : 1995-09-10  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	517	<p>76% 20%</p>
2	B	4	<p>25% 75%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IUM	A	522	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4553 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called OLIGO-PEPTIDE BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	517	4165	2666	700	794	5	4	0	0

- Molecule 2 is a protein called LYS-LYS-LYS-ALA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	4	33	21	7	5	0	0	0

- Molecule 3 is URANYL (VI) ION (three-letter code: IUM) (formula: O<sub>2</sub>U).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	U		
3	A	1	3	2	1	0	0
3	A	1	3	2	1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O U 3 2 1	0	0
3	A	1	Total O U 3 2 1	0	0
3	A	1	Total O U 3 2 1	0	0
3	A	1	Total U 1 1	0	0
3	A	1	Total U 1 1	0	0
3	A	1	Total U 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	335	Total O 335 335	0	0
4	B	2	Total O 2 2	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: OLIGO-PEPTIDE BINDING PROTEIN



- Molecule 2: LYS-LYS-LYS-ALA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.73Å 77.08Å 71.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.10 19.99 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (10.00-2.10) 99.7 (19.99-2.00)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.61 (at 2.01Å)	Xtrriage
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.143 , 0.197 0.136 , 0.184	Depositor DCC
$R_{free}$ test set	2111 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.9	Xtrriage
Anisotropy	0.173	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 81.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4553	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IUM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.91	3/4276 (0.1%)	1.66	54/5830 (0.9%)
2	B	1.11	0/32	1.59	1/37 (2.7%)
All	All	0.92	3/4308 (0.1%)	1.66	55/5867 (0.9%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	17	ARG	CD-NE	-12.01	1.26	1.46
1	A	170	SER	CB-OG	6.10	1.50	1.42
1	A	12	LYS	CB-CG	-5.58	1.37	1.52

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	17	ARG	CD-NE-CZ	39.30	178.62	123.60
1	A	17	ARG	CG-CD-NE	16.62	146.69	111.80
1	A	369	ASP	CB-CG-OD1	-13.78	105.90	118.30
1	A	133	ASP	CB-CG-OD2	-12.63	106.94	118.30
1	A	77	ARG	NE-CZ-NH2	9.84	125.22	120.30
1	A	288	ARG	NE-CZ-NH1	9.54	125.07	120.30
1	A	81	LYS	CA-CB-CG	8.38	131.84	113.40
1	A	290	ARG	NE-CZ-NH1	8.25	124.42	120.30
1	A	235	ARG	CD-NE-CZ	7.92	134.69	123.60
1	A	137	LYS	CA-CB-CG	7.87	130.72	113.40
1	A	489	ARG	NE-CZ-NH1	7.79	124.20	120.30
1	A	242	ASP	CB-CG-OD2	7.76	125.28	118.30
1	A	102	ASP	CB-CG-OD1	7.66	125.19	118.30
1	A	133	ASP	OD1-CG-OD2	7.45	137.45	123.30
1	A	32	GLU	OE1-CD-OE2	7.41	132.19	123.30
1	A	81	LYS	CB-CG-CD	7.40	130.84	111.60

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	84	ASP	CB-CG-OD1	7.35	124.92	118.30
1	A	53	GLU	OE1-CD-OE2	7.34	132.10	123.30
1	A	251	GLU	CA-CB-CG	-7.31	97.32	113.40
1	A	342	GLU	CA-CB-CG	7.16	129.16	113.40
1	A	236	TYR	CB-CG-CD2	7.03	125.22	121.00
1	A	459	ASP	CB-CG-OD2	-7.02	111.98	118.30
1	A	410	ASP	CB-CG-OD2	-7.00	112.00	118.30
1	A	51	ASP	CB-CG-OD1	6.82	124.43	118.30
1	A	273	TYR	CB-CG-CD2	-6.57	117.06	121.00
1	A	433	ASP	CB-CG-OD1	6.45	124.10	118.30
1	A	474	ASP	CB-CG-OD2	-6.10	112.81	118.30
1	A	36	GLU	OE1-CD-OE2	-5.99	116.11	123.30
1	A	396	GLU	OE1-CD-OE2	-5.97	116.13	123.30
1	A	245	TYR	CB-CG-CD1	5.94	124.56	121.00
1	A	122	ASP	CB-CG-OD1	5.82	123.54	118.30
1	A	41	ARG	NE-CZ-NH2	-5.69	117.45	120.30
1	A	288	ARG	NH1-CZ-NH2	-5.68	113.15	119.40
1	A	513	TYR	CB-CG-CD2	-5.68	117.59	121.00
1	A	245	TYR	CB-CG-CD2	-5.66	117.60	121.00
1	A	53	GLU	CA-CB-CG	-5.65	100.97	113.40
1	A	508	TYR	CB-CG-CD2	5.59	124.36	121.00
2	B	3	LYS	CA-CB-CG	-5.59	101.11	113.40
1	A	459	ASP	CB-CG-OD1	5.41	123.17	118.30
1	A	236	TYR	CB-CG-CD1	-5.40	117.76	121.00
1	A	342	GLU	N-CA-CB	-5.39	100.89	110.60
1	A	331	GLU	N-CA-CB	5.29	120.11	110.60
1	A	5	ALA	CB-CA-C	5.24	117.96	110.10
1	A	218	ILE	CA-CB-CG1	-5.23	101.06	111.00
1	A	242	ASP	CB-CG-OD1	-5.22	113.60	118.30
1	A	102	ASP	CB-CG-OD2	-5.21	113.61	118.30
1	A	109	TYR	CB-CG-CD1	-5.21	117.87	121.00
1	A	396	GLU	CB-CG-CD	5.20	128.24	114.20
1	A	117	HIS	CA-CB-CG	-5.18	104.79	113.60
1	A	413	ARG	NE-CZ-NH2	5.16	122.88	120.30
1	A	62	GLU	CA-CB-CG	5.14	124.70	113.40
1	A	386	LEU	CA-CB-CG	5.14	127.12	115.30
1	A	191	TYR	CB-CG-CD1	-5.07	117.96	121.00
1	A	400	PHE	N-CA-CB	-5.04	101.54	110.60
1	A	87	PRO	N-CA-CB	5.00	109.30	103.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4165	0	4076	54	0
2	B	33	0	46	2	0
3	A	18	0	0	5	0
4	A	335	0	0	15	0
4	B	2	0	0	1	0
All	All	4553	0	4122	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:519:IUM:O2	4:A:537:HOH:O	1.84	0.95
1:A:489:ARG:NH2	4:A:728:HOH:O	2.02	0.91
1:A:326:LYS:HE2	4:A:753:HOH:O	1.76	0.83
1:A:398:LYS:HG3	4:A:822:HOH:O	1.81	0.80
1:A:300:ASP:OD1	4:A:770:HOH:O	2.01	0.79
3:A:522:IUM:U	3:A:522:IUM:O1	1.67	0.75
1:A:396:GLU:HB3	1:A:398:LYS:HZ3	1.53	0.73
1:A:505:ASP:OD2	4:A:637:HOH:O	2.06	0.73
1:A:21:SER:HB2	4:A:715:HOH:O	1.91	0.70
3:A:522:IUM:U	3:A:522:IUM:O2	1.75	0.67
3:A:518:IUM:U	3:A:518:IUM:O2	1.73	0.67
1:A:492:LYS:HE2	4:A:785:HOH:O	1.99	0.62
1:A:286:ASP:OD1	1:A:288:ARG:HD3	2.00	0.62
1:A:428:ASN:HA	1:A:431:LEU:HD22	1.80	0.62
1:A:396:GLU:HB3	1:A:398:LYS:NZ	2.13	0.61
1:A:229:GLU:OE1	1:A:369:ASP:HB2	2.00	0.61
3:A:521:IUM:O2	3:A:521:IUM:U	1.81	0.61
1:A:113:LEU:HD22	1:A:118:ILE:HD12	1.83	0.60
1:A:354:THR:OG1	1:A:357:LYS:HE3	2.02	0.60
1:A:301:ILE:HA	1:A:305:LYS:HD2	1.85	0.59
1:A:24:GLN:HG2	1:A:36:GLU:HG2	1.86	0.56
1:A:354:THR:OG1	1:A:357:LYS:HG2	2.09	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:161:HIS:CD2	1:A:162:PRO:HD2	2.44	0.52
1:A:323:ASP:O	1:A:423:PRO:HD3	2.10	0.52
1:A:354:THR:HG1	1:A:357:LYS:HE3	1.75	0.51
1:A:43:LEU:O	1:A:187:THR:HB	2.11	0.50
1:A:383:LYS:HE2	1:A:389:ASN:OD1	2.12	0.50
1:A:257:LYS:O	1:A:261:PRO:HB3	2.13	0.48
1:A:354:THR:HG1	1:A:356:ASP:HB3	1.78	0.48
1:A:210:TYR:CE2	1:A:212:ASP:HB3	2.48	0.48
1:A:346:LYS:O	1:A:350:GLU:HG2	2.13	0.48
1:A:358:PRO:HA	4:A:652:HOH:O	2.13	0.48
1:A:398:LYS:HD3	1:A:399:THR:H	1.80	0.47
4:A:528:HOH:O	2:B:2:LYS:HE3	2.15	0.47
1:A:57:SER:HB3	1:A:58:PRO:HD2	1.97	0.46
1:A:3:VAL:HA	1:A:4:PRO:HD3	1.84	0.46
1:A:439:ALA:HB1	4:A:783:HOH:O	2.16	0.45
2:B:1:LYS:NZ	4:B:208:HOH:O	2.18	0.45
1:A:230:VAL:HG21	1:A:251:GLU:OE1	2.16	0.45
1:A:501:LYS:HE3	4:A:691:HOH:O	2.15	0.45
1:A:262:ASN:HB2	4:A:816:HOH:O	2.16	0.45
1:A:161:HIS:HA	1:A:162:PRO:HD3	1.88	0.45
1:A:296:ALA:HB3	1:A:386:LEU:HD22	1.98	0.45
1:A:398:LYS:HD2	4:A:616:HOH:O	2.17	0.44
1:A:130:PRO:O	1:A:133:ASP:HB2	2.17	0.44
1:A:373:LYS:HA	1:A:373:LYS:HD3	1.79	0.43
1:A:398:LYS:HD3	1:A:398:LYS:N	2.34	0.43
1:A:498:TYR:CE1	1:A:507:ILE:HD11	2.53	0.43
1:A:516:LYS:HE3	1:A:516:LYS:HB3	1.72	0.43
1:A:384:LYS:HD3	1:A:385:ASN:OD1	2.20	0.42
1:A:354:THR:OG1	1:A:357:LYS:N	2.49	0.42
1:A:462:ARG:HD2	1:A:462:ARG:HH11	1.66	0.42
1:A:336:SER:OG	1:A:338:GLN:HG2	2.20	0.42
1:A:363:LEU:HD13	1:A:411:VAL:HG12	2.00	0.42
1:A:342:GLU:HA	1:A:342:GLU:OE1	2.20	0.42
1:A:253:PHE:CD2	1:A:309:GLN:HG2	2.55	0.41
1:A:356:ASP:HB3	1:A:357:LYS:HE3	2.02	0.41
1:A:208:PRO:HD2	4:A:849:HOH:O	2.20	0.41
1:A:281:LYS:O	1:A:284:PHE:N	2.53	0.41
1:A:264:VAL:HG22	1:A:490:LEU:CD2	2.51	0.41
1:A:294:LYS:HA	1:A:480:VAL:HG13	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	515/517 (100%)	499 (97%)	15 (3%)	1 (0%)	47	49
2	B	2/4 (50%)	2 (100%)	0	0	100	100
All	All	517/521 (99%)	501 (97%)	15 (3%)	1 (0%)	47	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	455/455 (100%)	423 (93%)	32 (7%)	15	12
2	B	3/3 (100%)	3 (100%)	0	100	100
All	All	458/458 (100%)	426 (93%)	32 (7%)	15	12

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ASP
1	A	12	LYS
1	A	17	ARG
1	A	57	SER
1	A	63	LYS

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	67	LYS
1	A	123	ASP
1	A	133	ASP
1	A	137	LYS
1	A	173	GLU
1	A	206	ARG
1	A	215	LYS
1	A	258	LYS
1	A	262	ASN
1	A	297	LEU
1	A	328	VAL
1	A	331	GLU
1	A	342	GLU
1	A	346	LYS
1	A	348	LEU
1	A	350	GLU
1	A	357	LYS
1	A	369	ASP
1	A	383	LYS
1	A	398	LYS
1	A	402	ASP
1	A	413	ARG
1	A	431	LEU
1	A	442	LYS
1	A	461	GLN
1	A	468	LYS
1	A	516	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	199	ASN
1	A	209	GLN
1	A	461	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are modelled with single atom - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	IUM	A	520	-	0,2,2	0.00	-	-		
3	IUM	A	521	-	0,2,2	0.00	-	-		
3	IUM	A	519	-	0,2,2	0.00	-	-		
3	IUM	A	518	-	0,2,2	0.00	-	-		
3	IUM	A	522	-	0,2,2	0.00	-	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	521	IUM	1	0
3	A	519	IUM	1	0
3	A	518	IUM	1	0
3	A	522	IUM	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	517/517 (100%)	-0.73	3 (0%) 89   91	7, 16, 46, 87	1 (0%)
2	B	4/4 (100%)	-1.02	0 100   100	11, 12, 12, 13	0
All	All	521/521 (100%)	-0.73	3 (0%) 89   91	7, 16, 46, 87	1 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	356	ASP	4.1
1	A	354	THR	3.0
1	A	357	LYS	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	IUM	A	523	1/3	0.99	0.04	37,37,37,37	0
3	IUM	A	525	1/3	0.99	0.05	51,51,51,51	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	IUM	A	524	1/3	0.99	0.05	45,45,45,45	0
3	IUM	A	520	3/3	1.00	0.04	6,6,14,19	0
3	IUM	A	522	3/3	1.00	0.04	3,3,4,19	0
3	IUM	A	521	3/3	1.00	0.04	11,11,21,23	0
3	IUM	A	518	3/3	1.00	0.03	2,2,2,13	0
3	IUM	A	519	3/3	1.00	0.03	7,7,18,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.