



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2023 – 01:02 PM EDT

PDB ID : 2OL7
Title : The crystal structure of OspA mutant
Authors : Makabe, K.; Terechko, V.; Koide, S.
Deposited on : 2007-01-18
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	
1	B	251	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer surface protein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	1921	1197	309	412	3	0	15	0
1	B	246	1846	1145	300	400	1	0	7	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	GLY	-	expression tag	UNP Q45040
A	24	SER	-	expression tag	UNP Q45040
A	25	HIS	-	expression tag	UNP Q45040
A	26	MET	-	expression tag	UNP Q45040
A	37	SER	GLU	engineered mutation	UNP Q45040
A	45	SER	GLU	engineered mutation	UNP Q45040
A	46	SER	LYS	engineered mutation	UNP Q45040
A	48	ALA	LYS	engineered mutation	UNP Q45040
A	60	ALA	LYS	engineered mutation	UNP Q45040
A	64	SER	LYS	engineered mutation	UNP Q45040
A	83	ALA	LYS	engineered mutation	UNP Q45040
A	104	SER	GLU	engineered mutation	UNP Q45040
A	107	SER	LYS	engineered mutation	UNP Q45040
A	126	ILE	PHE	engineered mutation	UNP Q45040
A	127	ILE	ASN	engineered mutation	UNP Q45040
A	128	ILE	GLU	engineered mutation	UNP Q45040
A	129	ASP	LYS	engineered mutation	UNP Q45040
A	131	ILE	GLU	engineered mutation	UNP Q45040
A	132	ILE	VAL	engineered mutation	UNP Q45040
A	133	ILE	SER	engineered mutation	UNP Q45040
A	239	SER	LYS	engineered mutation	UNP Q45040
A	240	SER	GLU	engineered mutation	UNP Q45040
A	254	SER	LYS	engineered mutation	UNP Q45040
B	23	GLY	-	expression tag	UNP Q45040
B	24	SER	-	expression tag	UNP Q45040

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	25	HIS	-	expression tag	UNP Q45040
B	26	MET	-	expression tag	UNP Q45040
B	37	SER	GLU	engineered mutation	UNP Q45040
B	45	SER	GLU	engineered mutation	UNP Q45040
B	46	SER	LYS	engineered mutation	UNP Q45040
B	48	ALA	LYS	engineered mutation	UNP Q45040
B	60	ALA	LYS	engineered mutation	UNP Q45040
B	64	SER	LYS	engineered mutation	UNP Q45040
B	83	ALA	LYS	engineered mutation	UNP Q45040
B	104	SER	GLU	engineered mutation	UNP Q45040
B	107	SER	LYS	engineered mutation	UNP Q45040
B	126	ILE	PHE	engineered mutation	UNP Q45040
B	127	ILE	ASN	engineered mutation	UNP Q45040
B	128	ILE	GLU	engineered mutation	UNP Q45040
B	129	ASP	LYS	engineered mutation	UNP Q45040
B	131	ILE	GLU	engineered mutation	UNP Q45040
B	132	ILE	VAL	engineered mutation	UNP Q45040
B	133	ILE	SER	engineered mutation	UNP Q45040
B	239	SER	LYS	engineered mutation	UNP Q45040
B	240	SER	GLU	engineered mutation	UNP Q45040
B	254	SER	LYS	engineered mutation	UNP Q45040

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	350	Total O 350 350	0	0
2	B	218	Total O 218 218	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Outer surface protein A



- Molecule 1: Outer surface protein A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	37.13Å 55.33Å 64.79Å 85.68° 74.05° 86.45°	Depositor
Resolution (Å)	20.00 – 1.35 29.37 – 1.35	Depositor EDS
% Data completeness (in resolution range)	95.2 (20.00-1.35) 95.2 (29.37-1.35)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.27 (at 1.35Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.184 , 0.218 0.209 , 0.239	Depositor DCC
R_{free} test set	5140 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	13.7	Xtrriage
Anisotropy	0.378	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 54.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4335	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1975	0.78	0/2653
1	B	0.56	0/1875	0.70	0/2523
All	All	0.60	0/3850	0.74	0/5176

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	119	LYS	Peptide
1	B	137	ILE	Peptide
1	B	138	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1921	0	2041	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1846	0	1940	31	0
2	A	350	0	0	6	0
2	B	218	0	0	8	0
All	All	4335	0	3981	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:LYS:HB2	1:B:120:SER:HB3	1.29	1.15
1:A:69:LYS:HE2	2:A:524:HOH:O	1.52	1.07
1:B:119:LYS:HB2	1:B:120:SER:CB	1.95	0.97
1:B:139:ARG:HB2	1:B:143:THR:O	1.72	0.88
1:B:119:LYS:CB	1:B:120:SER:HB3	2.04	0.87
1:B:101:VAL:HG13	2:B:362:HOH:O	1.76	0.85
1:A:136:ILE:HD11	2:A:619:HOH:O	1.84	0.77
1:B:135:LYS:NZ	1:B:137:ILE:HD11	2.02	0.73
1:A:192:SER:OG	1:A:196[B]:GLU:HG2	1.94	0.67
1:B:144:ARG:HE	1:B:146:GLU:CD	2.01	0.63
1:B:135:LYS:HZ3	1:B:137:ILE:HD11	1.62	0.63
1:B:123:GLU:CD	1:B:135:LYS:HE2	2.22	0.60
1:B:139:ARG:O	1:B:140:ALA:HB2	2.02	0.60
1:B:121:SER:HA	1:B:139:ARG:O	2.03	0.59
1:B:119:LYS:CB	1:B:120:SER:CB	2.74	0.58
1:A:119:LYS:HD2	1:A:119:LYS:H	1.68	0.57
1:B:135:LYS:HA	2:B:347:HOH:O	2.05	0.56
1:A:148:THR:HG22	2:A:619:HOH:O	2.06	0.55
1:A:119:LYS:H	1:A:119:LYS:CD	2.19	0.55
1:A:24:SER:N	2:A:610:HOH:O	2.40	0.54
1:B:139:ARG:O	1:B:140:ALA:CB	2.56	0.53
1:B:148:THR:HG22	2:B:422:HOH:O	2.08	0.53
1:B:119:LYS:N	1:B:120:SER:OG	2.42	0.52
1:B:119:LYS:HB2	1:B:120:SER:OG	2.08	0.52
1:B:33:ASP:OD1	1:B:34:LEU:N	2.43	0.51
1:B:123:GLU:OE2	1:B:135:LYS:HE2	2.13	0.48
1:A:38[B]:MET:CE	1:A:58:VAL:HG22	2.44	0.48
1:A:159:LYS:HG3	1:A:166:VAL:CG1	2.44	0.48
1:A:38[A]:MET:HE2	2:A:470:HOH:O	2.14	0.47
1:A:148:THR:CG2	2:A:619:HOH:O	2.63	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:134:GLU:C	2:B:347:HOH:O	2.52	0.47
1:B:123:GLU:OE2	1:B:135:LYS:CE	2.63	0.47
1:A:133:ILE:O	1:A:134[A]:GLU:HG2	2.15	0.46
1:B:148:THR:O	2:B:408:HOH:O	2.21	0.44
1:B:118:ASP:N	1:B:118:ASP:OD1	2.51	0.44
1:B:136:ILE:HG22	1:B:138:THR:OG1	2.16	0.44
1:B:247:GLN:NE2	2:B:416:HOH:O	2.42	0.44
1:B:120:SER:O	1:B:139:ARG:O	2.34	0.44
1:A:192:SER:CB	1:A:196[B]:GLU:HG2	2.48	0.43
1:B:95:GLY:HA2	2:B:443:HOH:O	2.18	0.43
1:B:82:ASP:O	1:B:83:ALA:HB3	2.19	0.42
1:A:82:ASP:O	1:A:83:ALA:HB3	2.20	0.42
1:B:139:ARG:NH2	1:B:160:GLU:OE2	2.53	0.41
1:A:38[B]:MET:HE1	1:A:58:VAL:HG22	2.01	0.41
1:A:159:LYS:CG	1:A:166:VAL:CG1	2.99	0.41
1:B:163[A]:LYS:HG2	2:B:301:HOH:O	2.21	0.41
1:B:123:GLU:OE2	1:B:135:LYS:NZ	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/251 (105%)	259 (98%)	4 (2%)	0	100	100
1	B	251/251 (100%)	243 (97%)	5 (2%)	3 (1%)	13	1
All	All	514/502 (102%)	502 (98%)	9 (2%)	3 (1%)	25	6

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	120	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	140	ALA
1	B	138	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/216 (107%)	229 (99%)	2 (1%)	78	53
1	B	219/216 (101%)	216 (99%)	3 (1%)	67	36
All	All	450/432 (104%)	445 (99%)	5 (1%)	71	45

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	LYS
1	A	193	LYS
1	B	138	THR
1	B	139	ARG
1	B	144	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	241	ASN
1	A	246	GLN
1	A	251	ASN
1	A	270	ASN
1	B	96	GLN
1	B	247	GLN
1	B	270	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/251 (99%)	0.74	20 (8%) 12 13	9, 14, 23, 29	0
1	B	246/251 (98%)	2.08	100 (40%) 0 0	11, 22, 29, 34	0
All	All	496/502 (98%)	1.40	120 (24%) 0 0	9, 16, 28, 34	0

All (120) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	119	LYS	10.0
1	B	120	SER	9.7
1	B	28	ASN	9.1
1	B	152	SER	7.5
1	B	127	ILE	7.2
1	B	131	ILE	7.1
1	B	173	ALA	7.1
1	B	109	LEU	7.0
1	B	110	VAL	6.7
1	B	118	ASP	6.4
1	B	138	THR	6.2
1	B	154	GLY	5.8
1	B	126	ILE	5.7
1	B	33	ASP	5.5
1	B	129	ASP	5.4
1	B	108	THR	5.4
1	B	128	ILE	5.3
1	B	139	ARG	5.0
1	B	206	SER	5.0
1	B	133	ILE	4.9
1	B	102	PHE	4.7
1	B	94	LEU	4.6
1	B	153	ASP	4.6
1	B	37	SER	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	35	PRO	4.4
1	A	81	ALA	4.3
1	B	132	ILE	4.2
1	B	93	ASP	4.2
1	B	150	ILE	4.2
1	B	121	SER	4.2
1	B	36	GLY	4.2
1	B	140	ALA	4.1
1	A	59[A]	ASP	4.0
1	B	29	SER	4.0
1	B	44	LYS	4.0
1	B	106	GLY	4.0
1	B	148	THR	3.9
1	B	156	GLY	3.9
1	B	155	SER	3.9
1	B	107	SER	3.8
1	B	174	GLU	3.7
1	B	79	VAL	3.7
1	B	105	ASP	3.7
1	B	30	VAL	3.6
1	B	101	VAL	3.5
1	B	149	GLY	3.5
1	B	83	ALA	3.5
1	B	32	VAL	3.5
1	B	137	ILE	3.5
1	A	25	HIS	3.5
1	B	59	ASP	3.5
1	B	194	SER	3.4
1	B	172	THR	3.4
1	B	136	ILE	3.4
1	B	92	ASP	3.3
1	B	82	ASP	3.3
1	B	116	SER	3.3
1	B	208	ALA	3.3
1	B	151	LYS	3.2
1	B	88	LEU	3.2
1	B	225	ILE	3.1
1	B	96	GLN	3.1
1	B	268	ILE	3.1
1	B	134	GLU	3.1
1	A	206	SER	3.0
1	B	61	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	117	LYS	3.0
1	B	114	VAL	3.0
1	B	147	TYR	3.0
1	B	78	GLY	2.9
1	B	62	GLU	2.9
1	B	199	VAL	2.9
1	B	86	VAL	2.8
1	B	124	GLU	2.8
1	B	91	SER	2.8
1	B	104	SER	2.8
1	B	58	VAL	2.8
1	B	111	SER	2.8
1	A	268	ILE	2.7
1	B	115	THR	2.7
1	B	193	LYS	2.6
1	B	113	LYS	2.6
1	A	114	VAL	2.6
1	A	82	ASP	2.6
1	A	208	ALA	2.6
1	A	27	LYS	2.6
1	B	158	ALA	2.6
1	B	49	ASP	2.5
1	B	130	GLY	2.5
1	B	171	LEU	2.5
1	B	122	THR	2.5
1	B	272	LEU	2.5
1	B	135	LYS	2.4
1	B	98	THR	2.4
1	B	125	LYS	2.4
1	B	31	SER	2.4
1	A	60	ALA	2.3
1	A	119	LYS	2.3
1	A	83	ALA	2.3
1	B	75	VAL	2.3
1	A	243	ILE	2.3
1	B	71	ASN	2.3
1	B	81	ALA	2.3
1	B	207	SER	2.2
1	B	103	LYS	2.2
1	B	99	LEU	2.2
1	B	141	ASP	2.2
1	A	110	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	225	ILE	2.2
1	A	180	VAL	2.2
1	A	55	ILE	2.2
1	A	165	TYR	2.1
1	B	271	ALA	2.1
1	A	137	ILE	2.1
1	B	215	ALA	2.1
1	B	157	LYS	2.1
1	B	209	ALA	2.1
1	B	196	GLU	2.0
1	A	120	SER	2.0
1	B	142	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.