



## Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 02:31 am BST

PDB ID : 4OLD  
Title : Crystal structure of AmpC beta-lactamase in complex with the product form of (6R,7R)-7-amino-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid  
Authors : Shoichet, B.K.; Barelier, S.  
Deposited on : 2014-01-23  
Resolution : 1.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4690 (1.50-1.46)
Clashscore	141614	4955 (1.50-1.46)
Ramachandran outliers	138981	4846 (1.50-1.46)
Sidechain outliers	138945	4844 (1.50-1.46)
RSRZ outliers	127900	4614 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	 4% 87% 9% . .
1	B	358	 % 89% 10% .

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6386 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	351	Total	C	N	O	S	0	16	0
			2834	1823	479	524	8			
1	B	358	Total	C	N	O	S	0	12	0
			2878	1848	489	534	7			

- Molecule 2 is (2R)-2-[(R)-amino(carboxy)methyl]-3,6-dihydro-2H-1,3-thiazine-4-carboxylic acid (three-letter code: 2UZ) (formula: C<sub>7</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	Total	C	N	O	S	0	1
			28	14	4	8	2		
2	B	1	Total	C	N	O	S	0	0
			14	7	2	4	1		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	1
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	1


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	292	Total O 292 292	0	15
4	B	325	Total O 325 325	0	13


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.45Å 76.42Å 97.88Å 90.00° 115.95° 90.00°	Depositor
Resolution (Å)	29.34 – 1.48 29.28 – 1.48	Depositor EDS
% Data completeness (in resolution range)	99.3 (29.34-1.48) 99.3 (29.28-1.48)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.01 (at 1.48Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.164 , 0.188 0.164 , 0.188	Depositor DCC
$R_{free}$ test set	4891 reflections (3.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.3	Xtrriage
Anisotropy	0.020	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 49.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6386	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2UZ, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.37	5/2913 (0.2%)	1.29	13/3985 (0.3%)
1	B	1.39	11/2958 (0.4%)	1.27	9/4046 (0.2%)
All	All	1.38	16/5871 (0.3%)	1.28	22/8031 (0.3%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	92	TYR	CZ-OH	9.96	1.54	1.37
1	A	219	GLU	CD-OE1	-8.04	1.16	1.25
1	B	87	ASP	CB-CG	6.48	1.65	1.51
1	B	92	TYR	CE2-CZ	6.38	1.46	1.38
1	B	171	GLU	CD-OE1	-6.18	1.18	1.25
1	B	95	GLU	CD-OE1	-5.98	1.19	1.25
1	B	203	TYR	CE2-CZ	5.87	1.46	1.38
1	B	312	TRP	CD2-CE2	5.79	1.48	1.41
1	B	203	TYR	CB-CG	-5.52	1.43	1.51
1	B	142	TRP	CD2-CE2	5.49	1.48	1.41
1	B	325	TYR	CD1-CE1	5.29	1.47	1.39
1	A	44	GLY	N-CA	-5.14	1.38	1.46
1	A	171	GLU	CD-OE2	-5.11	1.20	1.25
1	B	87	ASP	CG-OD2	-5.03	1.13	1.25
1	A	276	TRP	CD2-CE2	5.02	1.47	1.41
1	A	276	TRP	NE1-CE2	-5.01	1.31	1.37

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	87	ASP	CB-CG-OD2	-11.23	108.19	118.30
1	A	87	ASP	CB-CG-OD1	-9.83	109.45	118.30
1	B	133	ARG	NE-CZ-NH1	8.99	124.79	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	133	ARG	NE-CZ-NH2	-7.45	116.58	120.30
1	B	10	ASP	CB-CG-OD1	6.65	124.28	118.30
1	A	349	ARG	NE-CZ-NH1	-5.90	117.35	120.30
1	A	293	LEU	CB-CG-CD1	5.80	120.86	111.00
1	A	232	ARG	CG-CD-NE	-5.69	99.85	111.80
1	A	92	TYR	CB-CG-CD2	-5.68	117.59	121.00
1	A	351	ASP	CB-CG-OD2	-5.61	113.25	118.30
1	B	332	LYS	CD-CE-NZ	-5.51	99.02	111.70
1	B	41	PHE	CG-CD2-CE2	-5.45	114.81	120.80
1	A	10	ASP	CB-CG-OD1	5.42	123.18	118.30
1	B	232	ARG	NE-CZ-NH1	-5.42	117.59	120.30
1	A	45	TYR	CD1-CE1-CZ	-5.38	114.95	119.80
1	B	177	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	217	ASP	CB-CG-OD2	-5.25	113.58	118.30
1	A	265[A]	MET	CG-SD-CE	-5.18	91.91	100.20
1	A	265[B]	MET	CG-SD-CE	-5.18	91.91	100.20
1	A	96	LEU	CB-CG-CD2	-5.17	102.22	111.00
1	A	349	ARG	NE-CZ-NH2	-5.05	117.78	120.30
1	B	332	LYS	CA-CB-CG	5.03	124.47	113.40

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2834	0	2756	27	0
1	B	2878	0	2815	19	0
2	A	28	0	16	2	0
2	B	14	0	8	0	0
3	A	10	0	0	0	0
3	B	5	0	0	0	0
4	A	292	0	0	6	0
4	B	325	0	0	3	0
All	All	6386	0	5595	48	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:401[B]:2UZ:SAI	2:A:401[B]:2UZ:O	2.20	0.99
1:A:28[B]:MET:HG3	1:A:29:ALA:N	2.04	0.71
1:B:87:ASP:OD2	1:B:92:TYR:OH	2.12	0.68
1:B:262[B]:THR:CG2	1:B:298:VAL:HG12	2.24	0.67
1:A:243:ILE:CD1	1:A:252:ILE:HD12	2.29	0.62
1:A:278[B]:VAL:HG21	1:A:283:ILE:HD11	1.82	0.62
1:A:262:THR:HG22	4:A:730:HOH:O	2.00	0.59
1:A:243:ILE:CD1	1:A:252:ILE:CD1	2.80	0.59
1:A:265[A]:MET:SD	4:A:730:HOH:O	2.58	0.56
1:A:262:THR:O	1:A:265[A]:MET:HG3	2.06	0.55
1:A:175[B]:GLN:HG3	1:A:180:GLN:OE1	2.08	0.54
1:A:33:ILE:HD12	1:A:33:ILE:N	2.23	0.53
1:A:243:ILE:HD12	1:A:252:ILE:HD12	1.91	0.53
1:B:296[B]:ARG:NH1	4:B:646:HOH:O	2.37	0.53
1:A:276:TRP:CE3	1:A:278[A]:VAL:HG12	2.44	0.52
1:A:87:ASP:OD2	1:A:92:TYR:OH	2.21	0.52
1:B:243:ILE:CD1	1:B:252:ILE:HD12	2.40	0.52
1:A:334:LEU:HG	1:A:357:LEU:HD22	1.93	0.51
1:A:262:THR:HG23	1:A:265[A]:MET:HG3	1.94	0.49
1:A:316[A]:THR:HG22	4:A:534:HOH:O	2.14	0.48
1:B:26:PRO:HB3	1:B:48:ILE:HD11	1.95	0.48
1:A:262:THR:CG2	1:A:265[A]:MET:SD	3.02	0.47
1:A:119:LEU:HA	1:A:151:ALA:HA	1.97	0.46
1:B:276:TRP:CD2	1:B:277:PRO:HA	2.51	0.46
1:B:262[B]:THR:HG23	1:B:298:VAL:HG12	1.95	0.46
1:A:344:TYR:CZ	1:A:349:ARG:HG2	2.51	0.45
1:B:334:LEU:HG	1:B:357:LEU:HD22	1.98	0.45
2:A:401[B]:2UZ:H6	4:A:766:HOH:O	2.15	0.44
1:B:243:ILE:CD1	1:B:252:ILE:CD1	2.95	0.44
1:A:276:TRP:CD2	1:A:277:PRO:HA	2.52	0.44
1:A:262:THR:HG23	1:A:265[A]:MET:CG	2.48	0.43
1:A:17:THR:HB	1:A:18:PRO:HD3	2.00	0.43
1:B:7[A]:GLN:HG2	1:B:8:ILE:N	2.33	0.43
1:B:33:ILE:HD12	1:B:33:ILE:N	2.34	0.42
1:A:240:PRO:O	1:A:249:GLN:HG3	2.19	0.42
1:A:204:ARG:NH2	4:A:546:HOH:O	2.52	0.42
1:A:262:THR:HG22	1:A:265[A]:MET:SD	2.60	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:265[A]:MET:HG3	4:B:504:HOH:O	2.20	0.42
1:A:256:GLN:HG3	4:A:639:HOH:O	2.19	0.42
1:B:262[B]:THR:HG22	1:B:298:VAL:HG12	1.99	0.42
1:B:344:TYR:CE1	1:B:349:ARG:HG2	2.55	0.41
1:B:278[A]:VAL:HG22	1:B:354:TRP:CZ3	2.55	0.41
1:A:125:VAL:HG11	1:A:131[A]:LEU:HD23	2.03	0.41
1:B:262[B]:THR:CG2	1:B:298:VAL:CG1	2.96	0.41
1:B:290:LYS:NZ	4:B:700:HOH:O	2.52	0.41
1:B:120[B]:GLN:NE2	1:B:152:ASN:HD22	2.17	0.41
1:A:293:LEU:HD23	1:A:293:LEU:C	2.41	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/358 (101%)	358 (99%)	5 (1%)	0	100	100
1	B	369/358 (103%)	363 (98%)	6 (2%)	0	100	100
All	All	732/716 (102%)	721 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/292 (100%)	290 (99%)	2 (1%)	84	68
1	B	299/292 (102%)	295 (99%)	4 (1%)	69	42
All	All	591/584 (101%)	585 (99%)	6 (1%)	78	54

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	314	HIS
1	A	351	ASP
1	B	262[A]	THR
1	B	262[B]	THR
1	B	309	ARG
1	B	314	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	B	402[B]	-	4,4,4	1.51	1 (25%)	6,6,6	1.20	1 (16%)
3	PO4	A	402[B]	-	4,4,4	1.80	1 (25%)	6,6,6	0.48	0
2	2UZ	B	401	-	2,14,14	3.08	1 (50%)	1,19,19	3.08	1 (100%)
2	2UZ	A	401[A]	-	2,14,14	5.49	2 (100%)	1,19,19	2.21	1 (100%)
3	PO4	A	403	-	4,4,4	0.47	0	6,6,6	1.47	1 (16%)
2	2UZ	A	401[B]	-	2,14,14	3.67	1 (50%)	1,19,19	1.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2UZ	B	401	-	-	0/0/22/22	0/0/1/1
2	2UZ	A	401[A]	-	-	0/0/22/22	0/0/1/1
2	2UZ	A	401[B]	-	-	0/0/22/22	0/0/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401[A]	2UZ	CAJ-CAL	-7.22	1.41	1.52
2	A	401[B]	2UZ	CAJ-CAL	-5.17	1.44	1.52
2	B	401	2UZ	CAJ-CAL	-4.33	1.45	1.52
2	A	401[A]	2UZ	CA-N	2.86	1.53	1.47
3	B	402[B]	PO4	P-O3	-2.39	1.47	1.54
3	A	402[B]	PO4	P-O3	-2.27	1.47	1.54

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	2UZ	CAG-SAI-CB	-3.08	88.45	94.45
3	A	403	PO4	O3-P-O2	2.30	115.34	107.97
2	A	401[A]	2UZ	CAG-SAI-CB	2.21	98.75	94.45
3	B	402[B]	PO4	O4-P-O2	2.07	114.61	107.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401[B]	2UZ	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	351/358 (98%)	0.08	16 (4%) 32 35	10, 17, 33, 48	0
1	B	358/358 (100%)	-0.15	3 (0%) 86 88	10, 16, 28, 48	0
All	All	709/716 (99%)	-0.04	19 (2%) 54 58	10, 16, 30, 48	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	ASN	4.6
1	A	278[A]	VAL	3.8
1	B	244	ASN	3.8
1	A	281	ASP	3.2
1	A	126	LYS	3.0
1	A	34	TYR	2.9
1	B	241	LEU	2.8
1	B	205	GLU	2.7
1	A	242	ASP	2.6
1	A	241	LEU	2.5
1	A	244	ASN	2.5
1	A	283	ILE	2.5
1	A	269	LEU	2.4
1	A	35	GLN	2.4
1	A	123[A]	ASP	2.3
1	A	102	ASN	2.3
1	A	4	ALA	2.1
1	A	282[A]	SER	2.1
1	A	24	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	2UZ	A	401[A]	14/14	0.85	0.20	23,32,36,41	14
2	2UZ	A	401[B]	14/14	0.85	0.20	19,37,49,49	14
2	2UZ	B	401	14/14	0.87	0.17	24,58,83,91	0
3	PO4	A	403	5/5	0.91	0.13	27,27,33,36	5
3	PO4	B	402[B]	5/5	0.95	0.11	18,19,21,22	5
3	PO4	A	402[B]	5/5	0.97	0.16	14,14,16,18	5

### 6.5 Other polymers [i](#)

There are no such residues in this entry.