wwPDB X-ray Structure Validation Summary Report (i)

Jan 8, 2024 – 01:07 pm GMT

PDB ID	:	5OMX
Title	:	X-ray Structure of the H2A-N38C Nucleosome Core Particle
Authors	:	Frouws, T.D.; Richmond, T.J.
Deposited on	:	2017-08-02
Resolution	:	2.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	I	147	11%, 27%, 73%
2	J	147	11%, 28%, 72%
3	A	135	.%, 60%, 11%, •, 28%
3	E	135	.%, 66%, 7%, 27%
4	B	103	3%, 60%, 17%, 23%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	103	3% 72% 12% 17%
5	C	129	3% 73% 9% 19%
5	G	129	2% 64% 16% 20%
6	D	122	% 66% 10% • 23%
6	H	122	% 63% 15% 22%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	MN	J	101	-	-	-	X

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 12250 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (147-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	I	147	3011	1440	546	879	146	0	0	0

- Molecule 2 is a DNA chain called DNA (147-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	J	147	3010	1440	543	881	146	0	0	0

- Molecule 3 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	97	801	506	155	138	2	0	0	0
			Total	C	N	O	S			
3	E	99	816	515	158	141	2	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	GLY	conflict	UNP P84233
A	110	ALA	CYS	engineered mutation	UNP P84233
E	102	ALA	GLY	conflict	UNP P84233
E	110	ALA	CYS	engineered mutation	UNP P84233

- Molecule 4 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	B	79	627	395	121	110	1	0	0	0
			Total	C	N	O	S			
4	F	86	694	436	140	117	1	0	0	0

- Molecule 5 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	C	105	807	509	157	140	1	0	0	0
5	G	103	793	500	154	138	1	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	38	CYS	ASN	engineered mutation	UNP Q6AZJ8
G	38	CYS	ASN	engineered mutation	UNP Q6AZJ8

- Molecule 6 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	D	94	736	463	132	139	2	0	0	0
6	H	95	746	469	136	139	2	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	32	THR	SER	conflict	UNP P02281
H	32	THR	SER	conflict	UNP P02281

- Molecule 7 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
7	I	19	19	19	0	0
7	J	13	13	13	0	0
7	E	1	1	1	0	0

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
8	A	1	1	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
8	C	1	1	1	0	0
			Total	Cl		
8	E	1	1	1	0	0
			Total	Cl		
8	G	1	1	1	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
9	I	17	17	17	0	0
			Total	O		
9	J	13	13	13	0	0
			Total	O		
9	A	7	7	7	0	0
			Total	O		
9	B	14	14	14	0	0
			Total	O		
9	C	25	25	25	0	0
			Total	O		
9	D	14	14	14	0	0
			Total	O		
9	E	34	34	34	0	0
			Total	O		
9	F	27	27	27	0	0
			Total	O		
9	G	15	15	15	0	0
			Total	O		
9	H	6	6	6	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (147-MER)

Chain I:



- Molecule 2: DNA (147-MER)

Chain J:



- Molecule 3: Histone H3.2

Chain A:



- Molecule 3: Histone H3.2

Chain E:



- Molecule 4: Histone H4

Chain B:



- Molecule 4: Histone H4

Chain F:



- Molecule 5: Histone H2A

Chain C:



- Molecule 5: Histone H2A

Chain G:



- Molecule 6: Histone H2B 1.1

Chain D:



- Molecule 6: Histone H2B 1.1

Chain H:

E113
A124
LYS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	106.75Å 182.58Å 109.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.14 – 2.32 47.12 – 2.32	Depositor EDS
% Data completeness (in resolution range)	86.7 (47.14-2.32) 86.4 (47.12-2.32)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.81 (at 2.32Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.222 , 0.259 0.220 , 0.258	Depositor DCC
R_{free} test set	4048 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å^2)	49.2	Xtriage
Anisotropy	0.511	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.29 , 49.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.46$, $< L^2 > = 0.28$	Xtriage
Estimated twinning fraction	0.027 for l,-k,h	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	12250	wwPDB-VP
Average B, all atoms (Å^2)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.45% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	I	0.43	0/3378	0.84	0/5212
2	J	0.43	0/3376	0.85	0/5209
3	A	0.34	0/813	0.54	0/1091
3	E	0.45	0/828	0.58	0/1110
4	B	0.39	0/634	0.58	0/848
4	F	0.43	0/702	0.62	0/937
5	C	0.40	0/817	0.58	0/1103
5	G	0.32	0/803	0.55	0/1085
6	D	0.42	0/747	0.59	0/1004
6	H	0.34	0/757	0.56	0/1018
All	All	0.41	0/12855	0.74	0/18617

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	3011	0	1662	113	0
2	J	3010	0	1663	120	0
3	A	801	0	841	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	816	0	858	6	0
4	B	627	0	663	17	0
4	F	694	0	742	12	0
5	C	807	0	862	9	0
5	G	793	0	844	19	0
6	D	736	0	760	11	0
6	H	746	0	773	15	0
7	E	1	0	0	0	0
7	I	19	0	0	0	0
7	J	13	0	0	0	0
8	A	1	0	0	0	0
8	C	1	0	0	0	0
8	E	1	0	0	0	0
8	G	1	0	0	0	0
9	A	7	0	0	0	0
9	B	14	0	0	0	0
9	C	25	0	0	0	0
9	D	14	0	0	0	0
9	E	34	0	0	0	0
9	F	27	0	0	0	0
9	G	15	0	0	0	0
9	H	6	0	0	0	0
9	I	17	0	0	0	0
9	J	13	0	0	0	0
All	All	12250	0	9668	306	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 306 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:24:DG:H2”	2:J:25:DG:H5’	1.11	1.10
2:J:24:DG:C2’	2:J:25:DG:H5’	1.96	0.95
2:J:-54:DT:H2”	2:J:-53:DA:H5’	1.48	0.94
2:J:56:DC:H2”	2:J:57:DT:H5’	1.48	0.93
2:J:63:DT:H2”	2:J:64:DG:H5’	1.56	0.88

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	95/135 (70%)	93 (98%)	2 (2%)	0	100	100
3	E	97/135 (72%)	96 (99%)	1 (1%)	0	100	100
4	B	77/103 (75%)	75 (97%)	2 (3%)	0	100	100
4	F	84/103 (82%)	82 (98%)	2 (2%)	0	100	100
5	C	103/129 (80%)	100 (97%)	3 (3%)	0	100	100
5	G	101/129 (78%)	99 (98%)	2 (2%)	0	100	100
6	D	92/122 (75%)	91 (99%)	0	1 (1%)	14	15
6	H	93/122 (76%)	91 (98%)	1 (1%)	1 (1%)	14	15
All	All	742/978 (76%)	727 (98%)	13 (2%)	2 (0%)	41	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
6	D	104	GLY
6	H	104	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	84/109 (77%)	83 (99%)	1 (1%)	71	83
3	E	85/109 (78%)	85 (100%)	0	100	100
4	B	64/79 (81%)	64 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	F	71/79 (90%)	71 (100%)	0	100	100
5	C	83/101 (82%)	81 (98%)	2 (2%)	49	65
5	G	82/101 (81%)	80 (98%)	2 (2%)	49	65
6	D	80/102 (78%)	77 (96%)	3 (4%)	33	46
6	H	81/102 (79%)	79 (98%)	2 (2%)	47	64
All	All	630/782 (81%)	620 (98%)	10 (2%)	62	77

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	G	81	ARG
6	H	36	SER
6	H	57	LYS
6	D	34	LYS
6	D	57	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
6	H	67	ASN
5	G	82	HIS
6	D	95	GLN
6	D	49	HIS
4	F	93	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 37 ligands modelled in this entry, 37 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	I	147/147 (100%)	0.81	16 (10%)	5	8	52, 98, 139, 156	0
2	J	147/147 (100%)	0.75	16 (10%)	5	8	54, 98, 146, 166	0
3	A	97/135 (71%)	0.48	2 (2%)	63	70	32, 50, 79, 105	0
3	E	99/135 (73%)	0.68	2 (2%)	65	72	22, 40, 75, 117	0
4	B	79/103 (76%)	0.61	3 (3%)	40	47	31, 47, 78, 121	0
4	F	86/103 (83%)	0.78	3 (3%)	44	51	26, 37, 72, 145	0
5	C	105/129 (81%)	0.62	4 (3%)	40	47	25, 46, 75, 99	0
5	G	103/129 (79%)	0.46	2 (1%)	66	73	33, 55, 86, 114	0
6	D	94/122 (77%)	0.66	1 (1%)	80	85	27, 45, 78, 133	0
6	H	95/122 (77%)	0.67	1 (1%)	80	85	37, 55, 91, 112	0
All	All	1052/1272 (82%)	0.66	50 (4%)	30	38	22, 55, 124, 166	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	135	ALA	9.4
4	F	17	ARG	4.8
1	I	-18	DA	3.8
3	A	39	HIS	3.8
2	J	22	DA	3.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
7	MN	J	102	1/1	-0.01	0.36	155,155,155,155	0
7	MN	J	113	1/1	0.07	0.19	156,156,156,156	0
7	MN	I	102	1/1	0.16	0.17	166,166,166,166	0
7	MN	J	106	1/1	0.23	0.17	172,172,172,172	0
7	MN	I	107	1/1	0.27	0.33	166,166,166,166	0
7	MN	J	101	1/1	0.28	0.53	179,179,179,179	0
7	MN	J	112	1/1	0.29	0.19	135,135,135,135	0
7	MN	I	117	1/1	0.40	0.12	126,126,126,126	0
7	MN	J	107	1/1	0.42	0.09	164,164,164,164	0
7	MN	J	104	1/1	0.43	0.13	152,152,152,152	0
7	MN	J	105	1/1	0.46	0.13	132,132,132,132	0
7	MN	I	108	1/1	0.50	0.11	157,157,157,157	0
7	MN	I	104	1/1	0.50	0.22	153,153,153,153	0
7	MN	I	110	1/1	0.60	0.10	130,130,130,130	0
7	MN	I	109	1/1	0.61	0.18	138,138,138,138	0
7	MN	I	101	1/1	0.63	0.14	143,143,143,143	0
7	MN	I	105	1/1	0.70	0.21	152,152,152,152	0
7	MN	I	103	1/1	0.72	0.28	152,152,152,152	0
7	MN	J	103	1/1	0.78	0.07	142,142,142,142	0
7	MN	I	111	1/1	0.82	0.08	118,118,118,118	0
7	MN	J	109	1/1	0.84	0.09	77,77,77,77	0
7	MN	I	119	1/1	0.84	0.18	115,115,115,115	0
7	MN	I	116	1/1	0.84	0.12	114,114,114,114	0
7	MN	I	112	1/1	0.88	0.06	135,135,135,135	0
7	MN	I	113	1/1	0.89	0.19	96,96,96,96	0
7	MN	I	114	1/1	0.92	0.19	89,89,89,89	0
7	MN	J	110	1/1	0.92	0.16	86,86,86,86	0
8	CL	E	202	1/1	0.94	0.10	69,69,69,69	0
8	CL	G	201	1/1	0.94	0.19	72,72,72,72	0
7	MN	I	106	1/1	0.95	0.04	109,109,109,109	0
7	MN	I	118	1/1	0.95	0.09	113,113,113,113	0
7	MN	I	115	1/1	0.96	0.16	88,88,88,88	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
8	CL	A	201	1/1	0.97	0.12	69,69,69,69	0
8	CL	C	201	1/1	0.98	0.18	56,56,56,56	0
7	MN	J	111	1/1	0.98	0.11	83,83,83,83	0
7	MN	J	108	1/1	0.98	0.18	71,71,71,71	0
7	MN	E	201	1/1	1.00	0.20	37,37,37,37	0

6.5 Other polymers (i)

There are no such residues in this entry.