



# Full wwPDB X-ray Structure Validation Report i

Feb 10, 2022 – 06:24 pm GMT

PDB ID : 7OMG  
Title : Crystal structure of KOD DNA Polymerase in a ternary complex with an Uracil containing template  
Authors : Betz, K.; Kropp, H.M.; Diederichs, K.; Marx, A.  
Deposited on : 2021-05-22  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.26
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 6906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	747	Total	C 6126	N 3946	O 1034	S 1130	16	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	141	ALA	ASP	engineered mutation	UNP P77933
A	143	ALA	GLU	engineered mutation	UNP P77933


- Molecule 2 is a DNA chain called Template.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	20	Total	C 409	N 194	O 72	P 123	20	0	0

- Molecule 3 is a DNA chain called Primer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	12	Total	C 239	N 114	O 48	P 66	11	0	0

- Molecule 4 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	30	10	5	12	3	0	0


- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total Mg 2		0	0

- Molecule 6 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total Mn 1		0	0

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 4 2 2	0	0
7	A	1	Total C O 4 2 2	0	0

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	2	Total Ca 2 2	0	0

- Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	T	1	Total Cl 1 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	81	Total O 81 81	0	0
10	T	6	Total O 6 6	0	0
10	P	1	Total O 1 1	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase



- Molecule 2: Template



- Molecule 3: Primer



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	108.40 Å    147.14 Å    71.13 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	46.25 – 2.10 46.25 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.9 (46.25-2.10) 98.9 (46.25-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	0.97 (at 2.10 Å)	Xtriage
Refinement program	PHENIX 1.19_4085, PHENIX 1.19_4085	Depositor
$R$ , $R_{free}$	0.210 , 0.248 0.208 , 0.246	Depositor DCC
$R_{free}$ test set	3179 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.5	Xtriage
Anisotropy	0.261	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6906	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, DOC, CL, EDO, MG, CA, DTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/6263	0.64	0/8452
2	T	1.90	8/457 (1.8%)	1.45	6/703 (0.9%)
3	P	0.88	0/248	0.79	0/380
All	All	0.67	8/6968 (0.1%)	0.74	6/9535 (0.1%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	4	DU	N1-C2	26.27	1.62	1.38
2	T	4	DU	C3'-O3'	-12.00	1.28	1.44
2	T	4	DU	C3'-C2'	10.79	1.65	1.52
2	T	4	DU	C4-C5	8.28	1.51	1.43
2	T	4	DU	C5-C6	8.28	1.41	1.34
2	T	4	DU	C2-N3	-7.64	1.32	1.37
2	T	5	DC	C1'-N1	5.18	1.55	1.49
2	T	4	DU	C5'-C4'	5.11	1.56	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	4	DU	N1-C2-O2	-15.61	111.88	122.80
2	T	4	DU	O5'-P-OP1	12.53	125.73	110.70
2	T	4	DU	C6-N1-C2	-10.32	114.81	121.00
2	T	4	DU	N3-C2-O2	9.59	128.91	122.20
2	T	4	DU	N1-C2-N3	7.11	119.17	114.90
2	T	3	DT	OP1-P-O3'	-5.89	92.25	105.20

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6126	0	6170	66	0
2	T	409	0	225	3	0
3	P	239	0	134	0	0
4	A	30	0	12	2	0
5	A	2	0	0	0	0
6	A	1	0	0	0	0
7	A	8	0	12	0	0
8	A	2	0	0	0	0
9	T	1	0	0	1	0
10	A	81	0	0	1	0
10	P	1	0	0	0	0
10	T	6	0	0	0	0
All	All	6906	0	6553	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:287:LYS:NZ	1:A:315:ASP:OD1	2.21	0.71
1:A:50:GLU:HG2	1:A:103:HIS:HE2	1.56	0.70
1:A:329:MET:HA	1:A:482:ARG:HD2	1.74	0.70
1:A:145:LEU:HB2	1:A:158:LEU:HD21	1.78	0.65
1:A:263:VAL:O	1:A:267:THR:HG23	1.96	0.64
1:A:299:TRP:HA	1:A:305:LEU:HD21	1.78	0.64
1:A:258:PHE:HE1	1:A:263:VAL:HG21	1.65	0.62
1:A:428:CYS:HB2	1:A:431:TYR:CZ	2.37	0.60
1:A:137:MET:HE3	1:A:327:LEU:HD22	1.86	0.57
1:A:159:MET:HG2	1:A:172:THR:HB	1.86	0.57
2:T:5:DC:H2'	9:T:101:CL:CL	2.41	0.57
1:A:419:SER:OG	1:A:421:ASP:OD1	2.18	0.57
1:A:467:MET:O	10:A:901:HOH:O	2.17	0.56
1:A:597:ILE:HD11	1:A:601:GLY:HA2	1.87	0.56
1:A:626:VAL:HG13	1:A:639:ALA:HB1	1.86	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:585:ARG:NH2	1:A:631:LEU:O	2.40	0.55
1:A:176:VAL:HG13	1:A:306:GLU:HG3	1.87	0.55
1:A:625:ARG:NH2	1:A:645:GLU:OE2	2.40	0.55
1:A:610:ILE:HG13	1:A:623:GLN:OE1	2.07	0.54
1:A:625:ARG:HB3	1:A:642:ILE:HD13	1.89	0.53
1:A:409:TYR:CG	4:A:801:DTP:H2'1	2.44	0.52
1:A:70:LYS:HD2	1:A:81:GLU:OE1	2.10	0.52
1:A:648:GLU:O	1:A:649:LYS:HB2	2.09	0.52
1:A:665:GLN:HG2	1:A:666:ILE:O	2.11	0.50
1:A:671:LYS:H	1:A:671:LYS:HD2	1.76	0.50
1:A:1:MET:HB3	1:A:130:GLU:O	2.11	0.50
1:A:409:TYR:CD2	4:A:801:DTP:H2'1	2.47	0.49
1:A:174:LYS:HB2	1:A:299:TRP:CZ3	2.48	0.49
1:A:525:ILE:HG23	1:A:536:VAL:HG21	1.94	0.49
1:A:293:GLU:HA	1:A:296:THR:HG22	1.95	0.49
1:A:633:ASP:OD1	1:A:633:ASP:N	2.46	0.48
1:A:325:GLU:HB3	1:A:478:LEU:HD11	1.96	0.47
1:A:429:LYS:HE3	1:A:429:LYS:HB2	1.65	0.47
1:A:50:GLU:HG2	1:A:103:HIS:NE2	2.27	0.47
1:A:603:ILE:HG22	1:A:605:THR:HG22	1.97	0.46
1:A:649:LYS:HB3	1:A:655:VAL:HG23	1.98	0.46
1:A:660:LEU:HB3	1:A:732:TYR:CZ	2.50	0.46
1:A:137:MET:HE2	1:A:207:ILE:HD11	1.97	0.46
1:A:391:GLU:HG3	1:A:394:ARG:HH12	1.81	0.46
1:A:310:ARG:NH1	1:A:314:GLU:OE2	2.48	0.45
1:A:57:GLU:HG2	1:A:58:ARG:H	1.81	0.45
1:A:416:HIS:NE2	1:A:520:TYR:OH	2.44	0.45
1:A:648:GLU:HG2	1:A:652:LYS:HD2	1.98	0.45
1:A:480:ASP:OD2	1:A:484:ARG:NH2	2.49	0.45
1:A:394:ARG:HH21	1:A:590:THR:HG22	1.81	0.45
1:A:396:LEU:HD11	1:A:585:ARG:HD2	1.99	0.45
1:A:72:GLN:HG2	1:A:81:GLU:HG3	1.99	0.44
1:A:49:GLU:O	1:A:53:LYS:HG3	2.17	0.44
2:T:19:DG:H2"	2:T:20:DT:H5'	1.99	0.44
1:A:16:ILE:HB	1:A:31:ASP:HB3	2.00	0.43
1:A:527:GLU:CG	1:A:566:TYR:HE2	2.31	0.43
1:A:391:GLU:OE1	1:A:591:LYS:HE3	2.18	0.43
1:A:627:LEU:HD23	1:A:627:LEU:HA	1.87	0.43
1:A:495:GLY:HA3	2:T:9:DT:O2	2.18	0.43
1:A:173:TRP:CH2	1:A:299:TRP:HZ2	2.37	0.42
1:A:737:VAL:O	1:A:741:VAL:HG22	2.19	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338:GLY:O	1:A:359:ARG:NH2	2.50	0.42
1:A:509:CYS:O	1:A:513:VAL:HG23	2.21	0.41
1:A:531:LYS:HE2	1:A:531:LYS:HB3	1.77	0.41
1:A:174:LYS:HB2	1:A:299:TRP:CH2	2.56	0.41
1:A:332:GLN:OE1	1:A:482:ARG:NH2	2.52	0.41
1:A:608:LEU:O	1:A:611:VAL:HG22	2.21	0.41
1:A:484:ARG:HE	1:A:484:ARG:HB2	1.48	0.41
1:A:258:PHE:CE1	1:A:263:VAL:HG21	2.51	0.40
1:A:78:ARG:HA	1:A:79:PRO:HD3	1.90	0.40
1:A:153:ALA:HA	1:A:218:TYR:CZ	2.56	0.40
1:A:188:ARG:HG3	1:A:228:ILE:HD11	2.02	0.40
1:A:419:SER:HB2	1:A:450:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	745/774 (96%)	717 (96%)	27 (4%)	1 (0%)	51 54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	649	LYS

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	649/672 (97%)	630 (97%)	19 (3%)	42    46

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	THR
1	A	215	ASP
1	A	244	MET
1	A	251	GLU
1	A	267	THR
1	A	274	THR
1	A	382	GLN
1	A	476	ARG
1	A	482	ARG
1	A	494	TYR
1	A	516	TRP
1	A	576	GLU
1	A	580	GLU
1	A	600	GLU
1	A	605	THR
1	A	630	LEU
1	A	633	ASP
1	A	713	ARG
1	A	746	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	DOC	P	12	2,3	14,19,20	2.65	5 (35%)	13,26,29	1.67	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DOC	P	12	2,3	-	0/4/18/19	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	12	DOC	C2-N3	5.45	1.49	1.38
3	P	12	DOC	C5-C4	5.14	1.53	1.41
3	P	12	DOC	C6-N1	3.88	1.40	1.35
3	P	12	DOC	C4-N4	2.96	1.43	1.35
3	P	12	DOC	C4-N3	-2.91	1.30	1.35

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	12	DOC	C4-N3-C2	4.59	120.99	116.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	DTP	A	801	6,5	26,32,32	0.60	0	30,50,50	0.91	1 (3%)
7	EDO	A	805	-	3,3,3	0.80	0	2,2,2	0.38	0
7	EDO	A	806	-	3,3,3	0.62	0	2,2,2	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTP	A	801	6,5	-	1/18/34/34	0/3/3/3
7	EDO	A	805	-	-	0/1/1/1	-
7	EDO	A	806	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	A	801	DTP	C5-C6-N6	2.42	124.04	120.35

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	801	DTP	PB-O3B-PG-O3G

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	801	DTP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	747/774 (96%)	0.71	69 (9%) 9 11	42, 69, 109, 156	0
2	T	19/21 (90%)	0.05	0 100 100	47, 77, 95, 126	0
3	P	11/12 (91%)	-0.17	0 100 100	54, 77, 89, 89	0
All	All	777/807 (96%)	0.68	69 (8%) 9 12	42, 70, 109, 156	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	299	TRP	9.1
1	A	292	ALA	8.7
1	A	289	LYS	7.1
1	A	723	THR	6.8
1	A	291	TYR	6.6
1	A	246	ASP	6.3
1	A	741	VAL	5.8
1	A	297	THR	5.6
1	A	301	THR	5.5
1	A	290	VAL	5.5
1	A	646	VAL	5.3
1	A	286	PRO	5.1
1	A	745	LEU	5.1
1	A	293	GLU	4.6
1	A	244	MET	4.5
1	A	675	ALA	4.4
1	A	471	ILE	4.3
1	A	243	ARG	4.2
1	A	302	GLY	4.1
1	A	692	LYS	4.0
1	A	296	THR	3.9
1	A	691	VAL	3.9
1	A	300	GLU	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	671	LYS	3.7
1	A	650	LEU	3.7
1	A	337	ILE	3.6
1	A	248	PHE	3.5
1	A	168	ALA	3.5
1	A	707	SER	3.3
1	A	303	GLU	3.2
1	A	643	VAL	3.2
1	A	294	GLU	3.1
1	A	105	ALA	3.0
1	A	726	LYS	2.9
1	A	717	PHE	2.9
1	A	305	LEU	2.8
1	A	648	GLU	2.7
1	A	548	ILE	2.6
1	A	374	GLU	2.6
1	A	642	ILE	2.6
1	A	456	LEU	2.6
1	A	288	GLU	2.5
1	A	554	GLU	2.5
1	A	747	ALA	2.5
1	A	247	ARG	2.5
1	A	285	GLN	2.5
1	A	295	ILE	2.4
1	A	708	GLY	2.4
1	A	661	VAL	2.4
1	A	689	ARG	2.4
1	A	375	LYS	2.4
1	A	394	ARG	2.4
1	A	688	ALA	2.4
1	A	147	HIS	2.3
1	A	715	ILE	2.3
1	A	47	ALA	2.3
1	A	658	GLU	2.3
1	A	731	TYR	2.2
1	A	693	ILE	2.2
1	A	674	LYS	2.2
1	A	636	VAL	2.2
1	A	61	THR	2.1
1	A	676	THR	2.1
1	A	739	PRO	2.0
1	A	60	GLY	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	690	GLY	2.0
1	A	742	GLU	2.0
1	A	635	ASP	2.0
1	A	357	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	DOC	P	12	18/19	0.96	0.19	48,50,55,55	0

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	CA	A	807	1/1	0.80	0.08	88,88,88,88	0
7	EDO	A	805	4/4	0.87	0.25	58,58,58,61	0
9	CL	T	101	1/1	0.87	0.16	82,82,82,82	0
7	EDO	A	806	4/4	0.88	0.16	64,65,65,67	0
5	MG	A	803	1/1	0.91	0.13	45,45,45,45	0
5	MG	A	802	1/1	0.94	0.12	57,57,57,57	0
4	DTP	A	801	30/30	0.96	0.16	44,48,53,54	0
8	CA	A	808	1/1	0.97	0.09	79,79,79,79	0
6	MN	A	804	1/1	0.99	0.15	51,51,51,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.