

Jan 20, 2024 - 07:44 am GMT

PDB ID	:	80PC
EMDB ID	:	EMD-17048
Title	:	Virus-like Particle based on PVY coat protein with helical architecture encap-
		sidating ssRNA
Authors	:	Kavcic, L.; Kezar, A.; Podobnik, M.
Deposited on		
Resolution	:	2.99 Å(reported)
Based on initial model	:	6HXZ

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev70
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f EM\ structures}\ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for $\geq=3, 2, 1$ and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq=5\%$ The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	Aa	267	84%	15%
1	Ac	267	84%	15%
1	Ae	267	84%	15%
1	Ag	267	84%	15%
1	Ai	267	84%	15%
1	Ak	267	84%	15%
1	Am	267	84%	15%



Chain Length Quality of chain Mol 2671 Ao 84% 15% 1 Aq 26784% 15% · 1 As 26784% 15% 2671 Au 84% 15% • 1 Aw 26784% 15% Ay 2671 84% 15% Ba 2671 84% 15% Bc 2671 84% 15% Be 2671 84% 15% 1 Bg 26784% 15% 1 Bi 26784% 15% ٠ Bk 2671 84% 15% 1 Bm 26784% 15% Во 2671 84% 15% Bq 2671 84% 15% • 1 Bs 26784% 15% Bu 2671 84% 15% 2671 Bw 84% 15% • 1 By 26784% 15% Ca2671 84% 15% 1 $\mathbf{C}\mathbf{c}$ 26784% 15% 2 Ab 5100% 2 Ad 5100% 2Af 5100% 2 Ah 5100%

Continued from previous page...



Chain Length Quality of chain Mol 25Aj 100% 2Al 5100% 2An 5100% 25Ap 100% 25Ar 100% 2 At 5100% 2Av 5100% 2 $\mathbf{A}\mathbf{x}$ 5100% 25 Az 100% 2 Bb 5100% $\mathbf{2}$ Bd 5100% Bf 25100% 2Bh5100% 2Вj 5100% 2Bl 5100% 2 Bn 5100% $\mathbf{2}$ Bp 5100% 2 Br 5100% 2Bt 5100% 25Bv 100% 25Bx 100% 2 Bz 5100% $\mathbf{2}$ Cb 5100% 2 Cd 5100%

Continued from previous page...



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 53088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	Mol	Chain	Residues		At	AltConf	Trace			
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	1	Δο	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		Aa	220	1796	1125	315	342	14	0	0
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	1	Δο	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		AC	220	1796	1125	315	342	14	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	Δο	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		Ae	220	1796			342		0	0
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	1	Δσ	226	Total	С	Ν	Ο	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		Ag	220	1796					0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	Δį	226	Total		Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		AI	220	1796	1125	315	342		0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	Alz	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		АК	220	1796	1125	315	342		0	0
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	1	1 m	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		AIII	220	1796	1125	315	342	14	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	Δο	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		AO	220	1796	1125	315	342	14	0	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	Δα	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		Aq	220	1796	1125	315	342	14		
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	Δα	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		AS	220	1796	1125	315	342	14	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	A	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	Au	220	1796	1125	315	342	14		0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	Δ	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		AW	220	1796	1125	315	342	14	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	۸	226	Total	С	Ν	0	S	0	0
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		Ау	220	1796	1125	315	342	14	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	D.	226	Total	С	Ν	0	S	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		Ba	220	1796	1125	315	342	14	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	D-	006	Total	С	Ν	0	S	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		BC	220	1796	1125	315	342		U	U
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	D-	006	Total	С	Ν	0	S	0	0
$ 1 B_{\sigma} 996 0 0 0$		Be	220	1796	1125	315	342	14	U	0
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	1	D	006	Total	С	Ν	0	S	0	0
		Bg	226	1796	1125	315	342	14	U	U

• Molecule 1 is a protein called Genome polyprotein (Fragment).



Mol	Chain	Residues		At		AltConf	Trace		
1	Bi	226	Total	С	Ν	0	S	0	0
		220	1796	1125	315	342	14	0	0
1	Bk	226	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
		220	1796	1125	315	342	14	0	0
1	Bm	226	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
		220	1796	1125	315	342	14	0	
1	Во	226	Total	С	Ν	Ο	\mathbf{S}	0	0
	D0	220	1796	1125	315	342	14	· · ·	
1	Bq	226	Total	С	Ν	Ο	\mathbf{S}	0	0
	Dq	220	1796	1125	315	342	14	Ŭ	0
1	Bs	226	Total	С	Ν	Ο	\mathbf{S}	0	0
	25		1796	1125	315	342	14	<u> </u>	
1	Bu	226	Total	С	Ν	Ο	\mathbf{S}	0	0
-	Du		1796	1125	315	342	14	<u> </u>	
1	Bw	226	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
-	D	220	1796	1125	315	342	14	0	0
1	By	226	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
-	Dy	220	1796	1125	315	342	14	0	0
1	Ca	226	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
		220	1796	1125	315	342	14	V	
1	Cc	226	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
		220	1796	1125	315	342	14	Ŭ	Ŭ

Continued from previous page...

• Molecule 2 is a RNA chain called RNA (5'-R(P*UP*UP*UP*UP*U)-3').

$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	Mol	Chain	Residues		At	oms			AltConf	Trace
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	9	Ab	Б	Total	С	Ν	0	Р	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	2	AD	5	100	45	10	40	5	0	0
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	2	Ad	5	Total	С	Ν	Ο	Р	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		Au	5	100	45	10	40	5	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	2	Δf	5	Total	С	Ν	Ο	Р	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	2	Π		100	45	10	40	5	0	U
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	9	Ab	К	Total	С	Ν	Ο	Р	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	2	ЛП	5	100	45	10	40	5	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	9	Λi	ц	Total	С	Ν	Ο	Р	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	2	лj	5	100	45	10	40	5	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	9	Δ1	ц	Total	С	Ν	Ο	Р	0	0
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	2	AI	5	100	45	10	40	5	0	0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	0	An	L.	Total	С	Ν	Ο	Р	0	0
			G	100	45	10	40	5	U	U
2 Ap 5 100 45 10 40 5 0 0	0	An	E.	Total	С	Ν	Ο	Р	0	0
		Ар	Ð	100	45	10	40	5	U	U



α \cdot \cdot \cdot	C	•	
Continued	trom	nremons	naae
Continucu	JIONE	precedus	page


Mol	Chain	<i>i previous pa</i> Residues	90	At	oms			AltConf	Trace
			Total	С	Ν	0	Р		
2	Ar	5	100	45	10	40	5	0	0
0	A	~	Total	С	Ν	Ο	Р	0	0
2	At	5	100	45	10	40	5	0	0
2	۸	5	Total	С	Ν	Ο	Р	0	0
	Av	5	100	45	10	40	5	0	0
2	Ax	5	Total	С	Ν	Ο	Р	0	0
	ЛЛ	5	100	45	10	40	5	0	0
2	Az	5	Total	С	Ν	Ο	Р	0	0
	112	0	100	45	10	40	5	0	0
2	Bb	5	Total	С	Ν	Ο	Р	0	0
		<u> </u>	100	45	10	40	5	Ŭ	
2	Bd	5	Total	С	Ν	0	Р	0	0
	24	Ŭ	100	45	10	40	5	Ŭ	
2	Bf	5	Total	С	Ν	0	Р	0	0
		-	100	45	10	40	5		
2	Bh	5	Total	С	Ν	0	Р	0	0
		_	100	45	10	40	5	-	
2	Bj	5	Total	С	Ν	0	Р	0	0
	5		100	45	10	40	5		
2	Bl	5	Total	C	N	0	P	0	0
			100	$\frac{45}{C}$	10 N	40	5		
2	Bn	5	Total	C	N 10	0	P	0	0
			100 Tatal	$\frac{45}{C}$	10 N	40 0	$\frac{5}{P}$		
2	Bp	5	Total				Р 5	0	0
			100 Total	$\frac{45}{C}$	10 N	40 0	$\frac{0}{P}$		
2	Br	5	10tai 100	$\frac{0}{45}$	10	40	г 5	0	0
			Total	40 C	N	$\frac{40}{0}$	P		
2	Bt	5	100a1 100	45	10	40	5	0	0
			Total	-40 C	N	0	P		
2	Bv	5	100	45	10	40	5	0	0
			Total	C	N	$\frac{10}{0}$	P		
2	Bx	5	100	45	10	40	5	0	0
			Total	C	N	0	<u>Р</u>		
2	Bz	5	100	45	10	40	5	0	0
	~		Total	C	N	$\overline{0}$	P		
2	Cb	5	100	45	10	40	5	0	0
	~ .		Total	C	N	0	P		
2	Cd	5		45	10	40		0	0
2	Cđ	Э	100	45	10	40	5	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Genome polyprotein (Fragment)













GLY ASP ASP ASP ASP ASP ASP ASP ASP ASP ASP	N266 M267
• Molecule 1: Genome polyprotein (Fragment)	
Chain By: 84% 15%	
GLY ASN THR ALF ALF ALF ALF ALA CLY CLY SER ALA ALA ALA ALA ALA ALA ALA ALA ALA AL	M267
• Molecule 1: Genome polyprotein (Fragment)	
Chain Ca: 84% 15%	
GLY ASP ASP ASP ASP ASP ASP ASP ASP GLY SSR ASP ASP ASP ASP GLY GLY GLY GLY GLY ASN LYS GLY ASN CLV GLY GLY ASN CLV GLY GLY ASS CLV GLY CLY GLY CLY GLY CLY CCU CLY GLY CLY CCU CCU CCU CCU CCU CCU CCU CCU CCU CC	M267
• Molecule 1: Genome polyprotein (Fragment)	
Chain Cc: 84% 15%	
GLY ASP ASP ASP ASP ASP ASP ASP CIL CVS CLYS CLYS CLVS ASP CLVS GLU CLVS GLU CLVS GLU CLVS GLU CLVS GLU CLVS GLU CLVS GLU CLVS CLV ASN ASN CLVS CLV CLVS CLV CLVS CLV CLVS CLV CLVS CLV CLVS CLV CLVS CLV CLVS CLV CLVS CLV CLVS CLV CLVS CLV CLV CLVS CLV CLV CLV CLV CLV CLV CLV CLV CLV CLV	M267
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')	
Chain Ab: 100%	
There are no outlier residues recorded for this chain.	
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')	
Chain Ad: 100%	
There are no outlier residues recorded for this chain.	
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')	
Chain Af: 100%	
There are no outlier residues recorded for this chain.	
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')	
Chain Ah: 100%	
There are no outlier residues recorded for this chain.	
• Molecule 2: RNA $(5'-R(P*UP*UP*UP*UP*U)-3')$	

WORLDWIDE PROTEIN DATA BANK

Chain Aj: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Al: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain An: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Ap: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Ar: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain At: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Av: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Ax: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Az: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA $(5'-R(P*UP*UP*UP*UP*U)-3')$



Chain Bb: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Bd: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Bf: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Bh: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Bj: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Bl: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Bn: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Bp: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')
Chain Br: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA (5'-R(P*UP*UP*UP*UP*U)-3')



Chain Bt: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA $(5'-R(P*UP*UP*UP*UP*U)-3')$
Chain Bv: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA $(5'-R(P*UP*UP*UP*UP*U)-3')$
Chain Bx: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA $(5'-R(P*UP*UP*UP*UP*U)-3')$
Chain Bz: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA $(5'-R(P*UP*UP*UP*UP*U)-3')$
Chain Cb: 100%
There are no outlier residues recorded for this chain.
• Molecule 2: RNA $(5'-R(P*UP*UP*UP*UP*U)-3')$
Chain Cd: 100%

There are no outlier residues recorded for this chain.



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=-40.960°, rise= 3.966 Å,	Depositor
	axial sym= $C1$	
Number of segments used	10266	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION; patchCTF correction	
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose $(e^-/\text{\AA}^2)$	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2100	Depositor
Magnification	150000	Depositor
Image detector	FEI FALCON III $(4k \ge 4k)$	Depositor
Maximum map value	4.379	Depositor
Minimum map value	-2.558	Depositor
Average map value	0.022	Depositor
Map value standard deviation	0.275	Depositor
Recommended contour level	0.665	Depositor
Map size (Å)	285.0, 285.0, 285.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^{\circ}$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.95, 0.95, 0.95	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
WIOI	Ullaili	RMSZ	# Z > 5	RMSZ	# Z > 5
1	Aa	0.31	0/1833	0.52	0/2482
1	Ac	0.31	0/1833	0.52	0/2482
1	Ae	0.31	0/1833	0.52	0/2482
1	Ag	0.31	0/1833	0.52	0/2482
1	Ai	0.31	0/1833	0.52	0/2482
1	Ak	0.30	0/1833	0.52	0/2482
1	Am	0.31	0/1833	0.52	0/2482
1	Ao	0.31	0/1833	0.52	0/2482
1	Aq	0.31	0/1833	0.52	0/2482
1	As	0.31	0/1833	0.52	0/2482
1	Au	0.31	0/1833	0.52	0/2482
1	Aw	0.31	0/1833	0.52	0/2482
1	Ay	0.31	0/1833	0.52	0/2482
1	Ba	0.31	0/1833	0.52	0/2482
1	Bc	0.30	0/1833	0.52	0/2482
1	Be	0.31	0/1833	0.52	0/2482
1	Bg	0.31	0/1833	0.52	0/2482
1	Bi	0.31	0/1833	0.52	0/2482
1	Bk	0.31	0/1833	0.52	0/2482
1	Bm	0.31	0/1833	0.52	0/2482
1	Bo	0.31	0/1833	0.52	0/2482
1	Bq	0.31	0/1833	0.52	0/2482
1	Bs	0.31	0/1833	0.52	0/2482
1	Bu	0.31	0/1833	0.52	0/2482
1	Bw	0.31	0/1833	0.52	0/2482
1	By	0.31	0/1833	0.52	0/2482
1	Ca	0.31	0/1833	0.52	0/2482
1	Cc	0.31	0/1833	0.52	0/2482
2	Ab	0.28	0/109	0.76	0/166
2	Ad	0.28	0/109	0.76	0/166
2	Af	0.28	0/109	0.76	0/166
2	Ah	0.28	0/109	0.76	0/166
2	Aj	0.27	0/109	0.76	0/166
2	Al	0.28	0/109	0.76	0/166



Mol	Chain	Bond	lengths	Bond	angles
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
2	An	0.28	0/109	0.76	0/166
2	Ap	0.28	0/109	0.76	0/166
2	Ar	0.28	0/109	0.76	0/166
2	At	0.28	0/109	0.76	0/166
2	Av	0.28	0/109	0.76	0/166
2	Ax	0.28	0/109	0.76	0/166
2	Az	0.28	0/109	0.76	0/166
2	Bb	0.28	0/109	0.76	0/166
2	Bd	0.28	0/109	0.76	0/166
2	Bf	0.28	0/109	0.76	0/166
2	Bh	0.28	0/109	0.76	0/166
2	Bj	0.28	0/109	0.76	0/166
2	Bl	0.28	0/109	0.76	0/166
2	Bn	0.28	0/109	0.76	0/166
2	Bp	0.27	0/109	0.76	0/166
2	Br	0.28	0/109	0.76	0/166
2	Bt	0.28	0/109	0.76	0/166
2	Bv	0.28	0/109	0.76	0/166
2	Bx	0.28	0/109	0.76	0/166
2	Bz	0.29	0/109	0.76	0/166
2	Cb	0.28	0/109	0.76	0/166
2	Cd	0.27	0/109	0.76	0/166
All	All	0.30	0/54376	0.54	0/74144

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Aa	1796	0	1775	0	0
1	Ac	1796	0	1775	0	0
1	Ae	1796	0	1775	0	0



	Continued from previous page								
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes			
1	Ag	1796	0	1775	0	0			
1	Ai	1796	0	1775	0	0			
1	Ak	1796	0	1775	0	0			
1	Am	1796	0	1775	0	0			
1	Ao	1796	0	1775	0	0			
1	Aq	1796	0	1775	0	0			
1	As	1796	0	1775	0	0			
1	Au	1796	0	1775	0	0			
1	Aw	1796	0	1775	0	0			
1	Ay	1796	0	1775	0	0			
1	Ba	1796	0	1775	0	0			
1	Bc	1796	0	1775	0	0			
1	Be	1796	0	1775	0	0			
1	Bg	1796	0	1775	0	0			
1	Bi	1796	0	1775	0	0			
1	Bk	1796	0	1775	0	0			
1	Bm	1796	0	1775	0	0			
1	Bo	1796	0	1775	0	0			
1	Bq	1796	0	1775	0	0			
1	Bs	1796	0	1775	0	0			
1	Bu	1796	0	1775	0	0			
1	Bw	1796	0	1775	0	0			
1	By	1796	0	1775	0	0			
1	Ca	1796	0	1775	0	0			
1	Cc	1796	0	1775	0	0			
2	Ab	100	0	51	0	0			
2	Ad	100	0	51	0	0			
2	Af	100	0	51	0	0			
2	Ah	100	0	51	0	0			
2	Aj	100	0	51	0	0			
2	Al	100	0	51	0	0			
2	An	100	0	51	0	0			
2	Ap	100	0	51	0	0			
2	Ar	100	0	51	0	0			
2	At	100	0	51	0	0			
2	Av	100	0	51	0	0			
2	Ax	100	0	51	0	0			
2	Az	100	0	51	0	0			
2	Bb	100	0	51	0	0			
2	Bd	100	0	51	0	0			
2	Bf	100	0	51	0	0			
2	Bh	100	0	51	0 Continu	0			



Mol	Chain	Non-H		H(added)	Clashes	Symm-Clashes
2	Bj	100	0	51	0	0
2	Bl	100	0	51	0	0
2	Bn	100	0	51	0	0
2	Bp	100	0	51	0	0
2	Br	100	0	51	0	0
2	Bt	100	0	51	0	0
2	Bv	100	0	51	0	0
2	Bx	100	0	51	0	0
2	Bz	100	0	51	0	0
2	Cb	100	0	51	0	0
2	Cd	100	0	51	0	0
All	All	53088	0	51128	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	Aa	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Ac	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Ae	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Ag	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Ai	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Ak	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Am	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Ao	224/267~(84%)	212 (95%)	12 (5%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	Aq	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	As	224/267~(84%)	212 (95%)	12~(5%)	0	100	100
1	Au	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Aw	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Ay	224/267~(84%)	212 (95%)	12~(5%)	0	100	100
1	Ba	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Bc	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Be	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Bg	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Bi	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Bk	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Bm	224/267~(84%)	212 (95%)	12~(5%)	0	100	100
1	Bo	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Bq	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Bs	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Bu	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Bw	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	By	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Ca	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
1	Cc	224/267~(84%)	212 (95%)	12 (5%)	0	100	100
All	All	6272/7476~(84%)	5936 (95%)	336 (5%)	0	100	100

Continued from previous page...

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric Outliers		Percentiles	
1	Aa	196/229~(86%)	195 (100%)	1 (0%)	88 96



α \cdot \cdot \cdot	C		
Continued	trom	previous	page
	J	1	1

Mol	Chain	Analysed	Rotameric	Outliers	Perc	centiles
1	Ac	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Ae	196/229~(86%)	195~(100%)	1 (0%)	88	96
1	Ag	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Ai	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Ak	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Am	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Ao	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Aq	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	As	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Au	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Aw	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Ay	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Ba	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bc	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Be	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bg	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bi	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bk	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bm	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bo	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bq	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bs	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bu	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Bw	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	By	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Ca	196/229~(86%)	195 (100%)	1 (0%)	88	96
1	Cc	196/229~(86%)	195 (100%)	1 (0%)	88	96
All	All	5488/6412 (86%)	5460 (100%)	28 (0%)	89	96

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Aa	267	MET
	a	7	

Mol	Chain	Res	Type
1	Ac	267	MET
1	Ae	267	MET
1	Ag	267	MET
1	Ai	267	MET
1	Ak	267	MET
1	Am	267	MET
1	Ao	267	MET
1	Aq	267	MET
1	As	267	MET
1	Au	267	MET
1	Aw	267	MET
1	Ay	267	MET
1	Ba	267	MET
1	Bc	267	MET
1	Be	267	MET
1	Bg	267	MET
1	Bi	267	MET
1	Bk	267	MET
1	Bm	267	MET
1	Bo	267	MET
1	Bq	267	MET
1	Bs	267	MET
1	Bu	267	MET
1	Bw	267	MET
1	By	267	MET
1	Ca	267	MET
1	Cc	267	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (73) such sidechains are listed below:

Mol	Chain	Res	Type
1	Aa	69	HIS
1	Aa	98	GLN
1	Aa	245	ASN
1	Ac	69	HIS
1	Ac	98	GLN
1	Ac	245	ASN
1	Ae	69	HIS
1	Ae	98	GLN
1	Ae	245	ASN
1	Ag	69	HIS
1	Ag	98	GLN



Mol	Chain	Res	Type
1	Ag	245	ASN
1	Ai	69	HIS
1	Ai	98	GLN
1	Ai	245	ASN
1	Ak	69	HIS
1	Ak	98	GLN
1	Ak	245	ASN
1	Am	69	HIS
1	Am	98	GLN
1	Am	245	ASN
1	Ao	69	HIS
1	Ao	98	GLN
1	Ao	245	ASN
1	Aq	69	HIS
1	Aq	98	GLN
1	As	69	HIS
1	As	98	GLN
1	As	245	ASN
1	Au	69	HIS
1	Au	245	ASN
1	Aw	69	HIS
1	Aw	98	GLN
1	Aw	245	ASN
1	Ay	69	HIS
1	Ay	98	GLN
1	Ay	245	ASN
1	Ba	69	HIS
1	Ba	98	GLN
1	Ba	245	ASN
1	Bc	69	HIS
1	Bc	98	GLN
1	Bc	245	ASN
1	Be	69	HIS
1	Be	98	GLN
1	Be	245	ASN
1	Bg	69	HIS
1	Bg	98	GLN
1	Bi	69	HIS
1	Bi	98	GLN
1	Bk	69	HIS
1	Bk	98	GLN
1	Bm	69	HIS



Mol	Chain	Res	Type
1	Bm	98	GLN
1	Bo	69	HIS
1	Bo	98	GLN
1	Bq	69	HIS
1	Bq	98	GLN
1	Bs	69	HIS
1	Bs	98	GLN
1	Bu	69	HIS
1	Bu	98	GLN
1	Bw	69	HIS
1	Bw	98	GLN
1	Bw	245	ASN
1	By	69	HIS
1	By	98	GLN
1	By	245	ASN
1	Ca	69	HIS
1	Ca	98	GLN
1	Ca	245	ASN
1	Cc	98	GLN
1	Cc	245	ASN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	Ab	4/5~(80%)	0	0
2	Ad	4/5~(80%)	0	0
2	Af	4/5~(80%)	0	0
2	Ah	4/5~(80%)	0	0
2	Aj	4/5~(80%)	0	0
2	Al	4/5~(80%)	0	0
2	An	4/5~(80%)	0	0
2	Ар	4/5~(80%)	0	0
2	Ar	4/5~(80%)	0	0
2	At	4/5~(80%)	0	0
2	Av	4/5~(80%)	0	0
2	Ax	4/5~(80%)	0	0
2	Az	4/5~(80%)	0	0
2	Bb	4/5~(80%)	0	0
2	Bd	4/5~(80%)	0	0
2	Bf	4/5~(80%)	0	0
2	Bh	4/5~(80%)	0	0



	Continuea from previous page				
Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	
2	Bj	4/5~(80%)	0	0	
2	Bl	4/5~(80%)	0	0	
2	Bn	4/5~(80%)	0	0	
2	Bp	4/5~(80%)	0	0	
2	Br	4/5~(80%)	0	0	
2	Bt	4/5~(80%)	0	0	
2	Bv	4/5~(80%)	0	0	
2	Bx	4/5~(80%)	0	0	
2	Bz	4/5~(80%)	0	0	
2	Cb	4/5~(80%)	0	0	
2	Cd	4/5~(80%)	0	0	
All	All	112/140~(80%)	0	0	

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-17048. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)

6.2.1 Primary map



6.2.2 Raw map



X Index: 150

Y Index: 150




The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map



6.3.2 Raw map



X Index: 118

Y Index: 182




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.665. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




Mask visualisation (i) 6.6

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

$emd_{17048}msk_{1.map}$ (i) 6.6.1






7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 560 $\rm nm^3;$ this corresponds to an approximate mass of 506 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.334 \AA^{-1}



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.334 \AA^{-1}



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
Resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	2.99	-	-
Author-provided FSC curve	2.99	3.37	3.01
Unmasked-calculated*	3.63	4.33	3.70


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.63 differs from the reported value 2.99 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-17048 and PDB model 80PC. Per-residue inclusion information can be found in section 3 on page 8.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.665 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.665).



9.4 Atom inclusion (i)



At the recommended contour level, 85% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.665) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7990	0.5790
Aa	0.7890	0.5760
Ab	0.9600	0.5890
Ac	0.7850	0.5770
Ad	0.9500	0.5960
Ae	0.7870	0.5780
Af	0.9600	0.5870
Ag	0.7930	0.5780
Ah	0.9600	0.5980
Ai	0.7900	0.5790
Aj	0.9500	0.5870
Ak	0.7880	0.5790
Al	0.9500	0.5960
Am	0.7890	0.5780
An	0.9500	0.5980
Ao	0.7900	0.5780
Ар	0.9600	0.5970
Aq	0.7960	0.5790
Ar	0.9400	0.5970
As	0.7920	0.5770
At	0.9500	0.5930
Au	0.7940	0.5780
Av	0.9400	0.5920
Aw	0.7870	0.5770
Ax	0.9500	0.5900
Ay	0.7900	0.5780
Az	0.9500	0.5950
Ba	0.7910	0.5790
Bb	0.9600	0.5920
Bc	0.7920	0.5810
Bd	0.9500	0.6000
Be	0.7900	0.5760
Bf	0.9500	0.5970
Bg	0.7960	0.5760
Bh	0.9500	0.5870



Chain	Atom inclusion	Q-score
Bi	0.7870	0.5780
Bj	0.9400	0.5910
Bk	0.7910	0.5790
Bl	0.9400	0.5940
Bm	0.7880	0.5780
Bn	0.9500	0.5960
Bo	0.7900	0.5790
Bp	0.9500	0.5960
Bq	0.7890	0.5780
Br	0.9500	0.5960
Bs	0.7880	0.5760
Bt	0.9600	0.5930
Bu	0.7890	0.5750
Bv	0.9300	0.5940
Bw	0.7900	0.5770
Bx	0.9600	0.5960
By	0.7910	0.5780
Bz	0.9400	0.5930
Ca	0.7890	0.5770
Cb	0.9500	0.5950
Cc	0.7890	0.5780
Cd	0.9400	0.5940

