



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 07:58 am BST

PDB ID : 5OQ1  
Title : Crystal structure of *Serratia marcescens* ChiX (used as MR model for superior PDB 5OPZ)  
Authors : Owen, R.A.; Fyfe, P.K.; Lodge, A.; Biboy, J.; Vollmer, W.; Hunter, W.N.; Sargent, F.  
Deposited on : 2017-08-10  
Resolution : 1.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11



## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2800 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ChiX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	133	1209	755	241	207	6	0	14	0
1	B	133	1189	744	233	205	7	11	12	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP A0A0U7UVX7
A	-2	PRO	-	expression tag	UNP A0A0U7UVX7
A	-1	LEU	-	expression tag	UNP A0A0U7UVX7
A	0	GLY	-	expression tag	UNP A0A0U7UVX7
A	1	SER	-	expression tag	UNP A0A0U7UVX7
B	-3	GLY	-	expression tag	UNP A0A0U7UVX7
B	-2	PRO	-	expression tag	UNP A0A0U7UVX7
B	-1	LEU	-	expression tag	UNP A0A0U7UVX7
B	0	GLY	-	expression tag	UNP A0A0U7UVX7
B	1	SER	-	expression tag	UNP A0A0U7UVX7

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	233	Total 233	O 233	0	0
4	B	166	Total 166	O 166	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ChiX



- Molecule 1: ChiX



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.56Å 55.50Å 51.60Å 90.00° 90.16° 90.00°	Depositor
Resolution (Å)	51.60 – 1.34 51.60 – 1.34	Depositor EDS
% Data completeness (in resolution range)	95.8 (51.60-1.34) 98.6 (51.60-1.34)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.57 (at 1.34Å)	Xtriage
Refinement program	REFMAC 5.8.0124	Depositor
R, $R_{free}$	0.132 , 0.163 0.134 , 0.166	Depositor DCC
$R_{free}$ test set	2561 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.7	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 34.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.096 for h,-k,-l	Xtriage
Reported twinning fraction	0.840 for H, K, L 0.160 for h,-k,-l	Depositor
Outliers	0 of 51229 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2800	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.66	0/1234	0.92	4/1665 (0.2%)
1	B	0.65	0/1213	0.94	5/1637 (0.3%)
All	All	0.66	0/2447	0.93	9/3302 (0.3%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	99	ARG	NE-CZ-NH1	7.12	123.86	120.30
1	A	99	ARG	NE-CZ-NH2	-6.03	117.28	120.30
1	B	19[A]	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	B	19[B]	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	B	89	ASP	CB-CG-OD1	5.48	123.23	118.30
1	B	30	ARG	NE-CZ-NH1	5.43	123.02	120.30
1	A	65[A]	ARG	NE-CZ-NH2	5.36	122.98	120.30
1	A	65[B]	ARG	NE-CZ-NH2	5.36	122.98	120.30
1	B	53	ARG	NE-CZ-NH2	-5.31	117.64	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1209	0	1213	6	0
1	B	1189	0	1194	6	2
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
4	A	233	0	0	2	3
4	B	166	0	0	4	1
All	All	2800	0	2407	12	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:99[A]:ARG:NH1	4:B:301:HOH:O	1.99	0.96
1:B:100[B]:ARG:HH11	1:B:100[B]:ARG:HG3	1.60	0.67
1:A:63:THR:HG22	4:A:403:HOH:O	1.97	0.63
1:A:57[B]:ARG:CZ	4:A:308:HOH:O	2.56	0.54
1:B:100[B]:ARG:CG	1:B:100[B]:ARG:HH11	2.21	0.52
1:B:17:ASN:HB2	4:B:302:HOH:O	2.15	0.46
1:A:117[B]:THR:HG22	4:B:363:HOH:O	2.16	0.45
1:A:23[B]:PRO:HB2	1:A:27[B]:ARG:HH12	1.83	0.44
1:B:37[B]:LEU:HD12	1:B:37[B]:LEU:N	2.33	0.43
1:A:47:ARG:O	1:A:69:LEU:HD23	2.19	0.42
1:A:63:THR:HG21	4:B:410:HOH:O	2.21	0.41
1:B:100[B]:ARG:CG	1:B:100[B]:ARG:NH1	2.84	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:492:HOH:O	4:B:457:HOH:O[1_556]	1.82	0.38
1:B:27[B]:ARG:NH2	4:A:441:HOH:O[1_655]	1.91	0.29
1:B:13:ARG:NH2	4:A:339:HOH:O[2_746]	2.07	0.13



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	146/137 (107%)	145 (99%)	1 (1%)	0	100	100
1	B	144/137 (105%)	143 (99%)	1 (1%)	0	100	100
All	All	290/274 (106%)	288 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/113 (112%)	124 (98%)	2 (2%)	62	29
1	B	124/113 (110%)	121 (98%)	3 (2%)	49	14
All	All	250/226 (111%)	245 (98%)	5 (2%)	67	20

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65[A]	ARG
1	A	65[B]	ARG
1	B	1	SER
1	B	99[A]	ARG
1	B	99[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	133/137 (97%)	0.28	5 (3%) 40 46	6, 10, 19, 36	0
1	B	133/137 (97%)	0.29	4 (3%) 50 57	6, 9, 20, 40	0
All	All	266/274 (97%)	0.29	9 (3%) 45 51	6, 10, 20, 40	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	SER	6.0
1	A	1	SER	4.9
1	A	133	ALA	3.9
1	A	3	ASN	3.9
1	B	2	LEU	3.4
1	B	3	ASN	2.7
1	A	65[A]	ARG	2.5
1	A	4	ASP	2.5
1	B	4	ASP	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	A	201	1/1	1.00	0.14	10,10,10,10	0
3	CL	A	202	1/1	1.00	0.04	19,19,19,19	0
2	ZN	B	201	1/1	1.00	0.11	9,9,9,9	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.