



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 1, 2023 – 02:07 PM EDT

PDB ID : 4OTT  
Title : Crystal structure of the gamma-glutamyltranspeptidase from *Bacillus licheniformis*.  
Authors : Merlino, A.  
Deposited on : 2014-02-14  
Resolution : 2.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2754 (3.00-2.96)
Clashscore	141614	3103 (3.00-2.96)
Ramachandran outliers	138981	2993 (3.00-2.96)
Sidechain outliers	138945	2996 (3.00-2.96)
RSRZ outliers	127900	2644 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	 60% 25% 10%
2	B	187	 70% 24%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4159 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma glutamyl transpeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	2744	1740	453	537	14	0	0	0

- Molecule 2 is a protein called Gamma-glutamyltranspeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	182	1409	892	233	280	4	0	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	O	0	0
			2	2		
4	B	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gamma glutamyl transpeptidase



- Molecule 2: Gamma-glutamyltranspeptidase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.90Å 61.97Å 148.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.25 – 2.98 26.25 – 2.99	Depositor EDS
% Data completeness (in resolution range)	97.2 (26.25-2.98) 97.2 (26.25-2.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.60 (at 3.00Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.191 , 0.250 0.191 , 0.250	Depositor DCC
$R_{free}$ test set	548 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.5	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , -6.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.109 for k,h,-l	Xtriage
Reported twinning fraction	0.858 for H, K, L 0.142 for K, H, -L	Depositor
Outliers	0 of 11634 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4159	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	1/2805 (0.0%)	0.82	2/3787 (0.1%)
2	B	0.63	0/1440	0.81	0/1959
All	All	0.61	1/4245 (0.0%)	0.81	2/5746 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	221	MET	CG-SD	-10.99	1.52	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	53	SER	N-CA-C	6.20	127.75	111.00
1	A	330	ASP	CB-CG-OD1	5.25	123.02	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	53	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2744	0	2684	112	0
2	B	1409	0	1363	36	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
All	All	4159	0	4047	132	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (132) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:161:LYS:CB	1:A:162:MET:HA	1.95	0.96
1:A:53:SER:N	1:A:54:LYS:CG	2.32	0.92
1:A:161:LYS:CG	1:A:162:MET:HA	1.99	0.91
1:A:161:LYS:HD3	1:A:162:MET:HG2	1.56	0.88
1:A:98:GLY:HA3	1:A:99:GLU:CB	2.05	0.86
1:A:161:LYS:HB3	1:A:162:MET:HA	1.56	0.86
1:A:161:LYS:HZ2	1:A:163:GLU:C	1.80	0.84
2:B:427:LEU:HD21	2:B:432:GLY:HA2	1.63	0.81
1:A:51:LEU:O	1:A:54:LYS:CG	2.31	0.79
2:B:405:THR:HG22	2:B:554:ILE:HD13	1.68	0.76
1:A:161:LYS:HG3	1:A:165:LEU:H	1.52	0.73
1:A:161:LYS:HG2	1:A:163:GLU:N	2.03	0.73
1:A:52:ALA:C	1:A:54:LYS:CG	2.57	0.72
1:A:98:GLY:HA3	1:A:99:GLU:HB3	1.71	0.71
1:A:51:LEU:C	1:A:54:LYS:CG	2.60	0.70
1:A:161:LYS:NZ	1:A:163:GLU:C	2.46	0.68
1:A:227:LEU:HD12	1:A:227:LEU:O	1.92	0.68
1:A:161:LYS:HD3	1:A:162:MET:CG	2.22	0.68
1:A:154:HIS:O	1:A:155:LYS:CB	2.42	0.68
1:A:333:ALA:HB2	1:A:372:GLU:HG2	1.75	0.67
1:A:161:LYS:HD2	1:A:165:LEU:CB	2.25	0.67

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:GLY:HA3	1:A:99:GLU:HB2	1.78	0.66
1:A:161:LYS:NZ	1:A:166:ILE:HG22	2.11	0.65
1:A:161:LYS:HZ3	1:A:166:ILE:H	1.42	0.65
1:A:161:LYS:CD	1:A:162:MET:HG2	2.24	0.65
2:B:399:THR:HB	2:B:480:PRO:HB3	1.78	0.63
1:A:161:LYS:CG	1:A:165:LEU:H	2.10	0.63
2:B:473:PRO:O	2:B:558:ARG:NH2	2.33	0.62
1:A:364:ASP:O	1:A:365:SER:HB3	2.00	0.62
1:A:161:LYS:HG2	1:A:162:MET:HA	1.83	0.61
1:A:161:LYS:CD	1:A:165:LEU:H	2.13	0.61
1:A:148:LYS:HB3	1:A:266:VAL:HG13	1.82	0.61
1:A:161:LYS:HD2	1:A:165:LEU:HB2	1.82	0.60
1:A:265:GLU:N	1:A:265:GLU:OE1	2.35	0.60
2:B:403:THR:OG1	2:B:476:THR:HG21	2.02	0.59
1:A:220:ASP:OD2	2:B:430:GLY:HA2	2.03	0.59
1:A:78:ASN:ND2	1:A:87:ILE:HG23	2.18	0.58
1:A:333:ALA:HB1	1:A:387:ILE:HG12	1.85	0.58
1:A:80:THR:HA	1:A:177:PHE:CZ	2.39	0.58
1:A:161:LYS:CE	1:A:162:MET:HG2	2.34	0.57
1:A:161:LYS:CD	1:A:162:MET:HA	2.35	0.57
1:A:137:HIS:HD2	1:A:139:ASN:H	1.52	0.57
1:A:286:PRO:HB2	1:A:338:PRO:HG3	1.86	0.57
1:A:87:ILE:HA	1:A:146:THR:HB	1.87	0.57
1:A:161:LYS:HZ3	1:A:166:ILE:HG22	1.69	0.56
1:A:161:LYS:HE3	1:A:162:MET:HG2	1.86	0.56
1:A:74:GLN:HE22	1:A:85:SER:HB2	1.69	0.56
2:B:511:ARG:NH1	2:B:548:ILE:O	2.41	0.54
1:A:161:LYS:HD2	1:A:165:LEU:H	1.73	0.54
1:A:318:HIS:O	1:A:322:GLU:HG2	2.07	0.54
2:B:550:ASN:HA	2:B:568:ASP:OD2	2.08	0.53
1:A:154:HIS:O	1:A:155:LYS:HB2	2.08	0.53
1:A:151:GLU:O	1:A:154:HIS:O	2.26	0.52
1:A:161:LYS:HB3	1:A:162:MET:CA	2.35	0.52
1:A:54:LYS:CG	1:A:55:ILE:H	2.23	0.52
1:A:161:LYS:HD2	1:A:165:LEU:CG	2.40	0.52
1:A:161:LYS:CE	1:A:166:ILE:HG22	2.41	0.51
1:A:364:ASP:O	1:A:365:SER:CB	2.59	0.51
1:A:284:MET:SD	1:A:285:PRO:HD2	2.52	0.50
1:A:358:GLN:HA	1:A:361:ILE:HD12	1.92	0.50
2:B:550:ASN:ND2	2:B:571:ARG:HD2	2.27	0.50
1:A:284:MET:HE3	1:A:285:PRO:O	2.11	0.50

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:GLN:HB3	1:A:221:MET:HB3	1.94	0.49
1:A:55:ILE:HD13	1:A:168:PRO:HB2	1.95	0.49
1:A:65:ASN:OD1	1:A:67:ILE:N	2.46	0.49
1:A:161:LYS:HG3	1:A:165:LEU:N	2.23	0.49
2:B:439:LEU:O	2:B:442:PHE:HB2	2.12	0.49
1:A:49:HIS:CG	1:A:50:PRO:HD2	2.48	0.48
1:A:62:LYS:O	1:A:160:LYS:NZ	2.47	0.48
1:A:287:PRO:O	2:B:459:LEU:HD12	2.13	0.48
1:A:210:LEU:H	1:A:210:LEU:HD12	1.79	0.48
1:A:118:PRO:HG3	2:B:453:GLN:NE2	2.28	0.47
1:A:163:GLU:O	1:A:167:SER:OG	2.31	0.47
2:B:441:ASP:HB2	2:B:458:PRO:HB2	1.95	0.47
1:A:49:HIS:CD2	1:A:51:LEU:HB2	2.49	0.47
1:A:333:ALA:CB	1:A:372:GLU:HG2	2.41	0.47
1:A:236:PHE:HA	1:A:241:ILE:HB	1.96	0.47
1:A:44:MET:HB3	2:B:405:THR:CG2	2.45	0.47
1:A:203:PHE:O	1:A:210:LEU:HD12	2.14	0.47
1:A:307:SER:OG	2:B:498:GLU:OE1	2.33	0.47
1:A:205:PRO:HD2	1:A:208:GLU:O	2.15	0.47
2:B:420:GLN:HB2	2:B:437:ASN:HD21	1.81	0.46
1:A:370:VAL:HG12	1:A:370:VAL:O	2.16	0.45
1:A:188:LYS:O	1:A:191:GLN:HG3	2.15	0.45
2:B:402:PHE:CZ	2:B:414:TYR:CD2	3.05	0.45
1:A:161:LYS:HG2	1:A:162:MET:CA	2.47	0.45
1:A:245:ILE:HG23	2:B:435:LEU:HD11	1.98	0.45
1:A:49:HIS:CD2	1:A:51:LEU:H	2.33	0.45
1:A:54:LYS:CG	1:A:55:ILE:N	2.80	0.44
1:A:236:PHE:CD1	1:A:237:TYR:CE1	3.05	0.44
1:A:83:MET:HB3	2:B:422:PHE:CE1	2.52	0.44
1:A:227:LEU:CD2	1:A:240:GLU:OE2	2.65	0.44
1:A:337:ASP:OD1	1:A:339:GLU:HB2	2.18	0.44
2:B:403:THR:OG1	2:B:554:ILE:HD12	2.17	0.44
1:A:161:LYS:HE2	1:A:166:ILE:HG22	2.01	0.43
1:A:365:SER:HA	2:B:537:PHE:CZ	2.53	0.43
1:A:297:MET:HE2	2:B:490:PHE:CG	2.53	0.43
1:A:250:GLN:HG3	1:A:254:GLY:O	2.19	0.43
2:B:551:VAL:O	2:B:568:ASP:HB2	2.18	0.43
1:A:115:GLY:HA3	1:A:255:SER:OG	2.19	0.43
1:A:287:PRO:HB3	1:A:337:ASP:HA	2.01	0.43
1:A:297:MET:HE2	2:B:490:PHE:CB	2.49	0.43
2:B:507:ILE:HG12	2:B:551:VAL:CG1	2.49	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:ILE:HD13	1:A:284:MET:SD	2.59	0.43
2:B:552:GLN:HG3	2:B:567:ALA:HA	2.00	0.43
2:B:399:THR:HB	2:B:480:PRO:CB	2.48	0.42
1:A:93:MET:HB3	1:A:106:ILE:HB	2.00	0.42
1:A:154:HIS:O	1:A:155:LYS:HB3	2.18	0.42
2:B:515:ASN:OD1	2:B:519:SER:HB3	2.20	0.42
1:A:161:LYS:NZ	1:A:163:GLU:O	2.52	0.42
1:A:257:THR:OG1	1:A:259:ASP:HB3	2.20	0.42
1:A:286:PRO:HA	1:A:288:SER:N	2.34	0.42
1:A:67:ILE:HD13	1:A:157:TRP:CB	2.50	0.42
1:A:182:VAL:HG21	2:B:571:ARG:HB3	2.01	0.42
2:B:544:ASN:HB2	2:B:545:PRO:CD	2.50	0.42
1:A:118:PRO:HA	2:B:452:VAL:HB	2.00	0.42
1:A:227:LEU:HD21	1:A:240:GLU:OE2	2.19	0.41
1:A:257:THR:HB	1:A:258:PRO:HD2	2.02	0.41
1:A:61:LYS:HG2	2:B:578:VAL:HG13	2.01	0.41
1:A:149:GLY:HA2	1:A:266:VAL:HG11	2.03	0.41
1:A:148:LYS:CD	1:A:265:GLU:HA	2.50	0.41
1:A:161:LYS:HG2	1:A:162:MET:C	2.40	0.41
1:A:166:ILE:HD11	1:A:225:PHE:CG	2.55	0.41
1:A:161:LYS:HG2	1:A:164:ASP:N	2.36	0.41
1:A:296:GLN:NE2	1:A:348:LEU:O	2.51	0.41
1:A:61:LYS:CG	2:B:578:VAL:HG13	2.51	0.40
1:A:135:SER:O	2:B:424:THR:HA	2.20	0.40
1:A:76:ALA:O	1:A:80:THR:OG1	2.38	0.40
1:A:161:LYS:HD2	1:A:165:LEU:HG	2.03	0.40
1:A:284:MET:O	1:A:291:GLY:HA3	2.22	0.40
1:A:140:ALA:HB1	2:B:437:ASN:HB2	2.03	0.40
2:B:475:LEU:HG	2:B:476:THR:N	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/398 (89%)	329 (93%)	19 (5%)	7 (2%)	7	32
2	B	180/187 (96%)	167 (93%)	11 (6%)	2 (1%)	14	47
All	All	535/585 (92%)	496 (93%)	30 (6%)	9 (2%)	9	36

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	99	GLU
1	A	155	LYS
1	A	365	SER
1	A	98	GLY
2	B	419	GLU
1	A	161	LYS
1	A	54	LYS
1	A	130	PRO
2	B	480	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/325 (89%)	256 (89%)	32 (11%)	6	23
2	B	158/162 (98%)	143 (90%)	15 (10%)	8	30
All	All	446/487 (92%)	399 (90%)	47 (10%)	7	25

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	THR
1	A	83	MET
1	A	87	ILE
1	A	119	ASP
1	A	124	GLU
1	A	125	ASP
1	A	132	SER

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	156	LYS
1	A	160	LYS
1	A	161	LYS
1	A	164	ASP
1	A	172	LEU
1	A	200	LYS
1	A	201	ASP
1	A	218	GLN
1	A	219	LYS
1	A	223	LYS
1	A	227	LEU
1	A	229	ARG
1	A	231	GLU
1	A	233	SER
1	A	240	GLU
1	A	248	VAL
1	A	255	SER
1	A	261	LEU
1	A	300	LEU
1	A	301	ILE
1	A	307	SER
1	A	345	LEU
1	A	372	GLU
1	A	377	LYS
1	A	385	TYR
2	B	399	THR
2	B	403	THR
2	B	405	THR
2	B	414	TYR
2	B	419	GLU
2	B	437	ASN
2	B	446	PRO
2	B	471	GLU
2	B	472	LYS
2	B	514	THR
2	B	543	SER
2	B	558	ARG
2	B	564	MET
2	B	574	THR
2	B	580	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	HIS
1	A	78	ASN
1	A	137	HIS
1	A	390	GLN
2	B	437	ASN
2	B	550	ASN
2	B	552	GLN
2	B	572	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	357/398 (89%)	-0.30	1 (0%) 94 87	20, 42, 59, 81	0
2	B	182/187 (97%)	-0.48	0 100 100	16, 31, 51, 74	0
All	All	539/585 (92%)	-0.36	1 (0%) 95 89	16, 37, 59, 81	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	162	MET	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	A	401	1/1	0.94	0.17	20,20,20,20	0
3	MG	B	601	1/1	0.97	0.21	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.