

Full wwPDB X-ray Structure Validation Report (i)

May 22, 2020 – 04:23 am BST

PDB ID : 10UT

Title : TROUT HEMOGLOBIN I

Authors : Tame, J.; Wilson, J.

Deposited on : 1996-06-21

Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp

with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain				
1	A	143	71%	27%			
2	В	146	77%	20%	•••		



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2475 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called HEMOGLOBIN I.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	143	Total	С	N	О	S	0	9	0
1	A	140	1080	701	184	191	4	0	2	U


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASP	DELETION	UNP P02019
A	?	-	LYS	DELETION	UNP P02019

• Molecule 2 is a protein called HEMOGLOBIN I.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	В	146	Total 1136	C 743	N 188	O 200	S 5	0	2	0

• Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).





Mol	Chain	Residues	${f Atoms}$				ZeroOcc	AltConf	
3	A	1	Total 43					0	0
3	В	1	Total 43		Fe 1		O 4	0	0

• Molecule 4 is water.

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
4	A	81	Total O 81 81	0	0
4	В	92	Total O 92 92	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: HEMOGLOBIN I





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants	63.28Å 63.28Å 312.74Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 - 2.30	Depositor
% Data completeness	(Not available) (8.00-2.30)	Depositor
(in resolution range)	(110t available) (0.00 2.00)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.162 , 0.247	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2475	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

ſ	Mol	Chain	Bond	lengths	Bond angles		
	MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
	1	A	0.76	0/1116	1.44	8/1510~(0.5%)	
	2	В	0.69	0/1176	1.30	4/1596~(0.3%)	
	All	All	0.72	0/2292	1.37	$12/3106 \ (0.4\%)$	

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	A	142	ARG	NE-CZ-NH2	-10.08	115.26	120.30
2	В	104	ARG	NE-CZ-NH2	9.78	125.19	120.30
1	A	142	ARG	NE-CZ-NH1	8.92	124.76	120.30
1	A	93	ARG	NE-CZ-NH2	-8.39	116.11	120.30
1	A	93	ARG	NE-CZ-NH1	7.75	124.18	120.30
1	A	141	TYR	CB-CG-CD1	-7.23	116.66	121.00
1	A	93	ARG	CD-NE-CZ	7.05	133.48	123.60
1	A	36	TYR	CB-CG-CD1	6.78	125.06	121.00
2	В	85	TYR	CB-CG-CD1	6.35	124.81	121.00
1	A	1	SER	N-CA-CB	6.02	119.53	110.50
2	В	144	ARG	NE-CZ-NH2	-5.24	117.68	120.30
2	В	134	MET	CA-CB-CG	5.03	121.85	113.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	1080	0	1101	28	0
2	В	1136	0	1152	26	0
3	A	43	0	30	5	0
3	В	43	0	30	4	0
4	A	81	0	0	3	0
4	В	92	0	0	3	0
All	All	2475	0	2313	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} ({\rm \AA}) \end{array}$	Clash overlap (Å)
2:B:15[A]:TRP:CH2	2:B:72:ASP:OD1	1.84	1.30
1:A:87:LEU:HA	1:A:91:LYS:HD3	1.54	0.90
1:A:91:LYS:HB3	4:A:171:HOH:O	1.70	0.89
2:B:15[A]:TRP:CZ3	2:B:72:ASP:OD1	2.27	0.86
1:A:0:ACE:O	1:A:1:SER:HB2	1.83	0.76
2:B:15[A]:TRP:CH2	2:B:72:ASP:CG	2.66	0.69
2:B:86:LYS:HG2	2:B:144:ARG:HD2	1.76	0.68
1:A:119:THR:HB	1:A:120:PRO:HD2	1.79	0.64
2:B:91:THR:HG21	4:B:176:HOH:O	1.96	0.64
2:B:91:THR:HG22	4:B:239:HOH:O	1.99	0.61
2:B:15[A]:TRP:CZ2	2:B:72:ASP:HB2	2.35	0.61
2:B:15[A]:TRP:HH2	2:B:72:ASP:OD1	1.74	0.61
1:A:97:GLY:HA2	1:A:100:LYS:HE3	1.86	0.58
2:B:144:ARG:CZ	2:B:146[B]:PHE:HZ	2.17	0.58
1:A:119:THR:HB	1:A:120:PRO:CD	2.34	0.57
1:A:62:ILE:HG21	3:A:143:HEM:HBA1	1.85	0.57
1:A:84:LEU:HD12	1:A:137:LEU:HD21	1.87	0.56
3:A:143:HEM:HBA2	3:A:143:HEM:HMA2	1.89	0.55
1:A:84:LEU:HD22	1:A:87:LEU:HD12	1.90	0.54
1:A:37:PRO:HB3	1:A:40:LYS:HZ3	1.72	0.54
1:A:89:ALA:O	1:A:93:ARG:HD3	2.08	0.54
3:A:143:HEM:HBC2	3:A:143:HEM:HMC2	1.91	0.53
1:A:17:ILE:HG13	1:A:114:PHE:CE2	2.43	0.53
2:B:15[A]:TRP:CZ2	2:B:72:ASP:CB	2.91	0.53
1:A:112:ILE:HG13	2:B:115:ALA:HB1	1.90	0.53

Continued on next page...



Continued from previous page...

A tom 1		Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap $(ext{Å})$
2:B:15[A]:TRP:CZ3	2:B:76:LYS:NZ	2.79	0.51
3:A:143:HEM:CMA	3:A:143:HEM:HBA2	2.42	0.50
2:B:45:PHE:HA	2:B:59:LYS:HE2	1.93	0.50
1:A:46:TRP:HE3	1:A:49:LEU:HD23	1.77	0.49
2:B:106:LEU:HD22	3:B:148:HEM:CBB	2.42	0.49
3:B:148:HEM:HBA1	3:B:148:HEM:HHA	1.94	0.48
1:A:48:ASP:HB3	1:A:53:SER:HB2	1.95	0.48
2:B:106:LEU:HD22	3:B:148:HEM:HBB2	1.96	0.48
1:A:0:ACE:H2	4:A:148:HOH:O	2.15	0.47
1:A:27:GLU:OE2	1:A:31:ARG:NE	2.44	0.45
2:B:74:ALA:HB2	2:B:85:TYR:CE2	2.51	0.45
1:A:54:GLY:N	1:A:55:PRO:HD2	2.32	0.45
1:A:73:LEU:HD13	1:A:80:GLY:HA3	1.98	0.45
3:B:148:HEM:HBC2	3:B:148:HEM:HMC2	1.99	0.45
1:A:0:ACE:CH3	4:A:148:HOH:O	2.66	0.44
2:B:15[A]:TRP:HZ2	2:B:72:ASP:HB2	1.79	0.44
1:A:17:ILE:HD11	1:A:114:PHE:CG	2.53	0.44
2:B:105:VAL:O	2:B:109:VAL:HG23	2.18	0.44
3:A:143:HEM:HBC2	3:A:143:HEM:CMC	2.48	0.44
2:B:24:GLY:N	2:B:25:PRO:HD2	2.33	0.44
1:A:114:PHE:N	1:A:115:PRO:CD	2.80	0.43
2:B:4:THR:O	2:B:8:LYS:HG3	2.19	0.43
1:A:20:LYS:O	1:A:24:VAL:HG23	2.18	0.43
1:A:33:LEU:O	1:A:40:LYS:NZ	2.46	0.43
1:A:87:LEU:HA	1:A:91:LYS:CD	2.38	0.43
2:B:18:VAL:HG13	2:B:23:ILE:HD12	2.00	0.43
1:A:17:ILE:HG13	1:A:114:PHE:CZ	2.53	0.42
1:A:37:PRO:HB3	1:A:40:LYS:NZ	2.33	0.42
1:A:84:LEU:HD13	1:A:88:HIS:CE1	2.54	0.42
2:B:15[B]:TRP:CZ2	2:B:20:ILE:HD11	2.54	0.41
2:B:17:LYS:HG3	4:B:185:HOH:O	2.21	0.41
2:B:15[B]:TRP:CH2	2:B:20:ILE:HD11	2.56	0.41
2:B:88:LEU:HD22	2:B:92:HIS:CE1	2.57	0.40
2:B:15[A]:TRP:CG	2:B:75:VAL:HG21	2.55	0.40

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{entiles}$
1	A	143/143 (100%)	139 (97%)	3 (2%)	1 (1%)	22	26
2	В	145/146~(99%)	143 (99%)	2 (1%)	0	100	100
All	All	288/289 (100%)	282 (98%)	5 (2%)	1 (0%)	34	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	\mathbf{ntiles}
1	A	112/110 (102%)	102 (91%)	10 (9%)	9	11
2	В	119/117~(102%)	111 (93%)	8 (7%)	16	21
All	All	$231/227 \ (102\%)$	213 (92%)	18 (8%)	12	16

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	SER
1	A	29	LEU
1	A	64	MET
1	A	69	LYS
1	A	82	SER

Continued on next page...



Continued from previous page...

7.6			1 0
Mol	Chain	${f Res}$	\mathbf{Type}
1	A	84	LEU
1	A	86	ASP
1	A	87	LEU
1	A	93	ARG
1	A	102	LEU
2	В	1	VAL
2	В	20	ILE
2	В	40	ARG
2	В	71	LEU
2	В	82	LEU
2	В	91	THR
2	В	110	LEU
2	В	144	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	${f Res}$	\mathbf{Type}
2	В	102	ASN
2	В	131	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res Link		Bo	ond leng	ths	В	ond ang	les
10101	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	В	148	2	27,50,50	1.95	6 (22%)	17,82,82	2.41	8 (47%)
3	HEM	A	143	1	27,50,50	1.92	6 (22%)	17,82,82	2.20	9 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	В	148	2	-	1/6/54/54	-
3	HEM	A	143	1	-	4/6/54/54	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	В	148	HEM	C3C-C2C	-5.44	1.32	1.40
3	A	143	HEM	C3B-C2B	-4.57	1.34	1.40
3	A	143	HEM	C3C-C2C	-4.31	1.34	1.40
3	В	148	HEM	C3B-C2B	-3.77	1.35	1.40
3	В	148	HEM	C3C-CAC	3.43	1.54	1.47
3	A	143	HEM	C3C-CAC	3.17	1.54	1.47
3	A	143	HEM	C3B-CAB	3.05	1.54	1.47
3	В	148	HEM	C3B-CAB	2.79	1.53	1.47
3	В	148	HEM	CAD-C3D	2.33	1.56	1.52
3	A	143	HEM	CAD-C3D	2.31	1.56	1.52
3	A	143	HEM	CMB-C2B	2.12	1.56	1.51
3	В	148	HEM	C4B-NB	2.07	1.40	1.36

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	${f Atoms}$	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
3	В	148	HEM	CAA-CBA-CGA	4.72	120.59	112.67
3	В	148	HEM	CBA-CAA-C2A	3.96	119.80	112.49
3	В	148	HEM	CMA-C3A-C4A	-3.96	122.37	128.46
3	A	143	HEM	CBA-CAA-C2A	3.56	119.06	112.49
3	A	143	HEM	CMA-C3A-C4A	-3.55	123.00	128.46
3	В	148	HEM	CMD-C2D-C1D	-3.49	123.10	128.46

Continued on next page...



$\alpha \cdots$	· ·	•	
Continued	trom	nromanne	naae
-	110111	picolous	payc

Mol	Chain	Res	Type	${f Atoms}$	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
3	A	143	HEM	CMD-C2D-C1D	-3.33	123.34	128.46
3	A	143	HEM	CAA-CBA-CGA	3.23	118.08	112.67
3	A	143	HEM	CAD-CBD-CGD	3.10	117.88	112.67
3	В	148	HEM	CMA-C3A-C2A	2.84	130.30	124.94
3	В	148	HEM	CMD-C2D-C3D	2.59	129.83	124.94
3	A	143	HEM	CMB-C2B-C3B	2.52	129.39	124.68
3	В	148	HEM	CMB-C2B-C3B	2.50	129.35	124.68
3	A	143	HEM	CMD-C2D-C3D	2.40	129.47	124.94
3	В	148	HEM	CMC-C2C-C3C	2.35	129.07	124.68
3	A	143	HEM	CMA-C3A-C2A	2.29	129.26	124.94
3	A	143	HEM	CMC-C2C-C3C	2.28	128.95	124.68

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	143	HEM	C1A-C2A-CAA-CBA
3	A	143	HEM	C3A-C2A-CAA-CBA
3	A	143	HEM	C2D-C3D-CAD-CBD
3	A	143	HEM	C4D-C3D-CAD-CBD
3	В	148	HEM	C3A-C2A-CAA-CBA


There are no ring outliers.

2 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	148	HEM	4	0
3	A	143	HEM	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

