

Full wwPDB X-ray Structure Validation Report (i)

Oct 5, 2023 – 10:25 PM EDT

PDB ID	:	6P1Z
Title	:	Bacteriophage phiKZ gp163.1 PAAR repeat protein in complex with the C-
		terminal part of the T4 gp5 beta-helical domain
Authors	:	Buth, S.A.; Shneider, M.M.; Leiman, P.G.
Deposited on	:	2019-05-20
Resolution	:	2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics Refmac CCP4 Ideal geometry (proteins) Ideal geometry (DNA, RNA)	: : : : :	20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove) Engh & Huber (2001) Parkinson et al. (1996)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	5197(2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	96	95% •	•
1	В	96	94% •	•
1	С	96	93% •	•
1	Е	96	92% •	•
1	F	96	93%	•



Continued from previous page...

Mol	Chain		Quality of chain	
1	G	96	91%	5% •
2	D	88	% 98%	••
2	Н	88	94%	5%•



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 11666 atoms, of which 5342 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues			Aton	ns			ZeroOcc	AltConf	Trace
1	А	92	Total	С	Н	Ν	Ο	\mathbf{S}	0	2	0
	A	92	1333	413	656	113	149	2	0	2	0
1	В	92	Total	С	Η	Ν	0	S	0	0	0
	D	92	1316	408	646	112	148	2	0	0	0
1	С	92	Total	С	Н	Ν	Ο	S	0	1	0
		92	1317	408	647	112	148	2	0		0
1	Е	92	Total	С	Н	Ν	0	S	0	4	0
	Ľ	92	1358	420	668	115	153	2	0	4	0
1	F	92	Total	С	Н	Ν	Ο	S	0	2	0
	I.	92	1329	412	651	113	151	2	0	2	0
1	G	92	Total	С	Η	Ν	Ο	S	0	4	0
	G	92	1345	416	662	114	151	2		4	0

• Molecule 1 is a protein called Baseplate central spike complex protein gp5.

There are 24 discrepancies between the modelled and reference sequences:

Residue	Modelled	Actual	$\operatorname{Comment}$	Reference
480	GLY	-	expression tag	UNP P16009
481	SER	-	expression tag	UNP P16009
482	GLY	-	expression tag	UNP P16009
483	SER	-	expression tag	UNP P16009
480	GLY	-	expression tag	UNP P16009
481	SER	-	expression tag	UNP P16009
482	GLY	-	expression tag	UNP P16009
483	SER	-	expression tag	UNP P16009
480	GLY	-	expression tag	UNP P16009
481	SER	-	expression tag	UNP P16009
482	GLY	-	expression tag	UNP P16009
483	SER	-	expression tag	UNP P16009
480	GLY	-	expression tag	UNP P16009
481	SER	-	expression tag	UNP P16009
482	GLY	-	expression tag	UNP P16009
483	SER	-	expression tag	UNP P16009
480	GLY	-	expression tag	UNP P16009
	481 482 483 480 481 482 483 480 481 482 483 480 481 482 483	481 SER 482 GLY 483 SER 480 GLY 481 SER 482 GLY 483 SER 480 GLY 481 SER 480 GLY 483 SER 484 SER 485 GLY 486 GLY 487 SER 488 SER 481 SER 482 GLY 483 SER 483 SER 483 SER 483 SER	481 SER - 482 GLY - 483 SER - 480 GLY - 481 SER - 482 GLY - 483 SER - 483 SER - 483 SER - 480 GLY - 481 SER - 483 SER - 483 SER - 483 SER - 481 SER - 483 SER -	481SER-expression tag482GLY-expression tag483SER-expression tag480GLY-expression tag481SER-expression tag482GLY-expression tag483SER-expression tag484GLY-expression tag485SER-expression tag480GLY-expression tag481SER-expression tag482GLY-expression tag483SER-expression tag481SER-expression tag481SER-expression tag481SER-expression tag481SER-expression tag483SER-expression tag </td



6P1Z

Commu	ea from pre	vious page			
Chain	Residue	Modelled	Actual	Comment	Reference
F	481	SER	-	expression tag	UNP P16009
F	482	GLY	-	expression tag	UNP P16009
F	483	SER	-	expression tag	UNP P16009
G	480	GLY	-	expression tag	UNP P16009
G	481	SER	-	expression tag	UNP P16009
G	482	GLY	-	expression tag	UNP P16009
G	483	SER	-	expression tag	UNP P16009


Continued from previous page...

• Molecule 2 is a protein called PAAR-repeat central spike tip protein.

Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace
2	а	87	Total	С	Η	Ν	0	S	0	2	
	2 D		1220	375	612	112	110	11	0	2	0
0	ц	87	Total	С	Η	Ν	0	S	0	0	0
	2 H	01	1202	370	602	111	109	10		0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Mg 1 1	0	0
3	Ε	1	Total Mg 1 1	0	0





Mol	Chain	Residues	I	Aton	ns		ZeroOcc	AltConf
4	В	1	Total				0	0
			49 Total					
4	Ε	1	49	-	31	$\frac{0}{2}$	0	0


• Molecule 5 is STEARIC ACID (three-letter code: STE) (formula: $C_{18}H_{36}O_2$).



Mo	Chain	Residues	Atoms				ZeroOcc	AltConf
5	С	1	Total	С	Η	0	0	0
	5 0	1	55	18	35	2	0	
5	С	1	Total	С	Η	0	0	0
5	G	L	55	18	35	2	0	

• Molecule 6 is 9-OCTADECENOIC ACID (three-letter code: ELA) (formula: $C_{18}H_{34}O_2$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	С	1	Total	С	Η	Ο	0	0
0	0 0	1	53	18	33	2	0	0
6	E	1	Total	С	Η	Ο	0	0
0		L	53	18	33	2	0	

• Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	1	Total Zn 1 1	0	0
7	Н	1	Total Zn 1 1	0	0

• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	155	Total O 155 155	0	0
8	В	121	Total O 121 121	0	0
8	С	135	Total O 135 135	0	0
8	D	93	Total O 93 93	0	0
8	Е	113	Total O 113 113	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	\mathbf{F}	108	Total O 108 108	0	0
8	G	114	Total O 114 114	0	0
8	Н	89	Total O 89 89	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Baseplate central spike complex protein gp5



Chain G:




5%

91%



• Molecule 2: PAAR-repeat central spike tip protein



• Molecule 2: PAAR-repeat central spike tip protein

Chain H:	94%	5% •
MET P2 P18 K23 P82 888		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	72.28Å 69.12Å 77.80Å	Depositor
a, b, c, α , β , γ	90.00° 102.48° 90.00°	Depositor
Resolution (Å)	44.60 - 2.10	Depositor
Resolution (A)	37.98 - 2.10	EDS
% Data completeness	95.2 (44.60-2.10)	Depositor
(in resolution range)	95.0 (37.98-2.10)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.82 (at 2.10 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
D D.	0.188 , 0.251	Depositor
R, R_{free}	0.189 , 0.251	DCC
R_{free} test set	1459 reflections (3.50%)	wwPDB-VP
Wilson B-factor $(Å^2)$	20.6	Xtriage
Anisotropy	0.567	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.38 , 52.2	EDS
L-test for twinning ²	$ \langle L \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11666	wwPDB-VP
Average B, all atoms $(Å^2)$	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 65.35 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.1864e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PLM, ELA, MG, STE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		lengths	Bond angles	
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.23	0/690	0.47	0/937
1	В	0.24	0/676	0.48	0/917
1	С	0.24	0/680	0.48	0/922
1	Е	0.23	0/703	0.47	0/955
1	F	0.23	0/692	0.47	0/939
1	G	0.23	0/702	0.47	0/953
2	D	0.25	0/625	0.46	0/843
2	Н	0.26	0/612	0.45	0/826
All	All	0.24	0/5380	0.47	0/7292

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	677	656	648	1	0
1	В	670	646	646	2	0
1	С	670	647	643	3	0
1	Е	690	668	658	3	0
1	F	678	651	639	4	0



6P1Z	
------	--

Mol	Chain	<i>n previous</i> Non-H	H(model)	H(added)	Clashes	Symm-Clashes
			· · · ·	· · ·		ů.
1	G	683	662	648	6	0
2	D	608	612	606	0	0
2	Н	600	602	602	4	0
3	А	1	0	0	0	0
3	Е	1	0	0	0	0
4	В	18	31	31	0	0
4	Е	18	31	31	2	0
5	С	20	35	35	2	0
5	G	20	35	35	0	0
6	С	20	33	33	2	0
6	Е	20	33	33	0	0
7	D	1	0	0	0	0
7	Н	1	0	0	0	0
8	А	155	0	0	0	0
8	В	121	0	0	0	1
8	С	135	0	0	0	0
8	D	93	0	0	0	1
8	Ε	113	0	0	1	0
8	F	108	0	0	4	0
8	G	114	0	0	1	0
8	Н	89	0	0	3	0
All	All	6324	5342	5288	20	1

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:484:GLY:N	8:E:702:HOH:O	2.30	0.64
2:H:18:PRO:O	8:H:201:HOH:O	2.14	0.64
1:F:570:SER:OG	8:F:601:HOH:O	2.17	0.60
1:G:484:GLY:N	8:G:702:HOH:O	2.37	0.57
1:F:484:GLY:N	8:F:605:HOH:O	2.36	0.57
2:H:44:THR:O	8:H:202:HOH:O	2.17	0.57
1:F:484:GLY:N	8:F:606:HOH:O	2.39	0.55
1:G:572:ILE:HD12	1:G:572:ILE:N	2.25	0.52
1:E:498:VAL:HG11	1:E:502[B]:VAL:HG12	1.93	0.51
1:B:504:ILE:HD11	6:C:602:ELA:H71	1.94	0.48
4:E:603:PLM:H41	1:G:512:ILE:HD11	1.97	0.46
1:C:504:ILE:HD11	5:C:601:STE:H71	1.99	0.44



6P	'1	Ζ

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:504:ILE:HG21	5:C:601:STE:H82	1.98	0.43
1:B:494:GLY:O	1:C:502:VAL:HA	2.20	0.41
1:E:561[B]:SER:OG	1:G:556:SER:HA	2.20	0.41
2:H:82:PRO:O	8:H:203:HOH:O	2.22	0.41
4:E:603:PLM:H72	1:G:504:ILE:HD13	2.03	0.41
1:C:512:ILE:HD11	6:C:602:ELA:H31	2.03	0.40
1:G:572:ILE:HD13	2:H:23:LYS:HG3	2.03	0.40
1:F:511[B]:ASP:OD2	8:F:602:HOH:O	2.21	0.40

Continued from previous page...

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:B:770:HOH:O	8:D:265:HOH:O[2_847]	2.14	0.06

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	92/96~(96%)	92 (100%)	0	0	100 100
1	В	90/96~(94%)	90 (100%)	0	0	100 100
1	С	91/96~(95%)	91 (100%)	0	0	100 100
1	Ε	94/96~(98%)	94 (100%)	0	0	100 100
1	F	92/96~(96%)	90~(98%)	2(2%)	0	100 100
1	G	94/96~(98%)	94 (100%)	0	0	100 100
2	D	87/88~(99%)	82 (94%)	5~(6%)	0	100 100
2	Н	85/88~(97%)	81 (95%)	4(5%)	0	100 100
All	All	725/752~(96%)	714 (98%)	11 (2%)	0	100 100



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	76/76~(100%)	76~(100%)	0	100 100
1	В	74/76~(97%)	74 (100%)	0	100 100
1	С	74/76~(97%)	74 (100%)	0	100 100
1	Ε	78/76~(103%)	78~(100%)	0	100 100
1	F	76/76~(100%)	76~(100%)	0	100 100
1	G	78/76~(103%)	78~(100%)	0	100 100
2	D	64/64~(100%)	63~(98%)	1 (2%)	62 69
2	Н	63/64~(98%)	63~(100%)	0	100 100
All	All	583/584~(100%)	582 (100%)	1 (0%)	93 96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type	
2	D	23	LYS	

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
4	PLM	Е	603	-	17,17,17	0.60	0	17,17,17	1.02	0
6	ELA	С	602	-	19,19,19	0.58	0	19,19,19	0.72	0
6	ELA	Е	602	-	19,19,19	0.57	0	19,19,19	0.67	0
4	PLM	В	601	-	17,17,17	0.58	0	17,17,17	1.06	0
5	STE	С	601	-	19,19,19	0.56	0	19,19,19	1.10	1 (5%)
5	STE	G	601	-	19,19,19	0.57	0	19,19,19	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PLM	Е	603	-	-	11/15/15/15	-
6	ELA	С	602	-	-	13/17/17/17	-
6	ELA	Е	602	-	-	12/17/17/17	-
4	PLM	В	601	-	-	9/15/15/15	-
5	STE	С	601	-	-	9/17/17/17	-
5	STE	G	601	-	-	10/17/17/17	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	С	601	STE	C3-C2-C1	-2.21	108.90	114.47

There are no chirality outliers.

All (64) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	Е	603	PLM	C6-C7-C8-C9
4	Е	603	PLM	C1-C2-C3-C4
6	Е	602	ELA	C4-C3-CA-C
6	Е	602	ELA	C5-C6-C7-C8
4	В	601	PLM	C8-C9-CA-CB
6	Е	602	ELA	C13-C14-C15-C16
5	G	601	STE	C4-C5-C6-C7
4	Е	603	PLM	C8-C9-CA-CB
4	Е	603	PLM	C3-C4-C5-C6
4	Е	603	PLM	C7-C8-C9-CA
6	Е	602	ELA	C12-C13-C14-C15
4	В	601	PLM	C6-C7-C8-C9
5	С	601	STE	C7-C8-C9-C10
6	С	602	ELA	CA-C3-C4-C5
5	G	601	STE	C6-C7-C8-C9
6	С	602	ELA	C5-C6-C7-C8
6	С	602	ELA	C14-C15-C16-C17
6	Е	602	ELA	C3-C4-C5-C6
4	Е	603	PLM	CA-CB-CC-CD
6	Е	602	ELA	C6-C7-C8-C9
5	G	601	STE	C12-C13-C14-C15
5	G	601	STE	C1-C2-C3-C4
4	В	601	PLM	C7-C8-C9-CA
4	В	601	PLM	C9-CA-CB-CC
5	С	601	STE	C14-C15-C16-C17
5	С	601	STE	C5-C6-C7-C8
5	С	601	STE	C10-C11-C12-C13
5	G	601	STE	C14-C15-C16-C17
6	С	602	ELA	C11-C12-C13-C14
4	Е	603	PLM	CB-CC-CD-CE
4	В	601	PLM	CB-CC-CD-CE
6	С	602	ELA	C3-C4-C5-C6
4	В	601	PLM	CD-CE-CF-CG
5	G	601	STE	C3-C4-C5-C6
4	В	601	PLM	C2-C3-C4-C5
5	G	601	STE	C2-C3-C4-C5
6	С	602	ELA	C15-C16-C17-C18



Mol	Chain	Res	Type	Atoms
5	С	601	STE	С11-С10-С9-С8
6	С	602	ELA	C10-C11-C12-C13
6	Е	602	ELA	C4-C5-C6-C7
6	Е	602	ELA	CA-C3-C4-C5
6	С	602	ELA	C13-C14-C15-C16
4	Е	603	PLM	C5-C6-C7-C8
4	В	601	PLM	C1-C2-C3-C4
5	G	601	STE	O1-C1-C2-C3
4	Е	603	PLM	O2-C1-C2-C3
5	С	601	STE	O1-C1-C2-C3
6	Е	602	ELA	C14-C15-C16-C17
5	С	601	STE	O2-C1-C2-C3
5	G	601	STE	O2-C1-C2-C3
6	С	602	ELA	C4-C3-CA-C
5	С	601	STE	C11-C12-C13-C14
4	Ε	603	PLM	O1-C1-C2-C3
6	С	602	ELA	C9-C10-C11-C12
6	Ε	602	ELA	OXT-C-CA-C3
6	С	602	ELA	O-C-CA-C3
6	С	602	ELA	C4-C5-C6-C7
5	G	601	STE	C13-C14-C15-C16
4	Е	603	PLM	CD-CE-CF-CG
6	С	602	ELA	OXT-C-CA-C3
6	Ε	602	ELA	O-C-CA-C3
6	Е	602	ELA	C9-C10-C11-C12
4	В	601	PLM	C4-C5-C6-C7
5	С	601	STE	C2-C3-C4-C5

Continued from previous page...

There are no ring outliers.


3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Е	603	PLM	2	0
6	С	602	ELA	2	0
5	С	601	STE	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	$\mathbf{Q}{<}0.9$
1	А	92/96~(95%)	-0.35	0 100 100	16, 22, 31, 51	0
1	В	92/96~(95%)	-0.35	0 100 100	16, 22, 34, 44	0
1	С	92/96~(95%)	-0.35	0 100 100	16, 22, 34, 42	0
1	Е	92/96~(95%)	-0.12	0 100 100	17, 30, 42, 59	0
1	F	92/96~(95%)	-0.24	0 100 100	21, 29, 45, 64	0
1	G	92/96~(95%)	-0.22	0 100 100	20, 29, 45, 58	0
2	D	87/88~(98%)	-0.38	1 (1%) 80 84	14, 21, 33, 84	0
2	Н	87/88~(98%)	-0.25	0 100 100	20, 28, 47, 80	0
All	All	726/752~(96%)	-0.28	1 (0%) 95 96	14, 25, 41, 84	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	43	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




6P1Z


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
6	ELA	Е	602	20/20	0.80	0.27	43,69,91,98	0
6	ELA	С	602	20/20	0.81	0.23	44,63,94,102	0
4	PLM	Е	603	18/18	0.85	0.35	48,70,87,88	0
5	STE	G	601	20/20	0.87	0.21	38,64,94,96	0
4	PLM	В	601	18/18	0.88	0.23	$38,\!65,\!91,\!96$	0
5	STE	С	601	20/20	0.90	0.16	31,63,80,88	0
3	MG	А	601	1/1	0.96	0.06	$15,\!15,\!15,\!15$	0
3	MG	Е	601	1/1	0.96	0.05	18,18,18,18	0
7	ZN	D	101	1/1	0.99	0.05	14,14,14,14	0
7	ZN	Н	101	1/1	0.99	0.02	24,24,24,24	0

median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















6.5 Other polymers (i)

There are no such residues in this entry.

