



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2024 – 08:12 AM EST

PDB ID : 4P38  
Title : Human 11beta-Hydroxysteroid Dehydrogenase Type 1 in complex with AZD8329  
Authors : Ogg, D.; Hargreaves, D.; Gerhardt, S.  
Deposited on : 2014-03-06  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	 2% 85% 14% .
1	B	265	 3% 82% 16% ..

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4266 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Corticosteroid 11-beta-dehydrogenase isozyme 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	265	2037	1299	346	378	14	0	0	0
1	B	263	2040	1299	348	379	14	0	1	0

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	179	LEU	MET	conflict	UNP P28845
A	262	ARG	LEU	conflict	UNP P28845
A	272	SER	CYS	conflict	UNP P28845
A	278	GLU	PHE	conflict	UNP P28845
A	286	TRP	MET	conflict	UNP P28845
B	179	LEU	MET	conflict	UNP P28845
B	262	ARG	LEU	conflict	UNP P28845
B	272	SER	CYS	conflict	UNP P28845
B	278	GLU	PHE	conflict	UNP P28845
B	286	TRP	MET	conflict	UNP P28845

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	48	21	7	17	3	0	0
2	B	1	48	21	7	17	3	0	0

- Molecule 3 is 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]benzoic acid (three-letter code: 21T) (formula: C<sub>25</sub>H<sub>31</sub>N<sub>3</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	31	25	3	3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	B	1	31	25	3	3	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	B	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	19	Total	O	0	0
			19	19		
5	B	10	Total	O	0	0
			10	10		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Corticosteroid 11-beta-dehydrogenase isozyme 1



- Molecule 1: Corticosteroid 11-beta-dehydrogenase isozyme 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.52Å 103.52Å 134.78Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.27 – 2.80 29.17 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.7 (30.27-2.80) 98.7 (29.17-2.80)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 2.80Å)	Xtrriage
Refinement program	BUSTER 2.11.2	Depositor
R, $R_{free}$	0.227 , 0.272 0.222 , 0.261	Depositor DCC
$R_{free}$ test set	1062 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	75.7	Xtrriage
Anisotropy	0.353	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 49.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.56$ , $\langle L^2 \rangle = 0.41$	Xtrriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4266	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 21T, CL, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/2073	0.75	0/2801
1	B	0.54	0/2075	0.75	0/2801
All	All	0.53	0/4148	0.75	0/5602

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2037	0	2068	24	0
1	B	2040	0	2069	28	0
2	A	48	0	26	1	0
2	B	48	0	26	3	0
3	A	31	0	30	3	0
3	B	31	0	30	7	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	19	0	0	1	0
5	B	10	0	0	0	0
All	All	4266	0	4249	55	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:218:ILE:CD1	1:B:241:CYS:SG	2.45	1.05
1:B:218:ILE:HD12	1:B:241:CYS:SG	2.03	0.98
1:A:270:ASN:HD21	1:A:272:SER:HB2	1.55	0.72
1:B:231:VAL:HG23	1:B:232:HIS:N	2.08	0.69
1:B:218:ILE:HD11	1:B:241:CYS:SG	2.35	0.65
1:A:175:VAL:HG22	1:B:273:ARG:HB2	1.79	0.63
1:A:36:LYS:HG2	1:A:110:MET:HB3	1.84	0.60
3:A:302:21T:N3	3:A:302:21T:H21	2.17	0.59
1:B:76:SER:O	1:B:80:GLU:HG2	2.04	0.58
1:B:36:LYS:HG2	1:B:110:MET:HB3	1.84	0.57
3:B:302:21T:H6	3:B:302:21T:C8	2.37	0.54
1:B:67:SER:OG	2:B:301:NDP:O2X	2.25	0.54
1:A:273:ARG:HG3	1:B:175:VAL:HG22	1.91	0.52
3:A:302:21T:H6	3:A:302:21T:C8	2.40	0.52
1:A:171:LEU:HD21	1:A:264:THR:HG21	1.93	0.51
1:B:218:ILE:HD13	1:B:241:CYS:HB2	1.92	0.51
3:A:302:21T:H21	3:A:302:21T:C15	2.40	0.50
1:B:171:LEU:HD21	1:B:264:THR:HG21	1.94	0.50
1:A:70:THR:HA	1:A:73:LYS:HE3	1.94	0.50
1:A:48:ARG:HD3	1:A:52:TYR:CE2	2.48	0.48
1:A:277:GLU:HG3	1:B:175:VAL:HG13	1.96	0.48
1:B:70:THR:HA	1:B:73:LYS:HE3	1.97	0.47
1:B:218:ILE:HD13	1:B:241:CYS:CB	2.45	0.47
1:A:220:THR:O	1:A:224:MET:HG2	2.15	0.46
3:B:302:21T:H21	3:B:302:21T:C15	2.45	0.46
1:A:199:LYS:HE2	5:A:418:HOH:O	2.15	0.46
1:A:37:VAL:HG13	1:A:115:MET:HB3	1.98	0.46
2:B:301:NDP:H5N	3:B:302:21T:H7	1.98	0.45
1:A:272:SER:HB3	1:B:267:LEU:O	2.17	0.45
1:A:77:HIS:HA	1:A:80:GLU:HG3	1.99	0.45
1:A:140:MET:HG2	1:B:140:MET:HG2	1.99	0.45
1:A:216:GLY:O	1:A:218:ILE:HG12	2.16	0.45
1:B:218:ILE:CD1	1:B:241:CYS:CB	2.95	0.44
1:A:175:VAL:HG13	1:B:277:GLU:HG3	1.99	0.44
1:B:105:GLN:O	1:B:109:LEU:HG	2.18	0.44
1:B:37:VAL:HG13	1:B:115:MET:HB3	1.99	0.43
3:B:302:21T:H21	3:B:302:21T:H29	2.01	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:PHE:O	1:A:147:TYR:HB2	2.19	0.43
1:A:267:LEU:O	1:B:272:SER:HB3	2.19	0.43
1:B:42:ALA:HB3	1:B:63:VAL:HB	2.01	0.42
1:B:248:GLY:HA3	1:B:256:VAL:HG21	2.00	0.42
3:B:302:21T:C8	3:B:302:21T:C3	2.97	0.42
1:B:194:PHE:HB3	1:B:210:ILE:HG21	2.01	0.42
1:A:194:PHE:HB3	1:A:210:ILE:HG21	2.01	0.42
1:B:171:LEU:HD22	1:B:216:GLY:HA2	2.01	0.42
1:B:75:VAL:HG21	1:B:88:TYR:HB3	2.02	0.42
1:B:231:VAL:HG23	1:B:232:HIS:H	1.79	0.41
3:B:302:21T:H6	3:B:302:21T:C9	2.49	0.41
2:B:301:NDP:H41N	3:B:302:21T:N3	2.36	0.41
1:A:270:ASN:ND2	1:A:272:SER:HB2	2.29	0.41
1:A:121:ILE:HG12	2:A:301:NDP:H3D	2.03	0.41
1:A:48:ARG:HD3	1:A:52:TYR:CZ	2.56	0.40
1:A:203:VAL:HG21	1:A:286:TRP:HB2	2.03	0.40
1:B:119:ASN:HD22	1:B:168:VAL:HG21	1.87	0.40
1:A:285:ASN:HD21	1:A:287:ASP:HB2	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/265 (99%)	247 (94%)	16 (6%)	0	100	100
1	B	260/265 (98%)	248 (95%)	12 (5%)	0	100	100
All	All	523/530 (99%)	495 (95%)	28 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/222 (99%)	208 (94%)	12 (6%)	21	52
1	B	221/222 (100%)	204 (92%)	17 (8%)	13	35
All	All	441/444 (99%)	412 (93%)	29 (7%)	16	44

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	ARG
1	A	97	THR
1	A	171	LEU
1	A	203	VAL
1	A	221	GLU
1	A	228	SER
1	A	239	GLU
1	A	262	ARG
1	A	274	LYS
1	A	276	LEU
1	A	282	THR
1	A	285	ASN
1	B	28	ARG
1	B	48	ARG
1	B	67	SER
1	B	68	LYS
1	B	97	THR
1	B	171	LEU
1	B	197	ILE
1	B	203	VAL
1	B	205	ARG
1	B	228	SER
1	B	232	HIS
1	B	262	ARG
1	B	273	ARG
1	B	274	LYS
1	B	282	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	285	ASN
1	B	288	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	A	101	GLN
1	A	123	ASN
1	A	253	GLN
1	A	270	ASN
1	A	285	ASN
1	B	77	HIS
1	B	101	GLN
1	B	119	ASN
1	B	123	ASN
1	B	285	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	21T	A	302	-	32,35,35	0.82	1 (3%)	47,54,54	0.92	3 (6%)
2	NDP	A	301	-	45,52,52	1.31	5 (11%)	53,80,80	0.98	3 (5%)
2	NDP	B	301	-	45,52,52	1.26	3 (6%)	53,80,80	0.85	1 (1%)
3	21T	B	302	-	32,35,35	0.83	1 (3%)	47,54,54	1.00	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	21T	A	302	-	-	8/20/50/50	0/6/5/5
2	NDP	A	301	-	-	7/30/77/77	0/5/5/5
2	NDP	B	301	-	-	6/30/77/77	0/5/5/5
3	21T	B	302	-	-	8/20/50/50	0/6/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	NDP	C4N-C3N	-5.22	1.39	1.49
2	A	301	NDP	C4N-C3N	-4.92	1.40	1.49
2	B	301	NDP	C4N-C5N	-3.76	1.39	1.48
2	A	301	NDP	C4N-C5N	-3.52	1.39	1.48
2	A	301	NDP	C6N-C5N	3.51	1.39	1.33
2	B	301	NDP	C6N-C5N	3.22	1.39	1.33
3	B	302	21T	C6-C15	-3.05	1.44	1.50
3	A	302	21T	C6-C15	-3.00	1.44	1.50
2	A	301	NDP	P2B-O1X	-2.36	1.42	1.50
2	A	301	NDP	C1D-N1N	2.23	1.52	1.46

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	21T	C6-C15-N3	-3.68	109.99	116.80
3	A	302	21T	C6-C15-N3	-2.58	112.02	116.80
3	A	302	21T	C8-N2-C5	2.47	132.55	129.90
3	B	302	21T	C8-N2-C5	2.39	132.46	129.90

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	NDP	C3N-C2N-N1N	-2.34	119.75	123.10
2	A	301	NDP	C5A-C6A-N6A	2.33	123.89	120.35
2	A	301	NDP	O3X-P2B-O2B	-2.28	95.76	105.99
3	A	302	21T	O1-C14-C11	-2.19	115.63	121.45
3	B	302	21T	O1-C14-C11	-2.18	115.65	121.45
3	B	302	21T	O3-C15-N3	2.17	126.45	122.45
2	B	301	NDP	C5A-C6A-N6A	2.00	123.39	120.35

There are no chirality outliers.

All (29) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	301	NDP	C5D-O5D-PN-O2N
3	A	302	21T	C1-C2-C5-C6
3	A	302	21T	C3-C2-C5-C6
3	A	302	21T	C4-C2-C5-C6
3	A	302	21T	O3-C15-C6-C7
3	A	302	21T	N3-C15-C6-C7
3	B	302	21T	C1-C2-C5-C6
3	B	302	21T	C3-C2-C5-C6
3	B	302	21T	C4-C2-C5-C6
3	B	302	21T	O3-C15-C6-C7
3	B	302	21T	N3-C15-C6-C7
2	A	301	NDP	O4D-C1D-N1N-C6N
2	B	301	NDP	O4D-C1D-N1N-C6N
3	A	302	21T	C1-C2-C5-N2
3	B	302	21T	C1-C2-C5-N2
3	B	302	21T	C3-C2-C5-N2
3	B	302	21T	C4-C2-C5-N2
2	A	301	NDP	C1B-C2B-O2B-P2B
2	B	301	NDP	C5D-O5D-PN-O3
2	B	301	NDP	C5D-O5D-PN-O1N
2	A	301	NDP	PN-O3-PA-O1A
3	A	302	21T	C3-C2-C5-N2
3	A	302	21T	C4-C2-C5-N2
2	A	301	NDP	PN-O3-PA-O2A
2	A	301	NDP	C3B-C2B-O2B-P2B
2	B	301	NDP	O4B-C4B-C5B-O5B
2	A	301	NDP	C2B-O2B-P2B-O3X
2	A	301	NDP	O4B-C4B-C5B-O5B
2	B	301	NDP	C2N-C3N-C7N-N7N

There are no ring outliers.


4 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	21T	3	0
2	A	301	NDP	1	0
2	B	301	NDP	3	0
3	B	302	21T	7	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	265/265 (100%)	0.01	6 (2%) 60 51	51, 73, 100, 132	0
1	B	263/265 (99%)	-0.06	8 (3%) 50 40	50, 72, 102, 121	0
All	All	528/530 (99%)	-0.02	14 (2%) 54 44	50, 73, 102, 132	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	167	VAL	2.9
1	A	118	LEU	2.8
1	A	289	PHE	2.6
1	A	117	ILE	2.5
1	B	160	GLN	2.4
1	B	28	ARG	2.3
1	B	166	VAL	2.3
1	A	134	HIS	2.2
1	B	80	GLU	2.2
1	B	117	ILE	2.2
1	B	191	ASP	2.1
1	A	88	TYR	2.1
1	A	116	LEU	2.1
1	B	263	TRP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	21T	A	302	31/31	0.94	0.21	60,68,83,88	0
4	CL	B	303	1/1	0.94	0.08	75,75,75,75	0
3	21T	B	302	31/31	0.95	0.18	57,69,83,91	0
4	CL	A	303	1/1	0.96	0.10	71,71,71,71	0
2	NDP	A	301	48/48	0.96	0.16	55,64,70,71	0
2	NDP	B	301	48/48	0.98	0.15	54,64,71,72	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around 21T B 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NDP A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.