



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2023 – 03:19 PM EDT

PDB ID : 3P4T
Title : Crystal structure of a putative acyl-CoA dehydrogenase from *Mycobacterium smegmatis*
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2010-10-07
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	 4% 87% 7% 5%
1	B	403	 3% 91% 5% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FAO	A	500	X	-	-	-
3	FAO	B	500	X	-	-	-

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6703 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative acyl-CoA dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	381	2904	1810	526	556	12	0	9	0
1	B	381	2903	1809	527	555	12	0	9	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP A0R3J1
A	2	ALA	-	expression tag	UNP A0R3J1
A	3	HIS	-	expression tag	UNP A0R3J1
A	4	HIS	-	expression tag	UNP A0R3J1
A	5	HIS	-	expression tag	UNP A0R3J1
A	6	HIS	-	expression tag	UNP A0R3J1
A	7	HIS	-	expression tag	UNP A0R3J1
A	8	HIS	-	expression tag	UNP A0R3J1
A	9	MET	-	expression tag	UNP A0R3J1
A	10	GLY	-	expression tag	UNP A0R3J1
A	11	THR	-	expression tag	UNP A0R3J1
A	12	LEU	-	expression tag	UNP A0R3J1
A	13	GLU	-	expression tag	UNP A0R3J1
A	14	ALA	-	expression tag	UNP A0R3J1
A	15	GLN	-	expression tag	UNP A0R3J1
A	16	THR	-	expression tag	UNP A0R3J1
A	17	GLN	-	expression tag	UNP A0R3J1
A	18	GLY	-	expression tag	UNP A0R3J1
A	19	PRO	-	expression tag	UNP A0R3J1
A	20	GLY	-	expression tag	UNP A0R3J1
A	21	SER	-	expression tag	UNP A0R3J1
B	1	MET	-	expression tag	UNP A0R3J1
B	2	ALA	-	expression tag	UNP A0R3J1
B	3	HIS	-	expression tag	UNP A0R3J1
B	4	HIS	-	expression tag	UNP A0R3J1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	5	HIS	-	expression tag	UNP A0R3J1
B	6	HIS	-	expression tag	UNP A0R3J1
B	7	HIS	-	expression tag	UNP A0R3J1
B	8	HIS	-	expression tag	UNP A0R3J1
B	9	MET	-	expression tag	UNP A0R3J1
B	10	GLY	-	expression tag	UNP A0R3J1
B	11	THR	-	expression tag	UNP A0R3J1
B	12	LEU	-	expression tag	UNP A0R3J1
B	13	GLU	-	expression tag	UNP A0R3J1
B	14	ALA	-	expression tag	UNP A0R3J1
B	15	GLN	-	expression tag	UNP A0R3J1
B	16	THR	-	expression tag	UNP A0R3J1
B	17	GLN	-	expression tag	UNP A0R3J1
B	18	GLY	-	expression tag	UNP A0R3J1
B	19	PRO	-	expression tag	UNP A0R3J1
B	20	GLY	-	expression tag	UNP A0R3J1
B	21	SER	-	expression tag	UNP A0R3J1


- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is [(2R,3S,4R,5R)-5-(6-amino-9H-purin-9-yl)-3,4-dihydroxytetrahydrofuran-2-yl]

methyl (2R,3S,4S)-5-[(4aS,10aR)-7,8-dimethyl-2,4-dioxo-1,3,4,4a,5,10a-hexahydrobenzo[g]pteridin-10(2H)-yl]-2,3,4-trihydroxypentyl dihydrogen diphosphate (three-letter code: FAO) (formula: C₂₇H₃₇N₉O₁₅P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	387	Total	O	0	1
			388	388		
4	B	393	Total	O	0	1
			394	394		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative acyl-CoA dehydrogenase



- Molecule 1: Putative acyl-CoA dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	161.93Å 64.75Å 83.81Å 90.00° 112.04° 90.00°	Depositor
Resolution (Å)	19.82 – 1.70 19.82 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.5 (19.82-1.70) 96.5 (19.82-1.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.21 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.153 , 0.185 0.160 , 0.193	Depositor DCC
R_{free} test set	4271 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtrriage
Anisotropy	0.388	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6703	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, FAO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/2981	0.80	5/4039 (0.1%)
1	B	0.72	0/2981	0.76	0/4040
All	All	0.73	0/5962	0.78	5/8079 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	226	ARG	NE-CZ-NH1	7.15	123.88	120.30
1	A	312	ARG	NE-CZ-NH1	6.34	123.47	120.30
1	A	226	ARG	NE-CZ-NH2	-6.16	117.22	120.30
1	A	30	ARG	NE-CZ-NH1	5.53	123.06	120.30
1	A	373	ARG	NE-CZ-NH1	-5.06	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2904	0	2881	22	0
1	B	2903	0	2866	23	0
2	A	4	0	6	0	0
2	B	4	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	53	0	33	6	0
3	B	53	0	33	9	0
4	A	388	0	0	3	0
4	B	394	0	0	2	0
All	All	6703	0	5825	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:380:ILE:HD11	3:B:500:FAO:C7M	1.73	1.18
1:B:380:ILE:HD11	3:B:500:FAO:H7M3	1.35	1.05
1:B:380:ILE:CD1	3:B:500:FAO:H7M3	1.88	1.03
1:A:380:ILE:HD11	3:A:500:FAO:H7M3	1.43	1.01
1:A:380:ILE:HD11	3:A:500:FAO:C7M	1.94	0.97
1:B:326[A]:ARG:NE	1:B:331[A]:GLU:OE1	2.06	0.89
1:B:380:ILE:CD1	3:B:500:FAO:C7M	2.48	0.83
1:A:380:ILE:CD1	3:A:500:FAO:H7M3	2.15	0.76
1:A:347:GLU:OE1	4:A:695:HOH:O	2.10	0.69
1:B:326[A]:ARG:HH21	1:B:331[A]:GLU:CD	1.98	0.67
1:B:380:ILE:HD11	3:B:500:FAO:C7	2.28	0.64
1:B:380:ILE:HD11	3:B:500:FAO:H7M1	1.72	0.63
1:A:199:ALA:HB1	1:A:253:PHE:CG	2.39	0.57
1:B:178:TYR:O	3:B:500:FAO:C4N	2.53	0.57
1:B:159:LEU:HD12	1:B:159:LEU:C	2.26	0.55
1:A:178:TYR:O	3:A:500:FAO:C4N	2.55	0.55
1:B:38[A]:ARG:NH2	4:B:567:HOH:O	2.38	0.54
1:A:322:HIS:NE2	1:A:326[A]:ARG:HD2	2.23	0.53
1:B:261:VAL:HG23	4:B:607:HOH:O	2.09	0.53
1:A:168:ASP:OD1	1:A:169:HIS:ND1	2.43	0.51
1:B:146:ALA:HA	1:B:179:ILE:HD12	1.92	0.51
1:A:380:ILE:CD1	3:A:500:FAO:C7M	2.79	0.50
1:B:64:ARG:O	1:B:68[A]:GLU:HG3	2.13	0.48
1:B:326[A]:ARG:CZ	1:B:331[A]:GLU:OE1	2.59	0.48
1:A:261:VAL:HG13	1:A:338:VAL:HG21	1.96	0.48
1:A:326[B]:ARG:NE	1:A:331:GLU:OE2	2.42	0.47
1:A:365:TYR:CE2	1:B:380:ILE:HD12	2.51	0.46
1:A:365:TYR:CZ	1:B:380:ILE:HB	2.52	0.46
3:A:500:FAO:HC'	1:B:359:LEU:O	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:380:ILE:CD1	3:B:500:FAO:H7M1	2.37	0.45
1:B:159:LEU:HD12	1:B:159:LEU:O	2.18	0.44
1:B:185:ALA:O	1:B:209:LYS:HD2	2.17	0.44
1:A:159:LEU:HD12	1:A:159:LEU:C	2.38	0.43
1:A:51:GLU:HG3	4:A:572:HOH:O	2.19	0.43
1:A:146:ALA:HA	1:A:179:ILE:HD12	2.00	0.42
1:B:326[A]:ARG:HE	1:B:331[A]:GLU:CD	2.17	0.42
1:A:269:THR:HA	1:A:272:TYR:CE1	2.55	0.41
1:A:380:ILE:HB	1:B:365:TYR:CZ	2.55	0.41
1:A:170:TYR:CE2	1:A:242[B]:VAL:HG12	2.55	0.41
3:B:500:FAO:H1	3:B:500:FAO:H11'	1.76	0.41
1:A:134:ARG:HB3	1:A:135:PRO:HD3	2.03	0.40
1:A:38[B]:ARG:NH1	4:A:678:HOH:O	2.54	0.40
1:A:56:GLY:HA2	1:A:226:ARG:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/403 (96%)	387 (100%)	1 (0%)	0	100	100
1	B	388/403 (96%)	387 (100%)	1 (0%)	0	100	100
All	All	776/806 (96%)	774 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/303 (97%)	294 (100%)	0	100	100
1	B	292/303 (96%)	290 (99%)	2 (1%)	84	77
All	All	586/606 (97%)	584 (100%)	2 (0%)	91	89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	64	ARG
1	B	294	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	270	GLN
1	A	343	ASN
1	A	374	GLN
1	B	270	GLN
1	B	343	ASN
1	B	374	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	501	-	3,3,3	0.48	0	2,2,2	0.49	0
3	FAO	A	500	-	52,58,58	1.42	6 (11%)	59,89,89	2.04	10 (16%)
2	EDO	B	501	-	3,3,3	0.48	0	2,2,2	0.35	0
3	FAO	B	500	-	52,58,58	1.53	9 (17%)	59,89,89	1.72	10 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	501	-	-	0/1/1/1	-
2	EDO	B	501	-	-	0/1/1/1	-
3	FAO	A	500	-	2/2/14/17	0/30/78/78	0/6/6/6
3	FAO	B	500	-	2/2/14/17	0/30/78/78	0/6/6/6

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	FAO	C10-N1	-5.24	1.40	1.46
3	B	500	FAO	C10-N1	-5.00	1.40	1.46
3	B	500	FAO	C4N-N5	-4.38	1.39	1.45
3	B	500	FAO	C10-N10	-4.36	1.41	1.45
3	A	500	FAO	C10-N10	-4.30	1.41	1.45
3	A	500	FAO	C4N-N5	-3.68	1.40	1.45
3	B	500	FAO	C4N-C4	-2.96	1.47	1.52
3	B	500	FAO	O2-C2	2.58	1.28	1.23
3	B	500	FAO	C2A-N3A	2.47	1.36	1.32
3	A	500	FAO	O4A'-C1A'	2.41	1.44	1.41
3	B	500	FAO	O4A'-C1A'	2.41	1.44	1.41
3	B	500	FAO	C1'-N10	2.34	1.51	1.47
3	A	500	FAO	C4N-C4	-2.22	1.49	1.52
3	B	500	FAO	C2-N1	2.04	1.38	1.34
3	A	500	FAO	C2A-N3A	2.03	1.35	1.32

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	FAO	C4-N3-C2	-6.70	117.28	126.25
3	B	500	FAO	C4-N3-C2	-6.15	118.02	126.25
3	A	500	FAO	O4-C4-N3	-5.92	110.46	121.01
3	A	500	FAO	N3-C2-N1	5.90	122.33	116.12
3	B	500	FAO	N3-C2-N1	5.01	121.39	116.12
3	A	500	FAO	N3A-C2A-N1A	-4.76	121.24	128.68
3	B	500	FAO	N3A-C2A-N1A	-4.61	121.47	128.68
3	A	500	FAO	O2-C2-N3	-4.21	113.87	121.82
3	A	500	FAO	O4-C4-C4N	4.20	128.44	119.86
3	A	500	FAO	C1'-N10-C9N	-3.95	111.40	120.03
3	B	500	FAO	O4-C4-N3	-3.91	114.04	121.01
3	A	500	FAO	C4N-C10-N10	3.02	120.30	110.94
3	B	500	FAO	C1'-N10-C9N	-2.90	113.69	120.03
3	A	500	FAO	O3A'-C3A'-C2A'	-2.83	102.65	111.82
3	B	500	FAO	O2-C2-N3	-2.77	116.60	121.82
3	B	500	FAO	C4N-C10-N10	2.70	119.31	110.94
3	A	500	FAO	P-O3P-PA	-2.40	124.58	132.83
3	B	500	FAO	O4-C4-C4N	2.35	124.66	119.86
3	B	500	FAO	P-O3P-PA	-2.09	125.67	132.83
3	B	500	FAO	O3A'-C3A'-C2A'	-2.07	105.12	111.82

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	500	FAO	C4N
3	A	500	FAO	C10
3	B	500	FAO	C4N
3	B	500	FAO	C10

There are no torsion outliers.


There are no ring outliers.


2 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	500	FAO	6	0
3	B	500	FAO	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	381/403 (94%)	-0.17	16 (4%) 36 40	9, 18, 34, 41	0
1	B	381/403 (94%)	-0.11	14 (3%) 41 46	11, 20, 38, 48	0
All	All	762/806 (94%)	-0.14	30 (3%) 39 44	9, 19, 37, 48	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	199	ALA	7.9
1	B	260	PHE	5.1
1	B	292	PHE	4.8
1	A	292	PHE	4.6
1	B	332	THR	4.2
1	A	260	PHE	4.0
1	B	293	GLY	3.8
1	A	332	THR	3.8
1	A	253	PHE	3.5
1	A	153	GLY	3.3
1	B	158	HIS	3.2
1	B	253	PHE	3.0
1	B	197	PRO	3.0
1	B	199	ALA	3.0
1	B	329	ALA	2.9
1	B	294	ARG	2.6
1	A	403	SER	2.5
1	A	154	SER	2.5
1	B	333	ASN	2.3
1	B	166	ASP	2.3
1	A	329	ALA	2.3
1	A	293	GLY	2.2
1	A	257	ALA	2.2
1	B	403	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	158	HIS	2.1
1	A	290	ASP	2.1
1	A	291	THR	2.1
1	B	167	GLY	2.0
1	A	79	ASP	2.0
1	A	198	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	FAO	A	500	53/53	0.83	0.17	17,24,27,29	53
3	FAO	B	500	53/53	0.87	0.15	18,22,26,26	53
2	EDO	A	501	4/4	0.93	0.10	23,26,28,32	0
2	EDO	B	501	4/4	0.96	0.06	21,25,27,30	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAO A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAO B 500:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.