

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 3, 2023 – 02:47 AM EDT

PDB ID : 6P6Y

Title: Crystal structure of voltage-gated sodium channel NavAb V100C/Q150C

disulfide crosslinked mutant in the activated state

Authors: Wisedchaisri, G.; Tonggu, L.; McCord, E.; Gamal El-din, T.M.; Wang, L.;

Zheng, N.; Catterall, W.A.

Deposited on : 2019-06-04

Resolution : 2.89 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.89 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2201 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Ion transport protein.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	۸	242	Total	С	N	О	S	0	0	0
1	A	242	1977	1334	301	328	14	0	U	

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	983	MET	-	initiating methionine	UNP A8EVM5
A	984	ASP	-	expression tag	UNP A8EVM5
A	985	TYR	-	expression tag	UNP A8EVM5
A	986	LYS	-	expression tag	UNP A8EVM5
A	987	ASP	-	expression tag	UNP A8EVM5
A	988	ASP	-	expression tag	UNP A8EVM5
A	989	ASP	-	expression tag	UNP A8EVM5
A	990	ASP	-	expression tag	UNP A8EVM5
A	991	LYS	-	expression tag	UNP A8EVM5
A	992	GLY	-	expression tag	UNP A8EVM5
A	993	SER	-	expression tag	UNP A8EVM5
A	994	LEU	-	expression tag	UNP A8EVM5
A	995	VAL	-	expression tag	UNP A8EVM5
A	996	PRO	-	expression tag	UNP A8EVM5
A	997	ARG	-	expression tag	UNP A8EVM5
A	998	GLY	-	expression tag	UNP A8EVM5
A	999	SER	-	expression tag	UNP A8EVM5
A	1000	HIS	-	expression tag	UNP A8EVM5
A	1100	CYS	VAL	engineered mutation	UNP A8EVM5
A	1150	CYS	GLN	engineered mutation	UNP A8EVM5


• Molecule 2 is 3-[(3-CHOLAMIDOPROPYL)DIMETHYLAMMONIO]-1-PROPANESULFO NATE (three-letter code: CPS) (formula: C<sub>32</sub>H<sub>58</sub>N<sub>2</sub>O<sub>7</sub>S).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 31				0	0
2	A	1	Total 29	C 24	N 1	O 4	0	0

• Molecule 3 is 1,2-DIMYRISTOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PX4) (formula: C<sub>36</sub>H<sub>73</sub>NO<sub>8</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total 46	C 36	N 1	O 8	P 1	0	0

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
3	Λ	1	Total C N O P	0	0	
3	A	1	31 21 1 8 1	U	0	
3	Λ	1	Total C N O P	0	0	
3	A	1	42 32 1 8 1	U	U	
3	Λ	1	Total C N O P	0	0	
3	A	1	32 23 1 7 1	U	U	
3	Λ	1	Total C	0	0	
$\begin{array}{ c c c c c } \hline 3 & A & \end{array}$		1	13 13	U		

MolProbity and EDS failed to run properly - this section is therefore empty.



# 3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source	
Space group	I 4 2 2	Depositor	
Cell constants	125.83Å 125.83Å 189.29Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	44.53 - 2.89	Depositor	
% Data completeness	97.6 (44.53-2.89)	Depositor	
(in resolution range)	31.0 (44.00-2.03)	Depositor	
$R_{merge}$	0.12	Depositor	
$R_{sym}$	(Not available)	Depositor	
$< I/\sigma(I) > 1$	2.18 (at 2.90Å)	Xtriage	
Refinement program	REFMAC 5.8.0238	Depositor	
$R, R_{free}$	0.214 , 0.273	Depositor	
Wilson B-factor (Å <sup>2</sup> )	76.2	Xtriage	
Anisotropy	0.004	Xtriage	
L-test for twinning <sup>2</sup>	$ < L > = 0.49, < L^2> = 0.33$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	2201	wwPDB-VP	
Average B, all atoms $(\mathring{A}^2)$	82.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 4 Model quality (i)

### 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

## 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 4.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PX4	A	1304	-	30,30,45	0.49	0	36,38,53	0.57	0
3	PX4	A	1307	_	12,12,45	0.26	0	11,11,53	0.22	0
3	PX4	A	1303	-	45,45,45	0.36	0	51,53,53	0.71	2 (3%)
2	CPS	A	1302	-	32,32,45	0.46	0	51,51,70	0.75	0
3	PX4	A	1305	-	41,41,45	0.39	0	47,49,53	0.95	1 (2%)
3	PX4	A	1306	-	31,31,45	0.48	0	36,38,53	0.52	0
2	CPS	A	1301	-	34,34,45	0.54	0	52,53,70	0.86	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PX4	A	1304	-	-	8/34/34/49	-
3	PX4	A	1307	-	-	6/10/10/49	-
3	PX4	A	1303	-	-	21/49/49/49	-
2	CPS	A	1302	-	-	0/9/74/90	0/4/4/4
3	PX4	A	1305	-	-	28/45/45/49	_
3	PX4	A	1306	-	-	17/34/34/49	_
2	CPS	A	1301	-	-	3/12/77/90	0/4/4/4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^{o})$	$\operatorname{Ideal}(^{o})$
3	A	1305	PX4	O7-C23-C24	4.30	120.76	111.50
3	A	1303	PX4	O7-C23-C24	3.36	118.75	111.50
3	A	1303	PX4	O7-C7-C6	2.66	118.03	108.40
2	A	1301	CPS	C6-C5-C4	-2.17	105.38	107.40

There are no chirality outliers.

5 of 83 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
3	A	1303	PX4	C1-O3-P1-O1
3	A	1303	PX4	C1-O3-P1-O2
3	A	1303	PX4	C1-O3-P1-O4
3	A	1304	PX4	O3-C1-C2-N1
3	A	1304	PX4	C7-C6-O4-P1


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















# 4.7 Other polymers (i)

There are no such residues in this entry.

## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 5 Fit of model and data (i)

### 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

## 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

