



## wwPDB EM Validation Summary Report ⓘ

Feb 27, 2022 – 06:13 am GMT

PDB ID : 7PDD  
EMDB ID : EMD-13334  
Title : Focus refinement of soluble domain of Adenylyl cyclase 9 in complex with DARPin C4 and MANT-GTP  
Authors : Qi, C.; Korkhov, V.M.  
Deposited on : 2021-08-05  
Resolution : 4.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev97  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1354	
2	B	147	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylate cyclase 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	449	3550	2240	601	677	32	0	0

- Molecule 2 is a protein called DARPin C4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	128	966	607	171	187	1	0	0



VAL	THR	ASN	ASP	PI038	G1129	E1211	THR	GLU
LEU	LYS	PHE	GLY	Y1039	S1130	E1212	ASP	VAL
SER	ARG	GLN	ILE	H1040	H1131	S1213	GLU	GLY
THR	LEU	LEU	GLU	V1041	F1132	Y1214	ALA	MET
VAL	LEU	SER	ALA	A1042	Q1133	R1215	SER	GLU
PHE	GLU	TRP	SER	E1043	F1140	V1216	LEU	ALA
LEU	TRP	ILE	LEU	Q1044	A1143	L1217	VAL	HIS
ILE	ALA	GLY	GLN	L1045	K1144	M1220	PRO	ASN
ALA	ARG	SER	LEU	V1047	E1145	G1221	SER	GLU
SER	TRP	SER	GLU	S1048	M1146	Y1222	THR	LEU
LEU	LEU	LEU	VAL	Q1049	M1147	F1223	LYS	LYS
ILE	PRO	ALA	ILE	T1050	R1148	F1224	ASN	LEU
THR	ARG	THR	VAL	Y1051	V1149	D1225	PRO	ASN
PHE	HIS	VAL	VAL	S1052	V1150	Y1226	ASP	VAL
ARG	PHE	VAL	PHE	I1061	N1154	R1227	VAL	GLU
ILE	ILE	GLY	PHE	F1062	N1155	G1228	PRO	ARG
TYR	GLY	ALA	LEU	A1063	N1156	T1229	PRO	ALA
GLY	ALA	PRO	LEU	M1067	V1232	R1231	GLY	ALA
ALA	ILE	PRO	LEU	F1071	W1159	V1232	GLU	GLU
ALA	VAL	LEU	LEU	E1077	F1160	G1236	ASN	ASN
ALA	VAL	LEU	VAL	G1078	N1161	Q1237	ALA	ALA
ALA	PRO	LEU	THR	G1079	F1162	M1238	GLN	ALA
ALA	PRO	LEU	PHE	K1080	K1163	K1239	THR	THR
ALA	ALA	VAL	LEU	E1081	L1164	T1240	ARG	ARG
ALA	TYR	VAL	GLU	C1082	R1165	Y1241	ASP	ALA
ALA	SER	PRO	GLU	Y1083	V1166	Y1243	HIS	ALA
ALA	HIS	ASP	VAL	R1084	G1167	P1244	PRO	PRO
PHE	VAL	VAL	VAL	Y1085	A1175	S1249	SER	SER
GLY	THR	SER	TYR	L1086	I1178	G1250	ALA	LYS
ALA	GLU	ALA	LEU	M1087	G1179	VAL	ARG	ARG
LEU	LEU	VAL	ILE	E1088	T1180	PRO	PRO	PRO
LEU	LEU	ILE	HIS	L1089	T1181	GLN	TRP	TRP
LEU	THR	ASN	HIS	I1090	K1182	HIS	LYS	LYS
GLU	THR	ILE	ASP	L1096	L1183	GLN	PRO	PRO
ILE	HIS	ALA	VAL	K1099	L1184	LEU	VAL	VAL
SER	SER	GLN	GLU	P1100	Y1185	SER	ALA	ALA
LEU	THR	THR	ALA	D1101	D1186	ILE	GLU	GLU
VAL	PHE	ASN	ASP	Y1102	I1187	SER	ILE	ILE
SER	THR	PHE	HIS	S1103	W1188	PRO	ARG	ARG
VAL	ARG	THR	SER	S1104	G1189	ASP	CYS	CYS
ARG	SER	THR	ARG	K1107	D1190	ILE	ARG	ARG
MET	ALA	ARG	T1020	I1108	V1192	VAL	PHE	PHE
VAL	VAL	LYS	S1024	T1114	N1193	GLN	GLY	GLY
LEU	THR	LEU	D1027	Y1115	S1196	VAL	ALA	ALA
PHE	THR	CYS	Y1116	M1116	D1199	ASP	ILE	ILE
LEU	VAL	ALA	M1122	A1123	T1200	GLY	GLU	GLU
GLU	VAL	ALA	M1204	R1205	E1204	ILE	LYS	LYS
ASP	VAL	SER	L1032	L1206	G1205	GLY	SER	SER
VAL	GLN	LEU	L1033	I1207	R1206	ARG	ASP	ASP
TYR	PRO	PRO	N1035	D1128	I1207	SER	CYS	CYS
THR	CYS	HIS				PRO	GLU	GLU

• Molecule 2: DARPin C4



MET	H9	H10	G11	S12	D13	L14	G15	K16	K17	E20	R23	A24	G25	Q26	D27	V30	R31	I32	N36	G37	A38	H48	T49	P50	L51	H52	L53	H59	L60	E61	I62	V63	E64	V65	L66	L67	K68	A71	A75	A76	D77	W78	L79	G80	
ARG	GLY	HIS	HIS	HIS	HIS	HIS	HIS	ARG	GLY	SER	ASN	ASN	ALA	GLN	THR	ARG	ALA	HIS	PRO	LEU	ALA	LYS	ARG	TRP	LYS	PRO	VAL	VAL	GLU	ALA	GLU	ARG	CYS	ARG	PHE	LYS	ALA	ILE	GLU	LYS	SER	ASP	CYS	CYS	GLU

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	210729	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.063	Depositor
Minimum map value	-0.034	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.01	Depositor
Map size ( $\text{\AA}$ )	243.0, 243.0, 243.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	0.81, 0.81, 0.81	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/3611	0.48	0/4852
2	B	0.25	0/984	0.40	0/1337
All	All	0.28	0/4595	0.46	0/6189

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3550	0	3486	142	0
2	B	966	0	938	43	0
All	All	4516	0	4424	183	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 183 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1154:ASN:HD21	1:A:1162:PHE:H	1.21	0.88
1:A:392:GLU:N	1:A:493:GLY:O	2.23	0.72

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1206:ARG:HB3	1:A:1242:LEU:HD22	1.72	0.71
1:A:1042:ALA:O	1:A:1046:LYS:HB2	1.90	0.71
1:A:389:GLN:NE2	1:A:494:THR:OG1	2.23	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/1354 (33%)	410 (93%)	31 (7%)	0	100	100
2	B	126/147 (86%)	117 (93%)	9 (7%)	0	100	100
All	All	567/1501 (38%)	527 (93%)	40 (7%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/1176 (33%)	383 (100%)	1 (0%)	92	95
2	B	96/114 (84%)	96 (100%)	0	100	100
All	All	480/1290 (37%)	479 (100%)	1 (0%)	93	96

All (1) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	348	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1154	ASN
2	B	26	GLN
2	B	48	HIS
1	A	1131	HIS
1	A	420	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-13334. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)

#### 6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 134



Y Index: 160




Z Index: 61


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)


### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 51 nm<sup>3</sup>; this corresponds to an approximate mass of 46 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of  $0.238 \text{\AA}^{-1}$

## 8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [\(i\)](#)



\*Reported resolution corresponds to spatial frequency of 0.238 Å<sup>-1</sup>



## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.20	-	-
Author-provided FSC curve	4.07	7.17	4.13
Unmasked-calculated*	-	-	-

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-13334 and PDB model 7PDD. Per-residue inclusion information can be found in section 3 on page 4.

### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Atom inclusion [i](#)



At the recommended contour level, 94% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.