

Full wwPDB X-ray Structure Validation Report (i)

Apr 8, 2024 – 11:04 am BST

PDB ID	:	8PME
Title	:	Structure of Nal1 indica cultivar IR64, construct 88-458
Authors	:	Huang, L.Y.; Rety, S.; Xi, X.G.
Deposited on	:	2023-06-28
Resolution	:	2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motrie	Whole archive	Similar resolution
WIEUTIC	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	$1560 \ (2.16-2.16)$
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	371	79%	16% • •
1	В	371	3% 77%	12% • 10%
1	С	371	2% 79%	14% 8%



8PME

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8165 atoms, of which 2 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	Δ	357	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	Se	0	0	0
1	11		2747	1739	477	520	7	4	0		
1	В	333	Total	С	Ν	Ο	\mathbf{S}	Se	0	0	0
1	I D	000	2557	1614	453	479	7	4	0	0	U
1	С	343	Total	С	Ν	Ο	S	Se	0	0	0
1		545	2632	1665	462	494	7	4		U	0

• Molecule 1 is a protein called Narrow leaf 1.

• Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
0	Λ	1	Total	С	Ν	Ο	Р	0	0	
	A	1	31	10	5	13	3	0	0	
0	D	1	Total	С	Ν	Ο	Р	0	0	
	2 D	1	31	10	5	13	3	0	0	
0	С	1	Total	С	Ν	Ο	Р	0	0	
	2 C	1	31	10	5	13	3	0	0	



• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0

• Molecule 4 is SUCCINIC ACID (three-letter code: SIN) (formula: $C_4H_6O_4$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 4 & 4 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 4 & 4 \end{array}$	0	0
4	В	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{H} & \text{O} \\ 10 & 4 & 2 & 4 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 4 & 4 \end{array}$	0	0

• Molecule 5 is water.

5	А	30	TotalO3030	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	28	TotalO2828	0	0
5	С	41	Total O 41 41	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Narrow leaf 1





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	79.28Å 103.73Å 143.47Å	Deperitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
$\mathbf{P}_{\text{accolution}}(\hat{\lambda})$	39.64 - 2.15	Depositor
Resolution (A)	84.06 - 2.15	EDS
% Data completeness	72.8 (39.64-2.15)	Depositor
(in resolution range)	72.8 (84.06-2.15)	EDS
R _{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.75 (at 2.14 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.18.2_3874+SVN, PHENIX 1.18.2_3874+SVN	Depositor
D D.	0.194 , 0.235	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.194 , 0.234	DCC
R_{free} test set	2321 reflections $(4.88%)$	wwPDB-V
Wilson B-factor $(Å^2)$	50.0	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.35 , 47.3	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8165	wwPDB-V
Average B, all atoms $(Å^2)$	61.0	wwPDB-V

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SIN, ATP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
MOI	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.27	0/2796	0.52	0/3784	
1	В	0.26	0/2599	0.50	0/3509	
1	С	0.26	0/2677	0.49	0/3618	
All	All	0.26	0/8072	0.50	0/10911	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	268	SER	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2747	0	2724	40	2
1	В	2557	0	2551	37	0
1	С	2632	0	2624	35	0
2	А	31	0	12	1	0
2	В	31	0	12	0	0
2	С	31	0	12	0	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
4	А	16	0	8	1	0
4	В	8	2	4	3	0
4	С	8	0	4	0	1
5	А	30	0	0	0	0
5	В	28	0	0	1	0
5	С	41	0	0	1	0
All	All	8163	2	7951	110	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (110)	close	$\operatorname{contacts}$	within	the	same	asymmetric	unit	are	listed	below,	sorted	by	their	clash
magnitud	e.													

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:413:GLY:HA3	1:A:426:TRP:HE1	1.41	0.86
1:A:209:GLU:OE2	1:A:238:ASP:HB3	1.83	0.79
1:B:245:LYS:HB3	1:B:259:TYR:HE1	1.47	0.78
1:B:258:VAL:HG13	1:C:144:TRP:HA	1.67	0.76
1:A:413:GLY:HA3	1:A:426:TRP:NE1	2.05	0.70
1:B:418:THR:HG22	1:B:420:ASP:H	1.57	0.70
1:A:138:ARG:NH1	1:A:169:GLU:OE2	2.24	0.70
1:B:107:ILE:HD12	1:B:107:ILE:H	1.57	0.70
1:B:211:PHE:HB3	4:B:1003:SIN:H21	1.73	0.70
1:B:245:LYS:HG3	1:B:262:ALA:HB2	1.74	0.69
1:B:448:GLU:OE1	1:B:448:GLU:N	2.27	0.68
1:A:95:LEU:O	1:A:99:THR:HG23	1.96	0.65
1:A:395:ASP:HB2	1:A:397:GLU:OE1	1.97	0.63
1:C:282:THR:O	1:C:284:PRO:HD3	2.00	0.62
1:B:245:LYS:HB3	1:B:259:TYR:CE1	2.33	0.62
1:B:172:TYR:CE1	1:B:409:THR:HG22	2.36	0.61
1:A:237:VAL:HG12	1:A:239:LEU:HG	1.82	0.61
1:B:320:LYS:NZ	5:B:1101:HOH:O	2.17	0.59
1:B:211:PHE:HB3	4:B:1003:SIN:C2	2.33	0.58



	lo ao pagom	Interatomic	Clash		
Atom-1	Atom-2	distance (Å)	overlap (Å)		
1:B:245:LYS:HG3	1:B:262:ALA:CB	2.34	0.57		
1:C:169:GLU:HB3	1:C:416:LYS:HB3	1.87	0.57		
1:A:107:ILE:HD12	1:A:107:ILE:H	1.70	0.56		
1:C:355:LEU:HD22	1:C:370:LEU:HD13	1.87	0.56		
1:B:172:TYR:HE1	1:B:409:THR:HG22	1.70	0.56		
1:B:447:ASN:O	1:B:451:GLN:HG3	2.05	0.55		
1:B:243:ASN:HB3	1:B:245:LYS:HD2	1.87	0.55		
1:B:146:ASN:O	1:B:149:GLN:HG2	2.06	0.55		
1:B:135:PHE:HB3	1:B:170:PHE:HB3	1.89	0.55		
1:C:246:MSE:HG3	1:C:295:ILE:HG12	1.88	0.55		
1:C:242:PRO:HB2	1:C:265:ARG:CZ	2.37	0.55		
1:A:211:PHE:HB3	4:A:1003:SIN:H21	1.89	0.54		
1:A:419:SER:OG	1:C:187:GLU:OE2	2.25	0.53		
1:A:240:ASP:O	1:A:243:ASN:N	2.41	0.53		
1:C:103:PHE:O	5:C:1101:HOH:O	2.18	0.53		
1:C:357:TYR:HB3	1:C:365:PHE:HB3	1.90	0.53		
1:C:218:VAL:HG21	1:C:441:LEU:HD13	1.90	0.53		
1:A:233:ARG:CZ	1:A:266:ALA:HB2	2.39	0.52		
1:B:288:VAL:O	1:B:365:PHE:HA	2.10	0.52		
1:C:289:ARG:HH11	1:C:289:ARG:HG2	1.73	0.52		
1:A:204:GLN:OE1	1:A:341:ARG:NE	2.42	0.51		
1:A:214:LEU:HG	1:A:229:PHE:CD1	2.46	0.51		
1:C:274:VAL:O	1:C:278:ILE:HG13	2.11	0.51		
1:A:320:LYS:HB3	1:A:402:ILE:HG22	1.93	0.51		
1:B:414:ARG:HB2	1:B:423:PRO:HB2	1.94	0.50		
1:C:220:ARG:HA	1:C:441:LEU:HD23	1.94	0.49		
1:A:98:MSE:HE1	1:A:324:LEU:HD13	1.93	0.49		
1:C:95:LEU:O	1:C:99:THR:HG23	2.13	0.49		
1:B:341:ARG:HG3	1:B:342:SER:N	2.26	0.49		
1:A:220:ARG:HB2	1:A:225:LYS:HA	1.96	0.48		
1:A:170:PHE:CE1	1:A:415:LEU:HD12	2.48	0.48		
1:B:94:LEU:O	1:B:98:MSE:HG3	2.13	0.48		
1:B:135:PHE:HB3	1:B:170:PHE:CB	2.44	0.48		
1:C:235:VAL:HG22	1:C:235:VAL:O	2.13	0.48		
1:A:341:ARG:HG3	1:A:342:SER:N	2.28	0.48		
1:C:415:LEU:HD22	1:C:417:LEU:HG	1.95	0.48		
1:B:245:LYS:HA	1:B:262:ALA:HA	1.96	0.47		
1:C:341:ARG:HG3	1:C:342:SER:N	2.29	0.47		
1:C:204:GLN:OE1	1:C:341:ARG:NE	2.44	0.47		
1:C:414:ARG:HB3	1:C:423:PRO:HB2	1.97	0.47		
1:C:105:SER:O	1:C:109:ARG:HG3	2.15	0.47		



		Interatomic	Clash		
Atom-1	Atom-2	distance (Å)	overlap (Å)		
1:A:310:VAL:HG12	1:A:313:VAL:HG13	1.96	0.47		
1:C:263:VAL:HA	1:C:295:ILE:HG13	1.97	0.46		
1:C:318:ASP:OD1	1:C:319:VAL:HG12	2.15	0.46		
1:A:415:LEU:HD23	1:A:415:LEU:O	2.15	0.46		
1:C:303:ILE:HG22	1:C:303:ILE:O	2.16	0.46		
1:B:387:SER:HB2	1:B:405:ILE:HD12	1.96	0.46		
1:A:240:ASP:H	1:A:243:ASN:HD22	1.61	0.46		
1:C:415:LEU:CD2	1:C:417:LEU:HG	2.47	0.45		
1:C:318:ASP:OD1	1:C:319:VAL:N	2.50	0.44		
1:A:170:PHE:HE1	1:A:415:LEU:HD12	1.83	0.44		
1:A:244:GLN:O	1:A:262:ALA:HA	2.17	0.44		
1:B:253:ASN:ND2	1:B:254:LEU:HG	2.32	0.44		
1:A:233:ARG:NE	1:A:266:ALA:HB2	2.33	0.44		
1:C:135:PHE:HA	1:C:168:VAL:O	2.18	0.44		
1:A:209:GLU:CD	1:A:243:ASN:HB3	2.39	0.43		
1:A:295:ILE:HD12	1:A:295:ILE:N	2.33	0.43		
1:B:120:ARG:HG3	1:B:166:ASP:OD1	2.18	0.43		
1:A:387:SER:HB2	1:A:405:ILE:HD12	2.00	0.43		
1:B:338:LYS:HD2	1:B:338:LYS:C	2.39	0.43		
1:C:107:ILE:HD12	1:C:107:ILE:N	2.34	0.43		
1:C:228:GLY:HA2	1:C:297:PHE:CD2	2.54	0.43		
1:C:319:VAL:HG23	1:C:432:LEU:HD23	2.01	0.43		
1:B:120:ARG:NH1	1:B:166:ASP:OD1	2.51	0.43		
1:A:373:GLY:HA3	1:A:377:GLN:O	2.18	0.43		
1:C:403:GLY:HA2	1:C:432:LEU:HB2	2.00	0.42		
1:B:124:GLY:HA2	1:B:417:LEU:O	2.19	0.42		
1:A:120:ARG:HG3	1:A:166:ASP:OD1	2.19	0.42		
1:A:199:ILE:O	1:A:306:VAL:HA	2.19	0.42		
1:A:223:GLY:O	1:A:458:ARG:HD2	2.19	0.42		
1:A:338:LYS:HG2	1:A:339:VAL:N	2.29	0.42		
1:A:135:PHE:HB3	1:A:170:PHE:CB	2.50	0.42		
1:A:303:ILE:HG22	1:A:303:ILE:O	2.20	0.42		
1:C:107:ILE:HD12	1:C:107:ILE:H	1.84	0.41		
1:A:145:LEU:HD11	2:A:1001:ATP:C4	2.56	0.41		
1:B:172:TYR:O	1:B:412:ARG:HB3	2.20	0.41		
1:C:170:PHE:HE2	1:C:426:TRP:NE1	2.18	0.41		
1:B:130:PRO:HB3	1:B:328:LEU:CD1	2.50	0.41		
1:B:246:MSE:HE3	1:B:246:MSE:HB3	1.93	0.41		
1:B:107:ILE:HD12	1:B:107:ILE:N	2.31	0.41		
1:B:141:HIS:CE1	1:B:143:LYS:HG3	2.55	0.41		
1:A:189:VAL:HG13	1:A:341:ARG:O	2.21	0.41		



8PME

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268:SER:O	1:A:269:PHE:CD2	2.74	0.41
1:A:336:VAL:HG12	1:A:391:LEU:HD23	2.02	0.40
1:C:376:ARG:HH11	1:C:421:HIS:CG	2.38	0.40
1:C:426:TRP:HZ3	1:C:428:SER:HG	1.68	0.40
1:B:254:LEU:CD1	4:B:1003:SIN:H32	2.52	0.40
1:B:370:LEU:HD11	1:B:426:TRP:HB2	2.04	0.40
1:C:289:ARG:HG2	1:C:289:ARG:NH1	2.36	0.40
1:A:357:TYR:HB3	1:A:365:PHE:HB3	2.03	0.40
1:B:385:SER:HA	1:B:405:ILE:HG22	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:A:259:TYR:OH	1:A:397:GLU:OE2[4_557]	2.05	0.15	
1:A:276:TYR:OH	4:C:1003:SIN:O4[2_455]	2.13	0.07	

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	351/371~(95%)	351 (100%)	0	0	100 100
1	В	323/371~(87%)	323 (100%)	0	0	100 100
1	С	333/371~(90%)	333 (100%)	0	0	100 100
All	All	1007/1113~(90%)	1007 (100%)	0	0	100 100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	299/305~(98%)	295~(99%)	4 (1%)	69 74
1	В	278/305~(91%)	273~(98%)	5(2%)	59 63
1	С	286/305~(94%)	284 (99%)	2 (1%)	84 89
All	All	863/915~(94%)	852 (99%)	11 (1%)	69 74

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	164	ASP
1	А	172	TYR
1	А	338	LYS
1	А	415	LEU
1	В	164	ASP
1	В	245	LYS
1	В	365	PHE
1	В	412	ARG
1	В	440	GLU
1	С	164	ASP
1	С	190	ASP

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	С	283	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tiple	Bo	ond leng	$_{\rm sths}$	B	ond ang	les
WIOI	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
4	SIN	А	1004	1	$7,\!7,\!7$	1.04	0	8,8,8	1.54	1 (12%)
2	ATP	А	1001	3	26,33,33	0.91	1 (3%)	31,52,52	1.62	5 (16%)
4	SIN	А	1003	-	$7,\!7,\!7$	1.02	0	8,8,8	1.69	1 (12%)
2	ATP	В	1001	3	26,33,33	0.92	1 (3%)	31,52,52	1.46	5 (16%)
4	SIN	В	1003	-	7,7,7	1.12	0	8,8,8	1.73	2 (25%)
2	ATP	С	1001	3	26,33,33	0.90	1 (3%)	31,52,52	1.51	6 (19%)
4	SIN	С	1003	-	7,7,7	1.07	0	8,8,8	1.58	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SIN	А	1004	1	-	4/5/5/5	-
2	ATP	А	1001	3	-	3/18/38/38	0/3/3/3
4	SIN	А	1003	-	-	3/5/5/5	-
2	ATP	В	1001	3	-	2/18/38/38	0/3/3/3
4	SIN	В	1003	-	-	4/5/5/5	-
2	ATP	С	1001	3	-	2/18/38/38	0/3/3/3
4	SIN	С	1003	-	-	2/5/5/5	-

All (3) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	С	1001	ATP	C5-C4	2.51	1.47	1.40
2	В	1001	ATP	C5-C4	2.49	1.47	1.40
2	А	1001	ATP	C5-C4	2.48	1.47	1.40

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	С	1001	ATP	PA-O3A-PB	-3.95	119.27	132.83
2	А	1001	ATP	PB-O3B-PG	-3.89	119.47	132.83
2	А	1001	ATP	PA-O3A-PB	-3.78	119.86	132.83
2	В	1001	ATP	PB-O3B-PG	-3.35	121.33	132.83
2	В	1001	ATP	N3-C2-N1	-3.25	123.61	128.68
2	С	1001	ATP	N3-C2-N1	-3.17	123.72	128.68
2	А	1001	ATP	C3'-C2'-C1'	3.14	105.71	100.98
2	А	1001	ATP	N3-C2-N1	-3.10	123.83	128.68
2	В	1001	ATP	PA-O3A-PB	-3.09	122.22	132.83
2	С	1001	ATP	PB-O3B-PG	-3.06	122.34	132.83
2	А	1001	ATP	C4-C5-N7	-2.71	106.58	109.40
2	В	1001	ATP	C4-C5-N7	-2.63	106.66	109.40
2	С	1001	ATP	C3'-C2'-C1'	2.55	104.82	100.98
2	С	1001	ATP	C4-C5-N7	-2.51	106.78	109.40
2	В	1001	ATP	C3'-C2'-C1'	2.51	104.76	100.98
4	В	1003	SIN	O4-C4-C3	2.30	121.41	114.03
4	В	1003	SIN	C3-C2-C1	-2.13	109.03	113.60
4	С	1003	SIN	O2-C1-C2	2.12	120.86	114.03
4	A	1004	SIN	C3-C2-C1	-2.12	109.03	113.60
2	С	1001	ATP	O3G-PG-O2G	2.06	115.51	107.64
4	А	1003	SIN	C2-C3-C4	-2.04	109.22	113.60

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	1003	SIN	C1-C2-C3-C4
4	В	1003	SIN	C1-C2-C3-C4
2	А	1001	ATP	PA-O3A-PB-O2B
2	В	1001	ATP	PA-O3A-PB-O2B
2	С	1001	ATP	PA-O3A-PB-O2B
2	А	1001	ATP	C3'-C4'-C5'-O5'
4	А	1004	SIN	C2-C3-C4-O4
4	А	1004	SIN	O1-C1-C2-C3
4	А	1004	SIN	C2-C3-C4-O3



		<u> </u>	1 0	
Mol	Chain	\mathbf{Res}	Type	Atoms
4	А	1004	SIN	O2-C1-C2-C3
4	С	1003	SIN	O2-C1-C2-C3
2	С	1001	ATP	PA-O3A-PB-O1B
4	А	1003	SIN	O2-C1-C2-C3
4	С	1003	SIN	O1-C1-C2-C3
4	А	1003	SIN	O1-C1-C2-C3
4	В	1003	SIN	O1-C1-C2-C3
4	В	1003	SIN	O2-C1-C2-C3
2	А	1001	ATP	PA-O3A-PB-O1B
2	В	1001	ATP	PA-O3A-PB-O1B
4	В	1003	SIN	C2-C3-C4-O4

Continued from previous page...

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	1001	ATP	1	0
4	А	1003	SIN	1	0
4	В	1003	SIN	3	0
4	С	1003	SIN	0	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	353/371~(95%)	0.32	20 (5%) 23 32	35, 53, 102, 140	0
1	В	329/371~(88%)	0.19	10 (3%) 50 59	40, 60, 103, 127	0
1	С	339/371~(91%)	0.09	8 (2%) 59 67	40, 54, 87, 132	0
All	All	1021/1113~(91%)	0.20	38 (3%) 41 49	35, 56, 98, 140	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	268	SER	8.9
1	А	395	ASP	5.2
1	В	410	ALA	5.0
1	С	265	ARG	4.9
1	А	267	THR	4.9
1	А	414	ARG	4.8
1	А	170	PHE	4.7
1	А	397	GLU	4.7
1	А	269	PHE	4.7
1	А	241	TYR	4.6
1	А	172	TYR	4.5
1	А	270	ILE	4.4
1	В	409	THR	4.4
1	А	171	SER	4.1
1	А	394	GLN	3.7
1	С	266	ALA	3.5
1	С	458	ARG	3.2
1	А	88	GLY	3.1
1	С	242	PRO	3.1
1	В	172	TYR	3.1
1	А	271	THR	3.1
1	В	408	GLY	3.0
1	В	110	ARG	2.9



Mol	Chain	Res	Type	RSRZ
1	В	244	GLN	2.8
1	С	208	HIS	2.8
1	А	239	LEU	2.8
1	С	243	ASN	2.5
1	А	138	ARG	2.5
1	А	209	GLU	2.5
1	В	411	ASN	2.5
1	А	408	GLY	2.4
1	А	183	GLN	2.4
1	С	170	PHE	2.2
1	А	158	PRO	2.2
1	В	88	GLY	2.2
1	В	111	PHE	2.2
1	C	417	LEU	2.1
1	B	234	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	\mathbf{Res}	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SIN	В	1003	8/8	0.58	0.34	103,125,131,133	0
4	SIN	А	1004	8/8	0.75	0.23	75,78,83,87	0
4	SIN	С	1003	8/8	0.76	0.19	56,65,72,77	0
4	SIN	А	1003	8/8	0.81	0.27	60,69,73,74	0
3	MG	В	1002	1/1	0.84	0.13	43,43,43,43	0
3	MG	А	1002	1/1	0.88	0.13	48,48,48,48	0
3	MG	С	1002	1/1	0.91	0.12	43,43,43,43	0
2	ATP	С	1001	31/31	0.97	0.12	43,51,56,59	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	ATP	А	1001	31/31	0.98	0.12	$45,\!53,\!58,\!65$	0
2	ATP	В	1001	31/31	0.98	0.11	44,51,54,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















6.5 Other polymers (i)

There are no such residues in this entry.

